



Full wwPDB X-ray Structure Validation Report i

Aug 10, 2020 – 07:52 AM BST

PDB ID : 4GW4
Title : Crystal structure of 3BNC60 Fab with P61A mutation
Authors : Diskin, R.; Fu, B.Z.; Bjorkman, P.J.
Deposited on : 2012-08-31
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 12825 atoms, of which 6198 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called 3BNC60 Fab Light-chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	L	198	Total	C 3075	H 978	N 1519	O 269	S 305	4	0	0
1	B	199	Total	C 3090	H 982	N 1526	O 271	S 307	4	0	0

- Molecule 2 is a protein called 3BNC60 Fab Heavy-chain.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	H	213	Total	C 3224	H 1043	N 1580	O 286	S 310	5	0	0
2	A	212	Total	C 3209	H 1038	N 1573	O 285	S 308	5	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	L	1	Total C N O 14 8 1 5	0	0
3	B	1	Total C N O 14 8 1 5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	L	55	Total O 55 55	0	0
4	H	36	Total O 36 36	0	0
4	A	42	Total O 42 42	0	0
4	B	66	Total O 66 66	0	0


3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 3BNC60 Fab Light-chain



- Molecule 1: 3BNC60 Fab Light-chain




- Molecule 2: 3BNC60 Fab Heavy-chain



- Molecule 2: 3BNC60 Fab Heavy-chain





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	64.57 Å 154.88 Å 74.24 Å 90.00° 109.75° 90.00°	Depositor
Resolution (Å)	39.35 – 2.65 39.69 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.1 (39.35-2.65) 92.9 (39.69-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.15 (at 2.65 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_966)	Depositor
R , R_{free}	0.213 , 0.256 0.200 , 0.243	Depositor DCC
R_{free} test set	1954 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	48.9	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 76.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12825	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	B	0.56	0/1598	0.70	0/2171
1	L	0.54	0/1590	0.72	0/2160
2	A	0.49	0/1684	0.62	0/2299
2	H	0.49	0/1685	0.64	0/2302
All	All	0.52	0/6557	0.67	0/8932

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1564	1526	1523	66	0
1	L	1556	1519	1517	58	0
2	A	1636	1573	1568	27	0
2	H	1644	1580	1575	32	0
3	B	14	0	13	4	0
3	L	14	0	13	5	0
4	A	42	0	0	2	0
4	B	66	0	0	15	0
4	H	36	0	0	8	0

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:152:PHE:CD1	2:H:153:PRO:N	2.87	0.42
1:L:59:ARG:NH2	1:L:79:GLU:OE2	2.53	0.42
2:A:27:TYR:CE2	2:A:29:ILE:HA	2.54	0.42
1:B:138:LYS:HA	1:B:142:ALA:O	2.20	0.41
1:B:182:ALA:HA	1:B:196:LYS:O	2.20	0.41
2:H:62:ARG:CZ	4:H:322:HOH:O	2.67	0.41
2:H:74(C):ASP:O	2:H:75:PHE:HB2	2.19	0.41
2:H:151:TYR:CZ	2:H:156:VAL:CG1	3.03	0.41
2:A:68:VAL:HG22	2:A:83:LEU:HD23	2.02	0.41
1:L:14:ARG:NH2	1:L:96:LYS:HZ3	2.17	0.41
1:B:95:LEU:HD12	1:B:95:LEU:HA	1.82	0.41
1:B:152:VAL:CG2	1:B:164:LEU:HD12	2.51	0.41
1:B:152:VAL:HG22	1:B:164:LEU:HD12	2.03	0.41
2:A:201:ILE:HG22	2:A:214:ASP:OD1	2.21	0.41
1:B:164:LEU:C	1:B:164:LEU:HD23	2.41	0.41
1:B:4:MET:HA	1:B:24:GLN:O	2.21	0.41
1:L:125:LEU:N	1:L:125:LEU:HD12	2.36	0.41
1:L:172:LYS:HG3	1:L:173:ALA:N	2.36	0.40
1:B:14:ARG:NH2	1:B:96:LYS:HZ3	2.19	0.40
1:B:139:VAL:CG1	1:B:144:GLN:HE22	2.33	0.40
2:H:152:PHE:CD1	2:H:153:PRO:HA	2.56	0.40
2:H:52:ASN:HA	2:H:53:PRO:HD3	1.93	0.40
1:B:125:LEU:HD12	1:B:125:LEU:N	2.37	0.40
1:B:94:ASP:OD1	1:B:94:ASP:C	2.60	0.40
2:H:201:ILE:O	2:H:203:ASN:ND2	2.54	0.40
1:L:173:ALA:HB2	4:L:439:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Continued from previous page...

Mol	Chain	Res	Type
2	H	28	LYS
2	H	60	ASN
2	H	83	LEU
2	H	130	LEU
2	H	155	PRO
2	H	156	VAL
2	H	159	SER
2	H	183	SER
2	H	213	VAL
2	A	2	VAL
2	A	60	ASN
2	A	72	ARG
2	A	83	LEU
2	A	130	LEU
2	A	155	PRO
2	A	156	VAL
2	A	159	SER
2	A	183	SER
2	A	213	VAL
1	B	9	SER
1	B	12	SER
1	B	17	ASP
1	B	19	VAL
1	B	64	ARG
1	B	95	LEU
1	B	174	ASP
1	B	177	LYS
1	B	199	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	L	178	HIS
1	L	187	HIS
2	H	3	HIS
2	H	65	GLN
2	H	203	ASN
2	A	3	HIS
2	A	203	ASN
1	B	187	HIS

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	301	NAG	O7-C7-N2-C2
3	B	301	NAG	C8-C7-N2-C2
3	L	301	NAG	C8-C7-N2-C2
3	L	301	NAG	O7-C7-N2-C2
3	L	301	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	L	301	NAG	5	0
3	B	301	NAG	4	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	B	199/206 (96%)	0.41	17 (8%) 10 8	22, 62, 131, 187	0
1	L	198/206 (96%)	0.68	25 (12%) 3 2	24, 62, 126, 192	0
2	A	212/229 (92%)	0.35	7 (3%) 46 43	22, 62, 128, 143	0
2	H	212/229 (92%)	0.50	17 (8%) 12 9	21, 69, 128, 144	0
All	All	821/870 (94%)	0.48	66 (8%) 12 9	21, 64, 129, 192	0

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	5	THR	6.7
1	L	4	MET	6.2
2	A	195	LEU	6.2
1	L	179	LYS	6.1
2	H	190	VAL	5.5
2	H	193	SER	4.8
1	L	138	LYS	4.6
2	A	190	VAL	4.5
2	H	197	THR	4.4
1	L	67	GLN	4.2
1	L	8	PRO	3.9
1	B	4	MET	3.9
2	H	195	LEU	3.8
1	L	7	SER	3.7
1	B	182	ALA	3.7
2	H	142	ALA	3.7
1	B	8	PRO	3.7
2	A	193	SER	3.7
1	L	140	ASP	3.7
1	B	198	PHE	3.4
1	B	180	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	A	207	LYS	3.3
1	B	140	ASP	3.3
1	L	65	TRP	3.2
1	B	5	THR	3.2
2	H	199	THR	3.2
1	L	192	SER	3.1
1	B	192	SER	3.1
1	B	138	LYS	3.0
1	L	197	SER	3.0
2	A	197	THR	3.0
1	B	65	TRP	2.9
1	B	179	LYS	2.9
1	L	24	GLN	2.9
2	H	207	LYS	2.8
1	L	181	TYR	2.8
1	L	194	VAL	2.8
1	L	176	GLU	2.7
1	L	198	PHE	2.7
1	B	181	TYR	2.6
2	H	191	PRO	2.6
2	H	75	PHE	2.6
1	B	194	VAL	2.6
1	L	190	LEU	2.6
1	B	141	ASN	2.5
1	L	6	GLN	2.5
2	H	76	ASP	2.5
2	H	74(B)	TRP	2.5
2	H	162	SER	2.5
1	L	141	ASN	2.4
1	L	191	SER	2.3
1	B	176	GLU	2.3
1	B	197	SER	2.3
2	H	201	ILE	2.3
1	L	182	ALA	2.3
2	A	188	VAL	2.3
1	L	185	VAL	2.3
1	L	136	GLN	2.3
2	A	210	ASN	2.2
2	H	165	LEU	2.2
2	H	194	SER	2.2
1	L	180	VAL	2.1
1	B	67	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	L	105	PHE	2.1
2	H	74(C)	ASP	2.1
2	H	122	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	H	1	8/9	0.85	0.26	87,111,130,144	0

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	L	301	14/15	0.91	0.22	26,64,79,84	0
3	NAG	B	301	14/15	0.91	0.16	49,79,95,104	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.