



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 6, 2023 – 10:33 PM EDT

PDB ID : 4H1Q
Title : Crystal structure of mutant MMP-9 catalytic domain in complex with a twin inhibitor.
Authors : Stura, E.A.; Vera, L.; Cassar-Lajeunesse, E.; Nuti, E.; Catalani, M.P.; Dive, V.; Rossello, A.
Deposited on : 2012-09-11
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	160	
1	B	160	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	0XX	B	306	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3133 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Matrix metalloproteinase-9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	160	1310	843	226	239	2	1	4	0
1	B	160	1330	854	231	243	2	7	5	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	227	GLN	GLU	engineered mutation	UNP P14780
B	227	GLN	GLU	engineered mutation	UNP P14780


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Ca	0	0
			4	4		
3	B	3	Total	Ca	0	0
			3	3		

- Molecule 4 is N-(4-([(3R)-3-[(biphenyl-4-ylsulfonyl)(propan-2-yloxy)amino]-4-(hydroxyamino)-4-oxobutyl]amino}-4-oxobutyl)-N'-(4-([(3S)-3-[(biphenyl-4-ylsulfonyl)(propan-2-yloxy)amino]-4-(hydroxyamino)-4-oxobutyl]amino}-4-oxobutyl)benzene-1,3-dicarboxamide (three-letter code: OXX) (formula: C₅₄H₆₆N₈O₁₄S₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	A	1	78	54	8	14	2	0	0
4	B	1	78	54	8	14	2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	A	156	156	156	0	1
5	B	170	170	170	0	0



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Matrix metalloproteinase-9



- Molecule 1: Matrix metalloproteinase-9



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	44.14Å 48.66Å 67.92Å 90.00° 102.55° 90.00°	Depositor
Resolution (Å)	43.09 – 1.59 43.09 – 1.59	Depositor EDS
% Data completeness (in resolution range)	96.8 (43.09-1.59) 96.8 (43.09-1.59)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.59Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, R_{free}	0.170 , 0.211 0.171 , 0.211	Depositor DCC
R_{free} test set	1846 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	12.2	Xtrriage
Anisotropy	0.175	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 45.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3133	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, OXX, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/1358	0.55	0/1849
1	B	0.39	0/1377	0.56	0/1874
All	All	0.38	0/2735	0.56	0/3723

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1310	0	1201	10	1
1	B	1330	0	1226	24	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	4	0	0	0	0
3	B	3	0	0	0	0
4	A	78	0	65	12	0
4	B	78	0	65	37	1
5	A	156	0	0	2	0
5	B	170	0	0	2	0
All	All	3133	0	2557	60	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:306:0XX:C46	4:B:306:0XX:H12	1.65	1.26
1:B:235:ASP:OD2	4:B:306:0XX:C67	1.93	1.15
4:B:306:0XX:H12	4:B:306:0XX:H35	1.07	1.03
4:B:306:0XX:H35	4:B:306:0XX:C68	1.91	0.99
4:B:306:0XX:C68	4:B:306:0XX:C47	2.45	0.94
4:B:306:0XX:C46	4:B:306:0XX:C68	2.51	0.87
4:B:306:0XX:C68	4:B:306:0XX:H34	2.06	0.86
4:B:306:0XX:H12	4:B:306:0XX:C47	2.05	0.85
4:A:307:0XX:H52	4:A:307:0XX:H36	1.59	0.84
1:B:235:ASP:HB3	4:B:306:0XX:H41	1.58	0.84
4:B:306:0XX:H34	4:B:306:0XX:C63	2.08	0.83
1:B:235:ASP:HA	4:B:306:0XX:H42	1.58	0.83
4:B:306:0XX:O62	4:B:306:0XX:H31	1.86	0.75
1:B:235:ASP:OD2	4:B:306:0XX:C66	2.35	0.74
5:A:523:HOH:O	4:B:306:0XX:H16	1.86	0.74
1:A:182:ASP:HB3	4:A:307:0XX:H22	1.71	0.73
1:B:235:ASP:HB3	4:B:306:0XX:C40	2.18	0.73
4:B:306:0XX:O62	4:B:306:0XX:C51	2.37	0.72
1:B:249[C]:ARG:NH1	5:B:541:HOH:O	2.25	0.70
1:B:235:ASP:CA	4:B:306:0XX:H42	2.26	0.65
4:A:307:0XX:H47	4:A:307:0XX:H40	1.80	0.64
1:B:235:ASP:OD2	4:B:306:0XX:C68	2.46	0.63
4:B:306:0XX:O62	4:B:306:0XX:C50	2.46	0.63
4:A:307:0XX:H12	4:A:307:0XX:O78	1.99	0.62
4:B:306:0XX:O62	4:B:306:0XX:H29	1.99	0.62
4:B:306:0XX:H34	4:B:306:0XX:S60	2.41	0.60
1:A:120[A]:ASN:OD1	1:A:120[A]:ASN:N	2.29	0.60
1:B:213:GLY:HA3	1:B:250:PHE:CE2	2.38	0.59
1:A:110:PHE:CD1	1:A:110:PHE:N	2.72	0.58
1:B:249[A]:ARG:NH1	1:B:250:PHE:O	2.38	0.57
1:B:191:ALA:O	4:B:306:0XX:H46	2.05	0.55
1:B:235:ASP:HB3	4:B:306:0XX:C41	2.37	0.55
4:B:306:0XX:H31	4:B:306:0XX:S60	2.46	0.54
1:B:187:LEU:HD11	4:B:306:0XX:H55	1.91	0.52
1:B:226:HIS:CD2	4:B:306:0XX:O23	2.62	0.52
1:A:190:HIS:CG	4:A:307:0XX:H37	2.45	0.52
1:B:235:ASP:CB	4:B:306:0XX:H42	2.41	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:HIS:HD1	4:A:307:0XX:H34	1.76	0.50
1:B:233:GLY:O	4:B:306:0XX:O53	2.31	0.49
1:A:258:LYS:HE2	5:A:529:HOH:O	2.14	0.48
4:B:306:0XX:H36	4:B:306:0XX:H11	1.95	0.48
1:B:230:HIS:CD2	4:B:306:0XX:O23	2.68	0.46
1:B:235:ASP:HA	4:B:306:0XX:C41	2.37	0.46
4:A:307:0XX:H29	4:A:307:0XX:H24	1.97	0.45
4:B:306:0XX:C75	4:B:306:0XX:H26	2.47	0.44
1:B:214:LYS:HA	1:B:214:LYS:HD2	1.45	0.43
4:A:307:0XX:O77	1:B:249[B]:ARG:NH1	2.51	0.43
1:B:162:ARG:HB3	1:B:162:ARG:CZ	2.48	0.42
4:B:306:0XX:O53	4:B:306:0XX:C43	2.67	0.42
1:A:148:TRP:O	1:A:152:THR:HG22	2.19	0.42
4:B:306:0XX:H12	4:B:306:0XX:C45	2.44	0.42
1:A:179:TYR:HB3	4:A:307:0XX:H38	2.02	0.42
4:A:307:0XX:H9	4:A:307:0XX:H28	1.74	0.42
4:B:306:0XX:C68	4:B:306:0XX:H33	2.42	0.42
1:B:179:TYR:O	1:B:190:HIS:HE1	2.02	0.41
1:A:226:HIS:CD2	4:A:307:0XX:O23	2.72	0.41
4:A:307:0XX:C67	1:B:241:GLU:HG3	2.50	0.41
1:B:266:HIS:ND1	5:B:454:HOH:O	2.32	0.41
4:B:306:0XX:H44	4:B:306:0XX:H43	1.29	0.41
1:A:246:PRO:HG2	4:B:306:0XX:C23	2.51	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:119[B]:HIS:NE2	4:B:306:0XX:N44[2_656]	1.87	0.33

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	162/160 (101%)	153 (94%)	8 (5%)	1 (1%)	25	8
1	B	164/160 (102%)	158 (96%)	6 (4%)	0	100	100
All	All	326/320 (102%)	311 (95%)	14 (4%)	1 (0%)	41	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	113	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/128 (103%)	126 (96%)	6 (4%)	27	8
1	B	134/128 (105%)	132 (98%)	2 (2%)	65	44
All	All	266/256 (104%)	258 (97%)	8 (3%)	44	16

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	110	PHE
1	A	120[A]	ASN
1	A	120[B]	ASN
1	A	147	LEU
1	A	198	ILE
1	A	252	GLU
1	B	147	LEU
1	B	214	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 11 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	0XX	A	307	2	78,82,82	2.53	18 (23%)	94,112,112	2.46	22 (23%)
4	0XX	B	306	2	78,82,82	2.68	21 (26%)	94,112,112	4.00	41 (43%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	0XX	A	307	2	-	29/82/94/94	0/5/5/5
4	0XX	B	306	2	-	22/82/94/94	0/5/5/5

All (39) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	307	0XX	C54-N55	9.11	1.61	1.48
4	B	306	0XX	C75-N76	9.05	1.45	1.33
4	A	307	0XX	C21-N22	7.87	1.44	1.33
4	A	307	0XX	C75-N76	7.35	1.43	1.33
4	B	306	0XX	C43-N44	7.27	1.49	1.33
4	B	306	0XX	O62-S60	7.20	1.51	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	306	0XX	C21-N22	7.17	1.43	1.33
4	A	307	0XX	C43-N44	6.82	1.48	1.33
4	B	306	0XX	C20-N16	6.48	1.57	1.48
4	A	307	0XX	C51-C54	6.47	1.63	1.53
4	B	306	0XX	C34-N33	6.01	1.47	1.33
4	B	306	0XX	O14-S1	-6.00	1.36	1.43
4	A	307	0XX	C48-N49	5.81	1.46	1.33
4	A	307	0XX	C34-N33	5.45	1.45	1.33
4	B	306	0XX	C48-N49	5.27	1.45	1.33
4	A	307	0XX	C29-N28	5.02	1.44	1.33
4	B	306	0XX	C26-C20	4.84	1.61	1.53
4	B	306	0XX	O23-N22	-4.56	1.28	1.40
4	A	307	0XX	O23-N22	-3.87	1.30	1.40
4	B	306	0XX	O77-N76	-3.69	1.30	1.40
4	B	306	0XX	O15-S1	-3.66	1.39	1.43
4	A	307	0XX	O77-N76	-3.56	1.31	1.40
4	B	306	0XX	C29-N28	3.48	1.41	1.33
4	B	306	0XX	O35-C34	-3.38	1.16	1.23
4	B	306	0XX	C20-C21	3.24	1.60	1.52
4	A	307	0XX	C54-C75	3.18	1.60	1.52
4	B	306	0XX	C8-C5	2.64	1.55	1.49
4	A	307	0XX	C7-C2	2.61	1.42	1.38
4	B	306	0XX	C37-C34	-2.55	1.44	1.50
4	A	307	0XX	C37-C34	2.46	1.55	1.50
4	B	306	0XX	S60-N55	-2.40	1.58	1.70
4	A	307	0XX	C63-S60	2.38	1.79	1.76
4	B	306	0XX	C7-C2	2.35	1.42	1.38
4	B	306	0XX	S1-N16	-2.33	1.59	1.70
4	A	307	0XX	C2-S1	2.25	1.79	1.76
4	A	307	0XX	O15-S1	2.16	1.45	1.43
4	B	306	0XX	C2-S1	2.10	1.79	1.76
4	A	307	0XX	C64-C63	2.07	1.42	1.38
4	A	307	0XX	C26-C20	2.03	1.56	1.53

All (63) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	306	0XX	O61-S60-C63	-17.23	86.24	108.05
4	B	306	0XX	O61-S60-O62	-14.43	96.13	119.52
4	A	307	0XX	O61-S60-O62	-12.56	99.16	119.52
4	A	307	0XX	O15-S1-O14	-11.43	100.99	119.52
4	B	306	0XX	O15-S1-O14	-10.80	102.02	119.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	306	OXX	O62-S60-C63	7.92	118.06	108.05
4	B	306	OXX	O15-S1-C2	-7.57	98.47	108.05
4	B	306	OXX	C37-C38-C39	-7.56	111.63	120.44
4	B	306	OXX	O14-S1-C2	6.98	116.88	108.05
4	B	306	OXX	C68-C63-S60	6.61	126.71	119.76
4	B	306	OXX	C64-C63-S60	-6.47	112.94	119.76
4	B	306	OXX	C32-N33-C34	6.41	136.69	122.08
4	B	306	OXX	C63-S60-N55	6.29	132.20	104.71
4	B	306	OXX	C47-C48-N49	5.99	126.50	116.42
4	B	306	OXX	C31-C32-N33	-5.96	95.18	112.21
4	B	306	OXX	C31-C30-C29	-5.68	97.32	113.26
4	A	307	OXX	C37-C34-N33	5.23	128.31	117.09
4	B	306	OXX	C40-C39-C38	5.21	125.39	119.24
4	B	306	OXX	C3-C2-S1	-5.19	114.29	119.76
4	B	306	OXX	C39-C43-N44	-4.98	106.42	117.09
4	B	306	OXX	C7-C2-S1	4.89	124.90	119.76
4	A	307	OXX	C31-C30-C29	-4.36	101.02	113.26
4	B	306	OXX	C27-N28-C29	-4.13	115.17	122.84
4	A	307	OXX	C67-C66-C69	-3.99	114.45	121.36
4	B	306	OXX	C42-C41-C40	-3.95	114.64	120.25
4	B	306	OXX	C42-C37-C38	3.90	123.85	119.24
4	A	307	OXX	O25-C21-N22	-3.87	118.26	123.27
4	B	306	OXX	C20-N16-S1	-3.87	110.96	118.57
4	B	306	OXX	C42-C37-C34	-3.83	108.21	120.62
4	B	306	OXX	C51-C54-C75	-3.79	103.98	112.45
4	A	307	OXX	C20-N16-S1	-3.70	111.29	118.57
4	A	307	OXX	C32-N33-C34	3.64	130.38	122.08
4	B	306	OXX	C40-C39-C43	-3.61	108.92	120.62
4	B	306	OXX	C45-C46-C47	3.49	123.20	112.84
4	A	307	OXX	O35-C34-N33	-3.39	115.85	122.61
4	B	306	OXX	O53-C48-C47	-3.39	115.82	122.02
4	B	306	OXX	C54-N55-S60	-3.29	112.10	118.57
4	A	307	OXX	C54-N55-S60	3.24	124.94	118.57
4	A	307	OXX	O36-C29-N28	-3.18	117.01	123.01
4	A	307	OXX	C64-C63-S60	3.12	123.05	119.76
4	B	306	OXX	O35-C34-N33	3.04	128.66	122.61
4	B	306	OXX	O52-C43-C39	3.00	126.29	120.94
4	A	307	OXX	O78-C75-N76	-2.99	119.40	123.27
4	A	307	OXX	C2-S1-N16	2.91	117.42	104.71
4	A	307	OXX	O35-C34-C37	-2.87	115.82	120.94
4	A	307	OXX	C27-N28-C29	-2.84	117.57	122.84
4	B	306	OXX	O53-C48-N49	-2.83	117.68	123.01

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	306	0XX	O35-C34-C37	-2.72	116.08	120.94
4	A	307	0XX	C9-C8-C5	-2.54	116.96	121.36
4	B	306	0XX	C51-C50-N49	2.51	119.11	111.87
4	B	306	0XX	C30-C29-N28	2.44	120.53	116.42
4	A	307	0XX	C63-S60-N55	2.43	115.31	104.71
4	A	307	0XX	O15-S1-C2	-2.42	104.98	108.05
4	B	306	0XX	C3-C4-C5	-2.36	117.74	121.13
4	B	306	0XX	O52-C43-N44	2.35	127.29	122.61
4	A	307	0XX	C65-C66-C69	2.34	125.41	121.36
4	B	306	0XX	C41-C42-C37	2.34	123.11	120.34
4	A	307	0XX	C30-C29-N28	2.31	120.31	116.42
4	B	306	0XX	C38-C37-C34	2.25	127.87	120.44
4	B	306	0XX	C50-N49-C48	2.16	126.86	122.84
4	B	306	0XX	C70-C69-C66	-2.07	117.78	121.36
4	B	306	0XX	C2-S1-N16	2.04	113.65	104.71
4	A	307	0XX	C26-C27-N28	-2.00	106.09	111.87

There are no chirality outliers.

All (51) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	307	0XX	C50-C51-C54-C75
4	A	307	0XX	O56-N55-S60-O61
4	A	307	0XX	C68-C63-S60-N55
4	A	307	0XX	C64-C63-S60-N55
4	A	307	0XX	C45-C46-C47-C48
4	A	307	0XX	C20-C26-C27-N28
4	A	307	0XX	C21-C20-C26-C27
4	B	306	0XX	C51-C54-N55-S60
4	B	306	0XX	O56-N55-S60-O62
4	B	306	0XX	O56-N55-S60-O61
4	B	306	0XX	C68-C63-S60-N55
4	B	306	0XX	C64-C63-S60-N55
4	B	306	0XX	C58-C57-O56-N55
4	B	306	0XX	C21-C20-C26-C27
4	A	307	0XX	C38-C39-C43-O52
4	A	307	0XX	C40-C39-C43-O52
4	A	307	0XX	C38-C39-C43-N44
4	A	307	0XX	C40-C39-C43-N44
4	A	307	0XX	O35-C34-N33-C32
4	B	306	0XX	C47-C48-N49-C50
4	A	307	0XX	C37-C34-N33-C32

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	307	0XX	C65-C66-C69-C70
4	B	306	0XX	C65-C66-C69-C74
4	A	307	0XX	C67-C66-C69-C70
4	A	307	0XX	C65-C66-C69-C74
4	B	306	0XX	C67-C66-C69-C74
4	B	306	0XX	C67-C66-C69-C70
4	B	306	0XX	C65-C66-C69-C70
4	A	307	0XX	C67-C66-C69-C74
4	B	306	0XX	O53-C48-N49-C50
4	B	306	0XX	C38-C39-C43-O52
4	B	306	0XX	C38-C39-C43-N44
4	B	306	0XX	N44-C45-C46-C47
4	B	306	0XX	C40-C39-C43-O52
4	B	306	0XX	C40-C39-C43-N44
4	A	307	0XX	C64-C63-S60-O61
4	A	307	0XX	C68-C63-S60-O61
4	A	307	0XX	C51-C50-N49-C48
4	B	306	0XX	C30-C31-C32-N33
4	A	307	0XX	C26-C20-N16-S1
4	B	306	0XX	C54-N55-S60-O61
4	A	307	0XX	O35-C34-C37-C38
4	A	307	0XX	C51-C54-C75-O78
4	A	307	0XX	C51-C54-C75-N76
4	A	307	0XX	O35-C34-C37-C42
4	B	306	0XX	C64-C63-S60-O61
4	A	307	0XX	N33-C34-C37-C38
4	B	306	0XX	C68-C63-S60-O61
4	A	307	0XX	N33-C34-C37-C42
4	A	307	0XX	C50-C51-C54-N55
4	A	307	0XX	C7-C2-S1-O15


There are no ring outliers.

2 monomers are involved in 50 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	307	0XX	12	0
4	B	306	0XX	37	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is

within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	160/160 (100%)	-0.06	7 (4%) 34 31	7, 13, 33, 61	1 (0%)
1	B	160/160 (100%)	-0.16	5 (3%) 49 46	8, 14, 33, 49	0
All	All	320/320 (100%)	-0.11	12 (3%) 40 37	7, 13, 33, 61	1 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	110	PHE	9.4
1	A	111	GLU	5.7
1	B	110	PHE	4.8
1	A	215	GLY	4.2
1	B	212	LEU	3.6
1	B	213	GLY	3.4
1	A	112	GLY	3.0
1	B	214	LYS	2.5
1	A	160	TYR	2.5
1	B	249[A]	ARG	2.4
1	A	249[A]	ARG	2.1
1	A	250	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	OXX	A	307	78/78	0.86	0.23	10,39,67,92	0
4	OXX	B	306	78/78	0.89	0.21	8,37,60,84	0
3	CA	A	305	1/1	0.98	0.04	14,14,14,14	0
3	CA	B	304	1/1	0.99	0.04	14,14,14,14	0
3	CA	A	304	1/1	0.99	0.09	23,23,23,23	0
3	CA	A	306	1/1	0.99	0.12	29,29,29,29	0
2	ZN	A	302	1/1	1.00	0.03	15,15,15,15	0
2	ZN	B	301	1/1	1.00	0.04	10,10,10,10	0
3	CA	B	303	1/1	1.00	0.05	9,9,9,9	0
2	ZN	B	302	1/1	1.00	0.04	13,13,13,13	0
3	CA	B	305	1/1	1.00	0.02	11,11,11,11	0
3	CA	A	303	1/1	1.00	0.06	10,10,10,10	0
2	ZN	A	301	1/1	1.00	0.04	11,11,11,11	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 0XX A 307:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 0XX B 306:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.