



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 25, 2020 – 04:00 PM BST

PDB ID : 5H3Z
Title : Crystal Structure of 1,2-beta-oligoglucan phosphorylase from *Lachnoclostridium phytofermentans*
Authors : Nakajima, M.; Tanaka, N.; Furukawa, N.; Nihira, T.; Kodutsumi, Y.; Takahashi, Y.; Sugimoto, N.; Miyanaga, A.; Fushinobu, S.; Taguchi, H.; Nakai, H.
Deposited on : 2016-10-28
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1122	 % 92% 6% ..
1	B	1122	 % 92% 7% ..

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1203	-	X	-	-
4	PEG	A	1207	-	-	X	-
4	PEG	B	1209	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 18799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uncharacterized protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	1113	8918	5712	1483	1691	14	18	0	0	0
1	B	1113	8926	5717	1484	1692	14	19	0	1	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MSE	-	expression tag	UNP A9KJS6
A	1	GLY	-	expression tag	UNP A9KJS6
A	1114	LEU	-	expression tag	UNP A9KJS6
A	1115	GLU	-	expression tag	UNP A9KJS6
A	1116	HIS	-	expression tag	UNP A9KJS6
A	1117	HIS	-	expression tag	UNP A9KJS6
A	1118	HIS	-	expression tag	UNP A9KJS6
A	1119	HIS	-	expression tag	UNP A9KJS6
A	1120	HIS	-	expression tag	UNP A9KJS6
A	1121	HIS	-	expression tag	UNP A9KJS6
B	0	MSE	-	expression tag	UNP A9KJS6
B	1	GLY	-	expression tag	UNP A9KJS6
B	1114	LEU	-	expression tag	UNP A9KJS6
B	1115	GLU	-	expression tag	UNP A9KJS6
B	1116	HIS	-	expression tag	UNP A9KJS6
B	1117	HIS	-	expression tag	UNP A9KJS6
B	1118	HIS	-	expression tag	UNP A9KJS6
B	1119	HIS	-	expression tag	UNP A9KJS6
B	1120	HIS	-	expression tag	UNP A9KJS6
B	1121	HIS	-	expression tag	UNP A9KJS6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Ca 2 2	0	0
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	B	1	Total C O 7 4 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	432	Total O 432 432	0	0
5	B	439	Total O 439 439	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Uncharacterized protein



- Molecule 1: Uncharacterized protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	87.76Å 94.78Å 157.94Å 90.00° 98.41° 90.00°	Depositor
Resolution (Å)	42.72 – 2.00 42.72 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.6 (42.72-2.00) 99.6 (42.72-2.00)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.64 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.168 , 0.204 0.177 , 0.211	Depositor DCC
R_{free} test set	8658 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	27.4	Xtrriage
Anisotropy	0.099	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	18799	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CA, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	1/9125 (0.0%)	0.86	11/12366 (0.1%)
1	B	0.92	3/9133 (0.0%)	0.87	14/12376 (0.1%)
All	All	0.92	4/18258 (0.0%)	0.86	25/24742 (0.1%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	540	GLN	CD-OE1	7.12	1.39	1.24
1	B	994	ASP	CB-CG	5.58	1.63	1.51
1	B	116	CYS	CB-SG	-5.49	1.72	1.81
1	A	311	GLU	CG-CD	5.29	1.59	1.51

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	182	ARG	NE-CZ-NH1	7.24	123.92	120.30
1	A	812	ARG	NE-CZ-NH1	7.08	123.84	120.30
1	A	936	MSE	CA-CB-CG	-6.74	101.85	113.30
1	B	812	ARG	NE-CZ-NH1	6.54	123.57	120.30
1	B	573	ARG	NE-CZ-NH2	6.26	123.43	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8918	0	8675	40	0
1	B	8926	0	8683	47	0
2	A	24	0	31	3	0
2	B	36	0	48	2	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
4	A	14	0	20	4	0
4	B	7	0	10	8	0
5	A	432	0	0	13	0
5	B	439	0	0	8	0
All	All	18799	0	17467	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 94 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1063:MSE:HE2	1:A:1112:LYS:HB3	1.44	0.98
1:B:516:LEU:HD11	1:B:534:LEU:HD11	1.56	0.87
1:B:540:GLN:HG3	5:B:1519:HOH:O	1.78	0.83
1:A:165:MSE:HE3	5:A:1633:HOH:O	1.80	0.80
2:A:1204:GOL:H2	5:A:1307:HOH:O	1.90	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1111/1122 (99%)	1073 (97%)	37 (3%)	1 (0%)	51 49

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	1112/1122 (99%)	1073 (96%)	38 (3%)	1 (0%)	51	49
All	All	2223/2244 (99%)	2146 (96%)	75 (3%)	2 (0%)	51	49

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	906	LEU
1	B	906	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	975/965 (101%)	946 (97%)	29 (3%)	41	41
1	B	976/965 (101%)	955 (98%)	21 (2%)	52	55
All	All	1951/1930 (101%)	1901 (97%)	50 (3%)	46	48

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	847	GLU
1	A	1106	ASN
1	B	1014	ARG
1	A	875	LEU
1	A	1014	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	525	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 3 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	1201	-	5,5,5	0.98	0	5,5,5	1.15	0
4	PEG	A	1206	-	6,6,6	0.61	0	5,5,5	0.48	0
4	PEG	A	1207	-	6,6,6	0.91	0	5,5,5	1.22	0
2	GOL	A	1202	-	5,5,5	1.00	0	5,5,5	0.86	0
2	GOL	B	1202	-	5,5,5	0.67	0	5,5,5	0.53	0
2	GOL	B	1206	-	5,5,5	0.72	0	5,5,5	0.38	0
4	PEG	B	1209	-	6,6,6	1.05	0	5,5,5	0.85	0
2	GOL	B	1201	-	5,5,5	0.64	0	5,5,5	0.54	0
2	GOL	A	1203	-	5,5,5	0.66	0	5,5,5	1.57	2 (40%)
2	GOL	B	1205	-	5,5,5	0.27	0	5,5,5	1.34	0
2	GOL	A	1204	-	5,5,5	1.34	1 (20%)	5,5,5	0.79	0
2	GOL	B	1203	-	5,5,5	0.32	0	5,5,5	1.32	1 (20%)
2	GOL	B	1204	-	5,5,5	0.58	0	5,5,5	1.34	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1201	-	-	4/4/4/4	-
4	PEG	A	1206	-	-	2/4/4/4	-
4	PEG	A	1207	-	-	1/4/4/4	-
2	GOL	A	1202	-	-	4/4/4/4	-
2	GOL	B	1202	-	-	0/4/4/4	-
2	GOL	B	1206	-	-	0/4/4/4	-
4	PEG	B	1209	-	-	1/4/4/4	-
2	GOL	B	1201	-	-	0/4/4/4	-
2	GOL	A	1203	-	-	4/4/4/4	-
2	GOL	B	1205	-	-	4/4/4/4	-
2	GOL	A	1204	-	-	2/4/4/4	-
2	GOL	B	1203	-	-	2/4/4/4	-
2	GOL	B	1204	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1204	GOL	O3-C3	2.35	1.52	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1204	GOL	O2-C2-C1	2.43	119.82	109.12
2	A	1203	GOL	C3-C2-C1	-2.35	102.58	111.70
2	B	1203	GOL	O2-C2-C1	2.15	118.60	109.12
2	A	1203	GOL	O2-C2-C1	2.08	118.28	109.12

There are no chirality outliers.

5 of 26 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1201	GOL	O1-C1-C2-O2
2	A	1201	GOL	O1-C1-C2-C3
2	A	1202	GOL	O1-C1-C2-C3
2	A	1203	GOL	O1-C1-C2-C3
2	A	1203	GOL	C1-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 17 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1207	PEG	4	0
2	A	1202	GOL	1	0
4	B	1209	PEG	8	0
2	A	1203	GOL	1	0
2	A	1204	GOL	1	0
2	B	1204	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1095/1122 (97%)	-0.30	11 (1%) 82 81	17, 29, 49, 70	0
1	B	1095/1122 (97%)	-0.31	11 (1%) 82 81	18, 29, 47, 70	0
All	All	2190/2244 (97%)	-0.30	22 (1%) 82 81	17, 29, 48, 70	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	836	HIS	3.6
1	A	711	LEU	3.2
1	B	259	GLN	3.0
1	A	798	HIS	2.7
1	A	1064	ASP	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PEG	B	1209	7/7	0.85	0.15	33,37,42,43	0
2	GOL	A	1204	6/6	0.85	0.14	35,38,41,42	0
2	GOL	B	1205	6/6	0.86	0.18	46,53,59,60	0
2	GOL	B	1204	6/6	0.86	0.17	36,48,50,51	0
4	PEG	A	1206	7/7	0.88	0.19	38,48,55,61	0
2	GOL	A	1202	6/6	0.89	0.17	33,39,46,53	0
4	PEG	A	1207	7/7	0.89	0.16	35,37,40,43	0
2	GOL	B	1202	6/6	0.91	0.15	31,43,45,47	0
2	GOL	A	1203	6/6	0.91	0.15	36,43,46,48	0
2	GOL	A	1201	6/6	0.93	0.14	27,30,33,33	0
2	GOL	B	1203	6/6	0.94	0.16	29,36,40,42	0
2	GOL	B	1206	6/6	0.95	0.12	31,33,36,39	0
3	CA	B	1207	1/1	0.98	0.18	59,59,59,59	0
3	CA	A	1205	1/1	0.98	0.05	39,39,39,39	0
2	GOL	B	1201	6/6	0.98	0.14	24,26,27,28	0
3	CA	B	1208	1/1	0.99	0.07	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.