

Full wwPDB X-ray Structure Validation Report (i)

May 20, 2024 – 11:46 PM EDT

PDB ID : 7H9J

Title: Group deposition for crystallographic fragment screening of Chikungunya

virus nsP3 macrodomain – Crystal structure of Chikungunya virus nsP3

macrodomain in complex with Z57376600 (CHIKV MacB-x1454)

Authors: Aschenbrenner, J.C.; Fairhead, M.; Godoy, A.S.; Balcomb, B.H.; Capkin,

E.; Chandran, A.V.; Dolci, I.; Golding, M.; Koekemoer, L.; Lithgo, R.M.; Marples, P.G.; Ni, X.; Oliva, G.; Thompson, W.; Tomlinson, C.W.E.; Wild,

C.; Winokan, M.; Xavier, M.-A.E.; Fearon, D.; von Delft, F.

Deposited on : 2024-04-26

Resolution : 1.88 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36.2

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$


CCP4 : 7.0.044 (Gargrove)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.88 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(\mathring{A}))$
R_{free}	130704	9470 (1.90-1.86)
Clashscore	141614	10282 (1.90-1.86)
Ramachandran outliers	138981	10152 (1.90-1.86)
Sidechain outliers	138945	10152 (1.90-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain	
1	A	163	91%	9%
1	В	163	85%	15%
1	С	163	91%	9%
1	D	163	89%	9% •

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.36.2



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 5633 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Non-structural protein 3.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	٨	163	Total	С	N	О	S	0	2	0
1	A	105	1262	787	220	245	10	0	2	U
1	В	163	Total	С	N	О	S	0	2	0
1	Ъ	105	1263	787	220	247	9	0	2	U
1	С	163	Total	С	N	О	S	0	3	0
1		105	1274	794	225	246	9	0	3	U
1	D	159	Total	С	N	О	S	0	3	0
1	ע	159	1249	778	220	243	8	0	3	U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q8JUX6
A	-1	ALA	-	expression tag	UNP Q8JUX6
A	0	MET	-	expression tag	UNP Q8JUX6
A	77	THR	SER	conflict	UNP Q8JUX6
В	-2	GLY	-	expression tag	UNP Q8JUX6
В	-1	ALA	-	expression tag	UNP Q8JUX6
В	0	MET	-	expression tag	UNP Q8JUX6
В	77	THR	SER	conflict	UNP Q8JUX6
С	-2	GLY	-	expression tag	UNP Q8JUX6
С	-1	ALA	-	expression tag	UNP Q8JUX6
С	0	MET	-	expression tag	UNP Q8JUX6
С	77	THR	SER	conflict	UNP Q8JUX6
D	-2	GLY	-	expression tag	UNP Q8JUX6
D	-1	ALA	-	expression tag	UNP Q8JUX6
D	0	MET	-	expression tag	UNP Q8JUX6
D	77	THR	SER	conflict	UNP Q8JUX6

• Molecule 2 is DIMETHYL SULFOXIDE (three-letter code: DMS) (formula: C₂H₆OS).






Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf
2	٨	1	Total	С	О	S	0	0
2	A	1	4	2	1	1	0	0
2	A	1	Total	С	О	S	0	0
2	A	1	4	2	1	1	0	U
2	A	1	Total	С	О	S	0	0
	11	1	4	2	1	1	0	U
2	A	1	Total	С	О	S	0	0
	71	1	4	2	1	1		U
2	A	1	Total	С	О	S	0	0
	11	1	4	2	1	1	Ü	0
2	A	1	Total	С	О	S	0	0
_		_	4	2	1	1		Ů
2	В	1	Total	С	0	S	0	0
			4	2	1	1		
2	В	1	Total	С	0	S	0	0
			4	2	1	1		
2	В	1	Total	С	0	S	0	0
			4	2	1	1		
2	В	1	Total	С	0	S	0	0
			4	$\frac{2}{C}$	1	$\frac{1}{S}$		
2	С	1	Total		0		0	0
			4	$\frac{2}{C}$	1 O	$\frac{1}{S}$		
2	С	1	Total	2	1		0	0
			4 Total	$\frac{Z}{C}$	$\frac{1}{O}$	$\frac{1}{S}$		
2		1	4	2	1	3 1	0	0
			Total	$\frac{2}{C}$	O	$\frac{1}{S}$		
2	D	1	10tai 4	2	1	3 1	0	0
			4		т		mtime and are m	



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total C O S 4 2 1 1	0	0
2	D	1	Total C O S 4 2 1 1	0	0
2	D	1	Total C O S 4 2 1 1	0	0

• Molecule 3 is 2-AMINO-2-HYDROXYMETHYL-PROPANE-1,3-DIOL (three-letter code: TRS) (formula: $C_4H_{12}NO_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 8 4 1 3	0	0
3	D	1	Total C N O 8 4 1 3	0	0

 \bullet Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	3	Total Cl 3 3	0	0
4	В	2	Total Cl 2 2	0	0
4	С	3	Total Cl 3 3	0	0
4	D	2	Total Cl 2 2	0	0



• Molecule 5 is 5-bromo-6-methylpyridin-2-amine (three-letter code: 4JO) (formula: $C_6H_7BrN_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	Λ	1	Total	Br	С	N	0	0
3	А	1	9	1	6	2	0	0
5	B	1	Total	Br	С	N	0	0
)	Ъ	1	9	1	6	2	0	0
5	С	1	Total	Br	С	N	0	0
9	C	1	9	1	6	2	0	0
5	D	1	Total	Br	С	N	0	0
	D	1	9	1	6	2	U	U

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	140	Total O 141 141	0	1
6	В	101	Total O 101 101	0	0
6	С	118	Total O 118 118	0	0
6	D	95	Total O 95 95	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Non-structural protein 3





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 31	Depositor
Cell constants	87.55Å 87.55Å 85.70Å	D t
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.01 - 1.88	Depositor
Resolution (A)	38.98 - 1.88	EDS
% Data completeness	94.8 (39.01-1.88)	Depositor
(in resolution range)	94.8 (38.98-1.88)	EDS
R_{merge}	0.33	Depositor
R_{sum}	(Not available)	Depositor
$< I/\sigma(I) > 1$	4.78 (at 1.88Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC5	Depositor
υ .	0.241 , 0.309	Depositor
R, R_{free}	0.287 , 0.321	DCC
R_{free} test set	2869 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	15.4	Xtriage
Anisotropy	0.570	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.24 , 49.3	EDS
L-test for twinning ²	$< L >=0.52, < L^2>=0.35$	Xtriage
	0.000 for -h,-k,l	
Estimated twinning fraction	0.040 for h,-h-k,-l	Xtriage
	0.003 for -k,-h,-l	
F_o, F_c correlation	0.89	EDS
Total number of atoms	5633	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 24.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.7427e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, DMS, TRS, 4JO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.75	0/1285	0.88	0/1739	
1	В	0.71	0/1286	0.89	0/1741	
1	С	0.71	0/1297	0.81	0/1754	
1	D	0.71	0/1272	0.84	0/1721	
All	All	0.72	0/5140	0.86	0/6955	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	1262	0	1256	11	0
1	В	1263	0	1253	16	0
1	С	1274	0	1272	10	0
1	D	1249	0	1240	8	0
2	A	24	0	36	5	0
2	В	16	0	24	0	0
2	С	12	0	18	0	0
2	D	16	0	24	0	0
3	A	8	0	12	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	8	0	12	0	0
4	A	3	0	0	0	0
4	В	2	0	0	0	0
4	С	3	0	0	0	0
4	D	2	0	0	0	0
5	A	9	0	7	1	0
5	В	9	0	7	1	0
5	С	9	0	7	1	0
5	D	9	0	7	1	0
6	A	141	0	0	1	0
6	В	101	0	0	3	0
6	С	118	0	0	3	0
6	D	95	0	0	1	0
All	All	5633	0	5175	45	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 4.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:91[A]:GLU:OE2	6:B:301:HOH:O	1.93	0.86
1:C:80:GLU:OE1	6:C:301:HOH:O	2.11	0.67
1:C:144:ARG:HG3	5:C:207:4JO:BR	2.51	0.66
1:B:121:LEU:HD12	1:B:148:TRP:CD2	2.31	0.65
1:A:144:ARG:HG2	5:A:210:4JO:BR	2.58	0.58
1:C:26:ARG:NH2	6:C:308:HOH:O	2.39	0.55
1:A:121:LEU:C	1:A:121:LEU:HD23	2.27	0.55
1:D:144[A]:ARG:HG2	5:D:208:4JO:BR	2.64	0.53
1:D:140:VAL:HG12	1:D:142:TYR:CE1	2.45	0.51
1:A:113:VAL:CG2	2:A:202:DMS:C2	2.88	0.51
1:B:7:LYS:HE2	1:B:140:VAL:HG11	1.94	0.50
1:B:144:ARG:HG2	5:B:207:4JO:C5	2.40	0.50
1:C:22:ALA:HB3	1:C:33:VAL:CG2	2.42	0.50
1:D:121:LEU:C	1:D:121:LEU:HD23	2.32	0.50
1:B:47:ASN:O	1:B:56:LYS:HE2	2.12	0.48
1:B:55:ALA:HA	1:B:67:HIS:O	2.15	0.47
1:D:108:LEU:HB3	1:D:111:THR:HG21	1.97	0.47
1:A:114:TYR:CZ	2:A:202:DMS:H12	2.50	0.47
1:D:8:ARG:HA	1:D:143:CYS:O	2.15	0.47
1:D:31:ASP:HA	1:D:35:LYS:HB2	1.96	0.46



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${ m distance}({ m \AA})$	overlap (Å)
1:B:121:LEU:HD12	1:B:148:TRP:CG	2.50	0.46
1:B:5:ARG:O	1:B:140:VAL:HA	2.15	0.46
1:B:103:SER:HA	1:B:138:ASP:O	2.15	0.46
1:B:76:TYR:HE2	6:B:338:HOH:O	1.98	0.46
1:D:3[A]:SER:OG	6:D:301:HOH:O	2.21	0.46
1:D:16:GLU:HG3	1:D:103:SER:HB2	1.98	0.46
1:B:52:VAL:HA	1:B:69:VAL:HG12	1.97	0.46
1:C:126:ASN:ND2	6:C:302:HOH:O	2.19	0.45
1:A:103:SER:HA	1:A:138:ASP:O	2.17	0.45
1:C:22:ALA:HB3	1:C:33:VAL:HG21	1.98	0.44
1:B:18:CYS:SG	1:B:104:VAL:HG23	2.58	0.44
1:B:128:LEU:C	1:B:128:LEU:HD23	2.38	0.44
1:A:57:THR:O	2:A:205:DMS:H13	2.18	0.44
1:B:11:ILE:HG21	1:B:33:VAL:HG13	2.00	0.43
1:A:55:ALA:HA	1:A:67:HIS:O	2.18	0.43
1:A:8:ARG:HA	1:A:143:CYS:O	2.19	0.43
1:A:113:VAL:CG2	2:A:202:DMS:H21	2.48	0.43
1:C:121:LEU:C	1:C:121:LEU:HD23	2.40	0.42
1:B:3:SER:HA	6:B:353:HOH:O	2.19	0.42
1:A:157:GLN:OE1	2:A:206:DMS:C2	2.67	0.42
1:B:121:LEU:C	1:B:121:LEU:HD23	2.40	0.42
1:C:106[B]:ILE:O	1:C:141:ILE:HA	2.19	0.42
1:A:126:ASN:HB3	6:A:409:HOH:O	2.22	0.40
1:C:26:ARG:HB3	1:C:28:LEU:HG	2.04	0.40
1:C:128:LEU:C	1:C:128:LEU:HD23	2.41	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/163 (100%)	161 (99%)	2 (1%)	0	100 100	7



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	В	163/163 (100%)	160 (98%)	3 (2%)	0	100	100
1	С	164/163 (101%)	160 (98%)	4 (2%)	0	100	100
1	D	160/163 (98%)	158 (99%)	2 (1%)	0	100	100
All	All	650/652 (100%)	639 (98%)	11 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	136/134 (102%)	134 (98%)	2 (2%)	65 59		
1	В	136/134~(102%)	134 (98%)	2 (2%)	65 59		
1	С	137/134 (102%)	133 (97%)	4 (3%)	42 32		
1	D	136/134 (102%)	135 (99%)	1 (1%)	84 83		
All	All	$545/536 \ (102\%)$	536 (98%)	9 (2%)	60 54		

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LYS
1	A	159	ARG
1	В	0	MET
1	В	15	ASP
1	С	0	MET
1	С	31	ASP
1	С	35	LYS
1	С	159	ARG
1	D	145	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:



Mol	Chain	Res	Type
1	С	157	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 33 ligands modelled in this entry, 10 are monoatomic - leaving 23 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Res	Link	В	ond leng	gths	В	ond ang	gles
Mol	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DMS	В	201	-	3,3,3	0.22	0	3,3,3	0.13	0
5	4JO	A	210	-	8,9,9	0.18	0	7,12,12	0.43	0
2	DMS	D	203	-	3,3,3	0.26	0	3,3,3	0.14	0
2	DMS	A	206	-	3,3,3	0.35	0	3,3,3	0.05	0
2	DMS	В	202	-	3,3,3	0.22	0	3,3,3	0.09	0
2	DMS	A	202	-	3,3,3	0.12	0	3,3,3	0.14	0
2	DMS	D	205	-	3,3,3	0.30	0	3,3,3	0.05	0
2	DMS	С	201	-	3,3,3	0.25	0	3,3,3	0.08	0
5	4JO	D	208	-	8,9,9	0.19	0	7,12,12	0.38	0
5	4JO	С	207	-	8,9,9	0.20	0	7,12,12	0.36	0
3	TRS	A	204	-	7,7,7	0.25	0	9,9,9	0.34	0
2	DMS	В	203	-	3,3,3	0.19	0	3,3,3	0.12	0
3	TRS	D	202	-	7,7,7	0.18	0	9,9,9	0.23	0
2	DMS	A	203	-	3,3,3	0.28	0	3,3,3	0.14	0



Mol	ol Type Chain Res Link		Tiple	В	ond leng	gths	Bond angles			
MIOI	туре	Chain	nes Lilik		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	DMS	D	204	-	3,3,3	0.21	0	3,3,3	0.12	0
2	DMS	D	201	-	3,3,3	0.22	0	3,3,3	0.12	0
2	DMS	A	205	-	3,3,3	0.32	0	3,3,3	0.15	0
5	4JO	В	207	-	8,9,9	0.20	0	7,12,12	0.39	0
2	DMS	С	205	-	3,3,3	0.27	0	3,3,3	0.03	0
2	DMS	С	206	-	3,3,3	0.27	0	3,3,3	0.05	0
2	DMS	A	201	-	3,3,3	0.35	0	3,3,3	0.15	0
2	DMS	В	204	-	3,3,3	0.24	0	3,3,3	0.10	0
2	DMS	A	211	-	3,3,3	0.27	0	3,3,3	0.12	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	4JO	D	208	-	-	-	0/1/1/1
5	4JO	С	207	-	-	-	0/1/1/1
5	4JO	A	210	_	-	-	0/1/1/1
3	TRS	A	204	-	-	2/9/9/9	-
3	TRS	D	202	-	-	4/9/9/9	-
5	4JO	В	207	-	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	202	TRS	N-C-C3-O3
3	D	202	TRS	C3-C-C1-O1
3	D	202	TRS	N-C-C1-O1
3	A	204	TRS	C2-C-C3-O3
3	D	202	TRS	C2-C-C1-O1
3	A	204	TRS	N-C-C3-O3


There are no ring outliers.

7 monomers are involved in 9 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	210	4JO	1	0
2	A	206	DMS	1	0
2	A	202	DMS	3	0
5	D	208	4JO	1	0
5	С	207	4JO	1	0
2	A	205	DMS	1	0
5	В	207	4JO	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

