

Full wwPDB X-ray Structure Validation Report (i)

Aug 20, 2020 – 05:50 PM BST

PDB ID : 4HAY

Title: Crystal structure of CRM1 inhibitor Leptomycin B in complex with CRM1(K

548E,K579Q)-Ran-RanBP1

Authors : Sun, Q.; Chook, Y.M.

Deposited on : 2012-09-27

Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.13.1 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	216	86%	6% 7%
2	В	140	79%	6% • 14%
3	С	1023	90%	9% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	EDO	С	1103	-	-	X	-
6	EDO	С	1104	-	-	X	-



2 Entry composition (i)

There are 9 unique types of molecules in this entry. The entry contains 11886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GTP-binding nuclear protein Ran.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	200	Total 1674	C 1079	N 289	O 300	S 6	0	8	0

• Molecule 2 is a protein called Ran-specific GTPase-activating protein 1.

AltConf	Trace
2	0
	2

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	98	LYS	ALA	CONFLICT	UNP P41920

• Molecule 3 is a protein called Exportin-1.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
3	С	1014	Total 8474	C 5460	N 1380	O 1591	S 43	0	48	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	-1	GLY	_	EXPRESSION TAG	UNP P30822
С	0	ALA	-	EXPRESSION TAG	UNP P30822
С	539	CYS	THR	ENGINEERED MUTATION	UNP P30822
С	548	GLU	LYS	ENGINEERED MUTATION	UNP P30822
С	579	GLN	LYS	ENGINEERED MUTATION	UNP P30822
С	1022	CYS	TYR	ENGINEERED MUTATION	UNP P30822

• Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-GUANYLATE ESTER (three-letter code: GNP) (formula: C₁₀H₁₇N₆O₁₃P₃).






Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
4	A	1	Total 32	C 10	N 6	O 13	P 3	0	0

• Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total Mg 1 1	0	0


 \bullet Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $\mathrm{C_2H_6O_2}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	С	1	Total C O 4 2 2	0	0
6	С	1	Total C O 4 2 2	0	0
6	С	1	Total C O 4 2 2	0	0
6	С	1	Total C O 4 2 2	0	0

 \bullet Molecule 7 is Leptomycin B, bound form (three-letter code: LMB) (formula: $\mathrm{C_{33}H_{52}O_{7}}).$



Mol	Chain	Residues	Atom	5	ZeroOcc	AltConf
7	С	1	Total C 40 33	O 7	0	0

 \bullet Molecule 8 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

\mathbf{N}	Iol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
	8	С	3	Total Cl 3 3	0	0

• Molecule 9 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	102	Total O 102 102	0	0
9	В	27	Total O 27 27	0	0
9	С	499	Total O 499 499	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GTP-binding nuclear protein Ran





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	106.17Å 106.17Å 305.66Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 - 2.30	Depositor
Resolution (A)	47.40 - 2.29	EDS
% Data completeness	98.0 (50.00-2.30)	Depositor
(in resolution range)	97.9 (47.40-2.29)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.59 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
D D.	0.187 , 0.230	Depositor
R, R_{free}	0.191 , 0.232	DCC
R_{free} test set	3900 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	41.7	Xtriage
Anisotropy	0.094	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.30 , 34.2	EDS
L-test for twinning ²	$ < L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11886	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: LMB, GNP, EDO, MG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Clasies	Bo	nd lengths	Bond angles	
Mol Chain		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.78	1/1722~(0.1%)	0.78	0/2327
2	В	0.69	1/1031~(0.1%)	0.75	0/1375
3	С	0.76	$1/8742 \ (0.0\%)$	0.75	0/11837
All	All	0.76	3/11495~(0.0%)	0.75	0/15539

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	В	103	TRP	CD2-CE2	5.26	1.47	1.41
3	С	502	TRP	CD2-CE2	5.09	1.47	1.41
1	A	163	TRP	CD2-CE2	5.08	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1674	0	1681	15	0
2	В	1014	0	1010	10	0
3	С	8474	0	8630	72	0
4	A	32	0	13	0	0
5	A	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	4	0	6	0	0
6	С	16	0	24	11	0
7	С	40	0	48	3	0
8	С	3	0	0	0	0
9	A	102	0	0	0	0
9	В	27	0	0	1	0
9	С	499	0	0	15	0
All	All	11886	0	11412	93	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (93) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:917[B]:ARG:HH11	3:C:917[B]:ARG:HG3	1.07	1.06
3:C:917[B]:ARG:NH1	3:C:917[B]:ARG:HG3	1.83	0.91
3:C:306:GLY:O	6:C:1104:EDO:H21	1.71	0.90
1:A:179:MET:HE2	2:B:133:LYS:HA	1.56	0.87
3:C:688:THR:HA	3:C:691:LEU:HD23	1.59	0.82
3:C:1019[B]:THR:HG21	6:C:1103:EDO:H21	1.71	0.72
3:C:229:ILE:HG13	3:C:262:LEU:HD11	1.76	0.68
3:C:977:GLN:O	3:C:979:ALA:N	2.30	0.65
7:C:1101:LMB:O6	7:C:1101:LMB:C33	2.47	0.62
3:C:134:TRP:CD1	3:C:137:PHE:HB2	2.35	0.62
3:C:1019[B]:THR:HG21	6:C:1103:EDO:H11	1.82	0.62
3:C:1052:ALA:HB2	9:C:1649:HOH:O	2.01	0.61
3:C:244:THR:HG22	3:C:285:GLN:OE1	2.01	0.61
3:C:1019[B]:THR:HG21	6:C:1103:EDO:C2	2.31	0.60
3:C:896:ASN:ND2	3:C:947:ASP:O	2.33	0.59
3:C:1052:ALA:CB	9:C:1649:HOH:O	2.51	0.57
3:C:917[B]:ARG:CG	3:C:917[B]:ARG:HH11	1.92	0.57
3:C:1019[B]:THR:CG2	6:C:1103:EDO:H21	2.35	0.56
3:C:690:LEU:HD22	3:C:738:MET:SD	2.46	0.56
1:A:179:MET:HE2	2:B:133:LYS:CA	2.33	0.55
3:C:427:LEU:HD23	3:C:475:LEU:HD11	1.89	0.54
3:C:674:MET:O	3:C:678:THR:HG23	2.07	0.54
3:C:703[B]:ILE:HD13	3:C:766:ILE:HG13	1.91	0.53
3:C:883[B]:LYS:HD3	9:C:1498:HOH:O	2.09	0.53
3:C:138:ILE:HB	3:C:139:PRO:HD3	1.89	0.53
3:C:690:LEU:CD2	3:C:738:MET:SD	2.98	0.52



Continued from previous page...

Continued from previous		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance} \ ({\rm \AA})$	$\text{overlap } (\text{\AA})$	
3:C:462[B]:ILE:HD12	9:C:1421:HOH:O	2.08	0.52	
3:C:701:ASN:HB3	9:C:1663:HOH:O	2.10	0.52	
3:C:1019[B]:THR:HG21	6:C:1103:EDO:C1	2.40	0.51	
3:C:473:VAL:HG13	3:C:518:THR:HG22	1.93	0.51	
1:A:81:ILE:HD11	3:C:65:PHE:CG	2.46	0.51	
3:C:500:TRP:CZ3	3:C:538:LEU:HD11	2.46	0.50	
3:C:192[B]:GLN:HG3	9:C:1592:HOH:O	2.11	0.50	
1:A:87:ILE:HG12	1:A:118[B]:VAL:CG2	2.42	0.50	
3:C:306:GLY:O	6:C:1104:EDO:C2	2.50	0.50	
3:C:525:LYS:O	3:C:529:VAL:HG23	2.12	0.50	
3:C:917[B]:ARG:CG	3:C:917[B]:ARG:NH1	2.58	0.50	
1:A:117:ILE:HB	1:A:144:LEU:HD22	1.93	0.50	
3:C:793:ASN:HB3	9:C:1585:HOH:O	2.12	0.49	
1:A:159:LYS:HB2	1:A:160:PRO:HD3	1.94	0.49	
3:C:366:TRP:O	3:C:370:VAL:HG22	2.13	0.48	
3:C:997:ASN:HA	9:C:1402:HOH:O	2.13	0.48	
3:C:791:LEU:O	3:C:795:VAL:HG22	2.14	0.48	
3:C:755[B]:LYS:HA	3:C:755[B]:LYS:NZ	2.29	0.47	
3:C:981:VAL:HG13	3:C:982:PRO:CD	2.44	0.47	
3:C:141:LEU:HD23	3:C:141:LEU:C	2.34	0.47	
3:C:293[A]:VAL:HG23	9:C:1319:HOH:O	2.15	0.47	
3:C:307:ASN:HA	6:C:1104:EDO:H22	1.98	0.46	
3:C:981:VAL:HG13	3:C:982:PRO:HD2	1.96	0.46	
1:A:179:MET:CE	2:B:134:ILE:H	2.28	0.46	
3:C:142:ILE:CD1	3:C:192[B]:GLN:HB3	2.44	0.46	
7:C:1101:LMB:O6	7:C:1101:LMB:H4	2.15	0.46	
1:A:209:LEU:HD23	2:B:103:TRP:CG	2.51	0.46	
3:C:264:ILE:HG22	3:C:264:ILE:O	2.16	0.46	
2:B:131:THR:O	2:B:132:LEU:HB2	2.16	0.45	
3:C:48:TRP:N	9:C:1247:HOH:O	2.45	0.45	
3:C:1021:GLN:CD	3:C:1027:VAL:HG12	2.37	0.45	
3:C:309:GLN:HB2	6:C:1104:EDO:H11	1.99	0.45	
3:C:690:LEU:HD13	3:C:738:MET:HE1	1.99	0.45	
2:B:80:GLU:N	9:B:327:HOH:O	2.50	0.45	
3:C:238:LEU:O	3:C:252:THR:HG21	2.17	0.45	
3:C:284[A]:GLN:NE2	3:C:288:THR:HG21	2.32	0.45	
2:B:106:ARG:NH2	2:B:166:GLU:HG2	2.32	0.44	
3:C:789[B]:GLU:HB2	3:C:790:PRO:HD3	1.99	0.44	
3:C:485:GLU:HB3	9:C:1467:HOH:O	2.17	0.44	
3:C:235:LEU:HA	3:C:235:LEU:HD23	1.89	0.44	
3:C:50:LYS:NZ	9:C:1605:HOH:O	2.50	0.44	



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	A tom-2	${f distance} \; ({f \AA})$	overlap (Å)
3:C:570:TRP:CE2	3:C:614:GLN:HG3	2.53	0.44
3:C:82:LEU:O	3:C:87:ARG:NH1	2.51	0.44
3:C:691:LEU:HD13	3:C:691:LEU:HA	1.92	0.43
1:A:13:LEU:HD23	1:A:13:LEU:C	2.38	0.43
3:C:307:ASN:CA	6:C:1104:EDO:H22	2.48	0.43
3:C:660[A]:GLU:OE1	3:C:660[A]:GLU:N	2.52	0.43
3:C:328:ALA:HB3	9:C:1593:HOH:O	2.17	0.43
1:A:118[B]:VAL:HG21	1:A:160:PRO:HB3	2.01	0.43
3:C:446[A]:ASN:C	3:C:446[A]:ASN:OD1	2.57	0.43
1:A:87:ILE:HG23	1:A:118[B]:VAL:HG23	2.01	0.43
3:C:107:VAL:HG13	3:C:111[B]:GLN:OE1	2.20	0.42
3:C:607:LYS:HG3	3:C:654:GLU:CG	2.50	0.42
3:C:173:GLN:NE2	9:C:1511:HOH:O	2.52	0.42
3:C:505:ILE:HG23	3:C:506:ASN:N	2.34	0.42
3:C:555:ILE:O	3:C:559:VAL:HG22	2.20	0.42
3:C:582:GLU:HG2	9:C:1326:HOH:O	2.19	0.42
1:A:209:LEU:HD23	2:B:103:TRP:CD1	2.55	0.42
3:C:293[A]:VAL:HG11	3:C:345:GLN:HB3	2.02	0.42
3:C:696:VAL:HG11	3:C:755[B]:LYS:HZ1	1.85	0.42
1:A:118[B]:VAL:O	1:A:118[B]:VAL:HG23	2.20	0.41
2:B:130:LYS:HA	2:B:130:LYS:HD2	1.90	0.41
1:A:204:ALA:HB1	2:B:103:TRP:CZ2	2.56	0.41
3:C:506:ASN:HB3	3:C:510:TRP:CZ2	2.55	0.41
1:A:167:LYS:NZ	6:C:1104:EDO:H12	2.35	0.41
3:C:679:ILE:HD13	3:C:699:ILE:HG13	2.03	0.41
7:C:1101:LMB:H8	7:C:1101:LMB:H3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percen	$_{ m tiles}$
1	A	$204/216 \ (94\%)$	196 (96%)	8 (4%)	0	100	100
2	В	121/140 (86%)	113 (93%)	8 (7%)	0	100	100
3	С	$1054/1023 \; (103\%)$	1039 (99%)	15 (1%)	0	100	100
All	All	1379/1379 (100%)	1348 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	182/185~(98%)	179 (98%)	3 (2%)	62 78
2	В	$106/122 \; (87\%)$	103 (97%)	3 (3%)	43 60
3	С	973/933 (104%)	960 (99%)	13 (1%)	69 82
All	All	1261/1240 (102%)	1242 (98%)	19 (2%)	67 79

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Α	179	MET
1	A	198	GLU
1	A	202	GLU
2	В	130	LYS
2	В	143	GLU
2	В	157	VAL
3	С	207	SER
3	С	247[A]	ASP
3	С	247[B]	ASP
3	С	278	PHE
3	С	291	MET
3	С	538	LEU
3	С	579	GLN
3	С	607	LYS
3	С	611	VAL



Continued from previous page...

Mol	Chain	Res	Type
3	С	690	LEU
3	С	755[A]	LYS
3	С	755[B]	LYS
3	С	983	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	48	HIS
3	С	199	GLN
3	С	203	GLN
3	С	494	GLN
3	С	579	GLN
3	С	813	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Tuno	Chain	in Res 1		Во	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	Res Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	GNP	A	301	5	28,34,34	2.74	7 (25%)	30,54,54	1.22	4 (13%)	
6	EDO	С	1102	-	3,3,3	0.72	0	2,2,2	0.16	0	
6	EDO	A	303	-	3,3,3	0.50	0	2,2,2	0.34	0	
6	EDO	С	1105	_	3,3,3	0.64	0	2,2,2	0.10	0	
6	EDO	С	1104	-	3,3,3	0.62	0	2,2,2	0.72	0	
7	LMB	С	1101	3	31,39,39	1.08	3 (9%)	32,51,51	2.32	9 (28%)	
6	EDO	С	1103	-	3,3,3	0.60	0	2,2,2	0.12	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GNP	A	301	5	-	3/17/38/38	0/3/3/3
6	EDO	С	1102	-	-	1/1/1/1	-
6	EDO	A	303	-	-	1/1/1/1	-
6	EDO	С	1105	-	-	0/1/1/1	-
6	EDO	С	1104	_	-	1/1/1/1	-
7	LMB	С	1101	3	-	8/50/54/54	1
6	EDO	С	1103	-	-	1/1/1/1	=

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	$\operatorname{Ideal}(ext{\AA})$
4	A	301	GNP	C4-N9	-12.09	1.31	1.47
4	A	301	GNP	C5-C6	-3.25	1.47	1.52
4	A	301	GNP	PG-N3B	3.18	1.71	1.63
4	A	301	GNP	PB-N3B	2.97	1.71	1.63
4	A	301	GNP	C8-N9	-2.83	1.35	1.45
4	A	301	GNP	C2-N1	-2.65	1.33	1.44
7	С	1101	LMB	C30-C16	-2.65	1.50	1.54
7	С	1101	LMB	C23-C22	2.30	1.36	1.33
7	С	1101	LMB	C16-C17	-2.22	1.50	1.53
4	A	301	GNP	PB-O3A	2.00	1.61	1.59

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
7	С	1101	LMB	C26-C8-C9	-6.25	117.95	126.39
7	С	1101	LMB	C11-C12-C13	-4.98	114.86	126.00



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
7	С	1101	LMB	C29-C14-C13	4.88	125.77	118.08
7	С	1101	LMB	C2-C3-C4	-4.09	109.21	114.72
4	A	301	GNP	PA-O3A-PB	-3.09	121.75	132.62
7	С	1101	LMB	C19-C18-C17	3.06	115.89	110.36
7	С	1101	LMB	C18-C19-C20	-3.03	109.69	114.73
4	Α	301	GNP	C4-C5-N7	3.02	106.47	102.46
4	Α	301	GNP	O1G-PG-N3B	-2.50	108.09	111.77
4	A	301	GNP	O2G-PG-O1G	-2.45	107.29	113.45
7	С	1101	LMB	C30-C16-C15	-2.33	108.57	110.75
7	С	1101	LMB	C5-C6-C7	-2.19	121.59	125.42
7	С	1101	LMB	C32-C20-C21	2.18	112.99	109.94

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	301	GNP	PG-N3B-PB-O1B
4	A	301	GNP	PG-N3B-PB-O3A
4	A	301	GNP	C2'-C1'-N9-C4
7	С	1101	LMB	C12-C13-C14-C15
7	С	1101	LMB	C27-C26-C8-C9
7	С	1101	LMB	C2-C3-C4-C5
7	С	1101	LMB	C2-C3-C4-C25
6	С	1103	EDO	O1-C1-C2-O2
7	С	1101	LMB	C6-C7-C8-C9
6	С	1104	EDO	O1-C1-C2-O2
7	С	1101	LMB	C12-C13-C14-C29
6	С	1102	EDO	O1-C1-C2-O2
6	A	303	EDO	O1-C1-C2-O2
7	С	1101	LMB	O2-C5-C6-C7
7	С	1101	LMB	C32-C20-C21-C22

There are no ring outliers.


3 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	С	1104	EDO	6	0
7	С	1101	LMB	3	0
6	С	1103	EDO	5	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,



bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	$200/216 \; (92\%)$	-0.04	4 (2%) 65 71	36, 48, 84, 129	0
2	В	121/140 (86%)	-0.27	0 100 100	53, 65, 88, 107	0
3	С	1014/1023 (99%)	-0.26	10 (0%) 82 86	34, 51, 81, 111	0
All	All	1335/1379 (96%)	-0.23	14 (1%) 82 86	34, 52, 82, 129	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	216	LEU	6.7
3	С	262	LEU	5.6
3	С	264	ILE	4.0
3	С	229	ILE	3.5
1	A	196	GLN	3.2
3	С	690	LEU	3.1
1	A	182	LEU	3.0
3	С	455	PHE	2.8
3	С	967	TYR	2.8
3	С	983	GLN	2.5
1	A	179	MET	2.4
3	С	984	GLY	2.2
3	С	542	LYS	2.1
3	С	448	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
6	EDO	С	1105	4/4	0.78	0.14	74,78,80,81	0
6	EDO	С	1103	4/4	0.86	0.51	80,81,85,86	0
6	EDO	С	1104	4/4	0.87	0.15	57,57,57,59	0
7	LMB	С	1101	40/40	0.90	0.22	51,66,98,109	0
6	EDO	С	1102	4/4	0.92	0.14	53,64,65,69	0
6	EDO	A	303	4/4	0.93	0.28	67,70,72,76	0
8	CL	С	1108	1/1	0.94	0.37	81,81,81,81	0
8	CL	С	1107	1/1	0.95	0.10	66,66,66,66	0
8	CL	С	1106	1/1	0.96	0.10	72,72,72,72	0
4	GNP	A	301	32/32	0.99	0.12	40,42,45,47	0
5	MG	A	302	1/1	0.99	0.12	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around LMB C 1101: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)

Electron density around GNP A 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)



6.5 Other polymers (i)

There are no such residues in this entry.

