



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 28, 2021 – 07:23 pm BST

PDB ID : 6HAF
Title : Pyruvate oxidase variant E59Q from *L. plantarum* in complex with phosphate
Authors : Funk, L.M.; Sautner, V.; Tittmann, K.
Deposited on : 2018-08-07
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.22
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.22


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	603	 94%
1	B	603	 95%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	GOL	B	709[B]	-	-	X	-

2 Entry composition i

There are 8 unique types of molecules in this entry. The entry contains 21366 atoms, of which 9847 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyruvate oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	585	9705	3064	4861	848	915	17	0	68	0
1	B	585	9751	3077	4882	855	920	17	0	72	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	59	GLN	GLU	engineered mutation	UNP A0A1A0DLW4
B	59	GLN	GLU	engineered mutation	UNP A0A1A0DLW4

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	53	27	9	15	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		


- Molecule 3 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	K	0	0
			1	1		
3	B	1	Total	K	0	0
			1	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		
4	B	1	Total	Mg	0	0
			1	1		

- Molecule 5 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C₁₂H₁₉N₄O₇P₂S) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
5	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
5	B	1	26	12	4	7	2	1	0	0

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
6	A	1	14	3	8	3	0	0
6	A	1	14	3	8	3	0	0
6	A	1	14	3	8	3	0	0
6	A	1	14	3	8	3	0	1
6	A	1	14	3	8	3	0	1
6	A	1	14	3	8	3	0	0
6	A	1	14	3	8	3	0	1
6	B	1	14	3	8	3	0	0
6	B	1	14	3	8	3	0	0
6	B	1	14	3	8	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	H	O	0	0
			14	3	8	3		
6	B	1	Total	C	H	O	0	1
			28	6	16	6		

- Molecule 7 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			5	4	1		
7	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	754	Total	O	0	43
			782	782		
8	B	742	Total	O	0	59
			774	774		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pyruvate oxidase



- Molecule 1: Pyruvate oxidase



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	119.18Å 154.35Å 165.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.36 – 1.30 48.36 – 1.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.36-1.30) 99.8 (48.36-1.30)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.30Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.134 , 0.156 0.135 , 0.157	Depositor DCC
R_{free} test set	18478 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	10.4	Xtrriage
Anisotropy	0.421	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.98	EDS
Total number of atoms	21366	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, TPP, FAD, K, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.53	0/5178	0.68	2/7037 (0.0%)
1	B	0.52	0/5233	0.69	0/7108
All	All	0.52	0/10411	0.68	2/14145 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	447	ASP	CB-CG-OD1	6.23	123.91	118.30
1	A	447	ASP	CB-CG-OD2	-5.39	113.45	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4844	4861	4634	13	1
1	B	4869	4882	4628	19	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	B	1	0	0	0	0
5	A	26	0	16	0	0
5	B	26	0	16	1	0
6	A	42	56	56	2	0
6	B	36	48	48	11	0
7	A	5	0	0	0	1
7	B	5	0	0	0	0
8	A	782	0	0	6	2
8	B	774	0	0	6	2
All	All	11519	9847	9460	34	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:48[A]:ARG:NH1	8:A:801:HOH:O	2.14	0.80
6:B:709[B]:GOL:H32	8:B:821[B]:HOH:O	1.82	0.79
1:A:225:LYS:NZ	8:A:802:HOH:O	2.25	0.68
1:B:27:ASP:H	6:B:709[B]:GOL:C1	2.07	0.67
1:B:489[B]:GLN:NE2	8:B:802:HOH:O	2.28	0.66
1:A:439:ARG:HH22	6:A:709[A]:GOL:H2	1.60	0.66
1:B:28:HIS:H	6:B:709[B]:GOL:H12	1.63	0.63
1:A:224:ARG:HB2	6:A:710:GOL:H11	1.82	0.62
1:A:498[B]:ASN:ND2	8:A:811:HOH:O	2.37	0.56
1:B:27:ASP:H	6:B:709[B]:GOL:H11	1.71	0.56
1:B:27:ASP:HB3	6:B:709[B]:GOL:H2	1.88	0.55
1:A:224:ARG:NE	8:A:806[A]:HOH:O	2.31	0.52
1:B:27:ASP:HB3	6:B:709[B]:GOL:H12	1.91	0.52
1:A:111[B]:PHE:CE2	1:A:115:GLY:HA3	2.46	0.51
1:B:75[A]:LYS:HD2	6:B:709[A]:GOL:H31	1.95	0.49
5:B:704:TPP:H2	8:B:885:HOH:O	2.14	0.48
1:B:28:HIS:N	6:B:709[B]:GOL:H12	2.28	0.47
1:B:27:ASP:N	6:B:709[B]:GOL:C1	2.77	0.47
1:B:328[B]:LYS:NZ	8:B:814:HOH:O	2.47	0.45
1:A:50[B]:ARG:NH2	8:A:824:HOH:O	2.49	0.44
1:A:39:SER:HB3	1:A:174:TRP:CD2	2.54	0.43
1:A:69:ASP:HB3	8:A:1163:HOH:O	2.18	0.42
1:B:39:SER:HB3	1:B:174:TRP:CD2	2.54	0.42
1:B:553:GLU:OE2	1:B:554[A]:LYS:HE2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283[B]:VAL:HG23	1:B:305:ILE:HB	2.02	0.41
1:A:554:LYS:HE3	1:A:556[A]:ARG:NH2	2.36	0.41
1:B:27:ASP:N	6:B:709[B]:GOL:H12	2.35	0.41
1:B:28:HIS:H	6:B:709[B]:GOL:C1	2.32	0.41
1:B:489[B]:GLN:HG2	8:B:805:HOH:O	2.20	0.41
1:B:232[B]:GLN:HG3	8:B:1136:HOH:O	2.19	0.41
1:B:263[B]:ASN:HB2	1:B:287:TYR:OH	2.21	0.41
1:B:553:GLU:HG2	1:B:554[A]:LYS:HG3	2.03	0.41
1:A:204:ARG:NH1	1:A:320[B]:ILE:CD1	2.84	0.40
1:A:263[B]:ASN:HB2	1:A:287:TYR:OH	2.21	0.40

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:B:933:HOH:O	8:B:933:HOH:O[4_565]	1.83	0.37
8:A:1101:HOH:O	8:A:1101:HOH:O[4_565]	1.85	0.35
1:A:122:GLN:HE22	7:A:712:PO4:O4[4_565]	1.46	0.14
8:A:1202:HOH:O	8:B:1366:HOH:O[4_565]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	651/603 (108%)	632 (97%)	17 (3%)	2 (0%)	41	17
1	B	655/603 (109%)	639 (98%)	14 (2%)	2 (0%)	41	17
All	All	1306/1206 (108%)	1271 (97%)	31 (2%)	4 (0%)	47	17

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	263[A]	ASN
1	A	263[B]	ASN
1	B	263[A]	ASN
1	B	263[B]	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	531/485 (110%)	531 (100%)	0	100	100
1	B	533/485 (110%)	532 (100%)	1 (0%)	93	79
All	All	1064/970 (110%)	1063 (100%)	1 (0%)	93	79

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	245	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 4 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	PO4	B	710	-	4,4,4	1.57	0	6,6,6	1.61	1 (16%)
6	GOL	A	711[A]	-	5,5,5	0.40	0	5,5,5	0.44	0
6	GOL	A	708[B]	-	5,5,5	0.57	0	5,5,5	1.10	1 (20%)
6	GOL	A	710	-	5,5,5	0.47	0	5,5,5	0.55	0
6	GOL	A	709[A]	-	5,5,5	0.54	0	5,5,5	0.75	0
6	GOL	B	708	-	5,5,5	0.25	0	5,5,5	0.37	0
6	GOL	B	706	-	5,5,5	0.35	0	5,5,5	0.24	0
2	FAD	B	701	-	51,58,58	1.46	7 (13%)	60,89,89	2.19	9 (15%)
6	GOL	A	706	-	5,5,5	0.36	0	5,5,5	0.51	0
6	GOL	B	705	-	5,5,5	0.70	0	5,5,5	0.49	0
7	PO4	A	712	-	4,4,4	1.50	0	6,6,6	1.38	1 (16%)
5	TPP	A	704	4	22,27,27	1.15	1 (4%)	29,40,40	1.61	7 (24%)
5	TPP	B	704	4	22,27,27	1.28	3 (13%)	29,40,40	1.37	4 (13%)
6	GOL	A	705	-	5,5,5	0.25	0	5,5,5	0.65	0
6	GOL	B	707	-	5,5,5	0.22	0	5,5,5	0.42	0
6	GOL	B	709[A]	-	5,5,5	0.38	0	5,5,5	0.60	0
2	FAD	A	701	-	51,58,58	1.28	6 (11%)	60,89,89	2.10	12 (20%)
6	GOL	B	709[B]	-	5,5,5	0.64	0	5,5,5	1.03	0
6	GOL	A	707	-	5,5,5	0.42	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	TPP	B	704	4	-	1/16/17/17	0/2/2/2
6	GOL	A	706	-	-	0/4/4/4	-
6	GOL	A	705	-	-	0/4/4/4	-
6	GOL	A	710	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	711[A]	-	-	4/4/4/4	-
6	GOL	A	709[A]	-	-	2/4/4/4	-
6	GOL	B	705	-	-	0/4/4/4	-
6	GOL	B	707	-	-	0/4/4/4	-
6	GOL	B	708	-	-	2/4/4/4	-
6	GOL	B	709[A]	-	-	2/4/4/4	-
2	FAD	A	701	-	-	4/30/50/50	0/6/6/6
6	GOL	B	709[B]	-	-	2/4/4/4	-
6	GOL	B	706	-	-	0/4/4/4	-
5	TPP	A	704	4	-	1/16/17/17	0/2/2/2
6	GOL	A	707	-	-	0/4/4/4	-
6	GOL	A	708[B]	-	-	0/4/4/4	-
2	FAD	B	701	-	-	4/30/50/50	0/6/6/6

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	FAD	C4X-C10	6.36	1.45	1.38
2	A	701	FAD	C4X-C10	4.96	1.43	1.38
5	A	704	TPP	C4-N3	-3.76	1.36	1.39
2	B	701	FAD	C4X-N5	2.84	1.37	1.33
2	A	701	FAD	C8-C7	2.76	1.47	1.40
5	B	704	TPP	C6-C5	2.73	1.52	1.50
2	B	701	FAD	O4B-C1B	2.73	1.44	1.41
5	B	704	TPP	C4-N3	-2.62	1.37	1.39
2	A	701	FAD	C9A-C5X	2.57	1.47	1.42
2	B	701	FAD	C8-C7	2.53	1.47	1.40
2	A	701	FAD	C9A-N10	2.42	1.41	1.38
2	B	701	FAD	C9A-N10	2.41	1.41	1.38
2	B	701	FAD	C9A-C5X	2.34	1.47	1.42
2	B	701	FAD	C4-C4X	2.33	1.45	1.41
2	A	701	FAD	C4-N3	2.29	1.37	1.33
2	A	701	FAD	C4-C4X	2.20	1.45	1.41
5	B	704	TPP	C7'-C5'	2.09	1.55	1.51

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	FAD	C4-N3-C2	9.77	123.39	115.14
2	A	701	FAD	C4-N3-C2	9.29	122.98	115.14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	FAD	C1'-N10-C9A	7.41	124.13	118.29
2	A	701	FAD	C1'-N10-C9A	5.41	122.55	118.29
2	B	701	FAD	C4-C4X-C10	-5.24	116.48	119.95
2	A	701	FAD	C4-C4X-C10	-5.23	116.49	119.95
2	B	701	FAD	C4X-C4-N3	-4.47	117.32	123.43
2	A	701	FAD	C4X-C4-N3	-4.44	117.36	123.43
5	A	704	TPP	C6-C5-C4	4.36	130.94	127.43
2	B	701	FAD	C4X-C10-N10	-3.72	116.48	120.30
7	B	710	PO4	O4-P-O3	3.20	118.25	107.97
2	A	701	FAD	C4X-C10-N10	-3.14	117.07	120.30
2	B	701	FAD	C1B-N9A-C4A	-3.12	121.16	126.64
5	B	704	TPP	C7'-N3-C2	-3.03	119.87	125.35
2	A	701	FAD	C1B-N9A-C4A	-2.99	121.39	126.64
5	A	704	TPP	C6'-N1'-C2'	2.97	121.02	115.96
2	A	701	FAD	C6-C5X-N5	2.92	122.27	119.05
5	B	704	TPP	C5-C4-N3	2.92	113.41	107.57
2	A	701	FAD	O4B-C1B-C2B	-2.81	102.82	106.93
5	B	704	TPP	C5'-C7'-N3	-2.68	108.82	113.28
2	A	701	FAD	N3A-C2A-N1A	-2.67	124.51	128.68
2	A	701	FAD	C9A-N10-C10	-2.59	118.52	121.91
5	A	704	TPP	C7'-N3-C2	-2.58	120.69	125.35
5	A	704	TPP	C5-C4-N3	2.54	112.66	107.57
7	A	712	PO4	O4-P-O3	2.54	116.11	107.97
2	B	701	FAD	C4-C4X-N5	2.51	121.46	118.60
5	A	704	TPP	C5'-C6'-N1'	-2.47	119.70	123.82
2	A	701	FAD	C4-C4X-N5	2.42	121.36	118.60
2	B	701	FAD	C1'-N10-C10	-2.40	116.26	118.41
2	A	701	FAD	C9A-C5X-N5	-2.38	118.64	122.36
5	A	704	TPP	C5'-C7'-N3	-2.33	109.40	113.28
5	A	704	TPP	N1'-C2'-N3'	-2.13	121.88	125.54
5	B	704	TPP	O3B-PB-O2B	2.10	115.67	107.64
2	B	701	FAD	C6-C5X-N5	2.09	121.35	119.05
6	A	708[B]	GOL	O3-C3-C2	2.07	120.15	110.20

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	701	FAD	C3'-C4'-C5'-O5'
2	B	701	FAD	O4'-C4'-C5'-O5'
5	A	704	TPP	PA-O3A-PB-O3B
6	A	711[A]	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	708	GOL	O1-C1-C2-C3
6	B	709[A]	GOL	O1-C1-C2-C3
6	B	709[B]	GOL	O1-C1-C2-C3
6	B	709[A]	GOL	O1-C1-C2-O2
6	A	709[A]	GOL	C1-C2-C3-O3
6	A	710	GOL	C1-C2-C3-O3
6	A	711[A]	GOL	C1-C2-C3-O3
2	A	701	FAD	O4'-C4'-C5'-O5'
6	A	711[A]	GOL	O1-C1-C2-O2
6	B	709[B]	GOL	O1-C1-C2-O2
2	A	701	FAD	C3'-C4'-C5'-O5'
6	A	711[A]	GOL	O2-C2-C3-O3
2	A	701	FAD	P-O3P-PA-O5B
2	B	701	FAD	P-O3P-PA-O5B
5	B	704	TPP	PA-O3A-PB-O2B
6	A	710	GOL	O2-C2-C3-O3
6	B	708	GOL	O1-C1-C2-O2
6	A	709[A]	GOL	O2-C2-C3-O3
2	A	701	FAD	O4B-C4B-C5B-O5B
2	B	701	FAD	O4B-C4B-C5B-O5B


There are no ring outliers.



6 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	710	GOL	1	0
6	A	709[A]	GOL	1	0
7	A	712	PO4	0	1
5	B	704	TPP	1	0
6	B	709[A]	GOL	1	0
6	B	709[B]	GOL	10	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	585/603 (97%)	-0.45	3 (0%) 91 91	7, 10, 19, 35	17 (2%)
1	B	585/603 (97%)	-0.57	4 (0%) 87 87	7, 10, 19, 42	18 (3%)
All	All	1170/1206 (97%)	-0.51	7 (0%) 89 88	7, 10, 19, 42	35 (2%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	560[A]	ALA	4.1
1	B	9	THR	3.6
1	A	9	THR	3.4
1	B	592	ASP	2.8
1	A	561	MET	2.3
1	B	593	ASP	2.2
1	B	591	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	GOL	A	710	6/6	0.65	0.22	30,36,38,39	14
6	GOL	A	709[A]	6/6	0.66	0.17	23,28,33,34	14
6	GOL	B	708	6/6	0.66	0.15	26,32,34,35	14
6	GOL	B	707	6/6	0.71	0.14	32,38,41,42	0
6	GOL	B	709[A]	6/6	0.72	0.30	18,21,24,25	14
6	GOL	B	709[B]	6/6	0.72	0.30	18,22,25,25	14
6	GOL	B	706	6/6	0.77	0.20	25,30,33,34	14
6	GOL	A	711[A]	6/6	0.78	0.15	28,34,35,35	14
6	GOL	A	708[B]	6/6	0.80	0.19	13,19,24,26	14
6	GOL	A	707	6/6	0.86	0.12	29,35,36,36	14
6	GOL	B	705	6/6	0.92	0.10	18,22,25,25	14
6	GOL	A	706	6/6	0.92	0.12	20,25,29,30	14
6	GOL	A	705	6/6	0.95	0.08	12,18,20,22	14
7	PO4	A	712	5/5	0.98	0.09	12,12,14,14	5
7	PO4	B	710	5/5	0.98	0.10	9,10,13,13	5
5	TPP	B	704	26/26	0.99	0.08	7,7,8,9	0
2	FAD	A	701	53/53	0.99	0.07	6,8,10,10	0
2	FAD	B	701	53/53	0.99	0.06	6,7,8,9	0
5	TPP	A	704	26/26	0.99	0.07	6,8,8,9	0
3	K	B	702	1/1	1.00	0.03	9,9,9,9	1
4	MG	A	703	1/1	1.00	0.05	8,8,8,8	0
4	MG	B	703	1/1	1.00	0.05	7,7,7,7	0
3	K	A	702	1/1	1.00	0.05	9,9,9,9	1

6.5 Other polymers [i](#)

There are no such residues in this entry.