



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 7, 2026 – 04:06 pm BST

PDB ID : 9HA9 / pdb\_00009ha9  
Title : Central domain fragment of Glucan Water Dikinase-1 from *S.tuberosum*  
Authors : Laffargue, T.; Cioci, G.; Cooper, N.; Remaud-Simeon, M.; Moulis, C.  
Deposited on : 2024-11-02  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*



The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	467	 86% 9% 5%
1	B	467	 86% 9% 5%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7156 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-glucan water dikinase, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	0	1	0
			3562	2242	630	673	17			
1	B	445	Total	C	N	O	S	0	1	0
			3541	2230	626	668	17			

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	461	MET	-	initiating methionine	UNP Q9AWA5
A	462	GLY	-	expression tag	UNP Q9AWA5
A	463	SER	-	expression tag	UNP Q9AWA5
A	464	SER	-	expression tag	UNP Q9AWA5
A	465	HIS	-	expression tag	UNP Q9AWA5
A	466	HIS	-	expression tag	UNP Q9AWA5
A	467	HIS	-	expression tag	UNP Q9AWA5
A	468	HIS	-	expression tag	UNP Q9AWA5
A	469	HIS	-	expression tag	UNP Q9AWA5
A	470	HIS	-	expression tag	UNP Q9AWA5
A	471	SER	-	expression tag	UNP Q9AWA5
A	472	SER	-	expression tag	UNP Q9AWA5
A	473	GLY	-	expression tag	UNP Q9AWA5
A	474	LEU	-	expression tag	UNP Q9AWA5
A	475	VAL	-	expression tag	UNP Q9AWA5
A	476	PRO	-	expression tag	UNP Q9AWA5
A	477	ARG	-	expression tag	UNP Q9AWA5
B	461	MET	-	initiating methionine	UNP Q9AWA5
B	462	GLY	-	expression tag	UNP Q9AWA5
B	463	SER	-	expression tag	UNP Q9AWA5
B	464	SER	-	expression tag	UNP Q9AWA5
B	465	HIS	-	expression tag	UNP Q9AWA5
B	466	HIS	-	expression tag	UNP Q9AWA5
B	467	HIS	-	expression tag	UNP Q9AWA5
B	468	HIS	-	expression tag	UNP Q9AWA5

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	469	HIS	-	expression tag	UNP Q9AWA5
B	470	HIS	-	expression tag	UNP Q9AWA5
B	471	SER	-	expression tag	UNP Q9AWA5
B	472	SER	-	expression tag	UNP Q9AWA5
B	473	GLY	-	expression tag	UNP Q9AWA5
B	474	LEU	-	expression tag	UNP Q9AWA5
B	475	VAL	-	expression tag	UNP Q9AWA5
B	476	PRO	-	expression tag	UNP Q9AWA5
B	477	ARG	-	expression tag	UNP Q9AWA5

- Molecule 2 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	29	Total O 29 29	0	0
2	B	24	Total O 24 24	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Alpha-glucan water dikinase, chloroplastic

Chain A: 



- Molecule 1: Alpha-glucan water dikinase, chloroplastic

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.31Å 110.31Å 209.18Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.00 50.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.00-3.00) 100.0 (50.00-3.00)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.01 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, $R_{free}$	0.215 , 0.238 0.211 , 0.233	Depositor DCC
$R_{free}$ test set	1336 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	100.8	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 65.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7156	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	116.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.44	0/3626	0.93	0/4899
1	B	0.44	0/3605	0.93	0/4872
All	All	0.44	0/7231	0.93	0/9771

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3562	0	3538	25	0
1	B	3541	0	3512	25	0
2	A	29	0	0	0	0
2	B	24	0	0	0	0
All	All	7156	0	7050	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 49 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:530[B]:ARG:HH11	1:B:530[B]:ARG:HG3	1.08	1.16

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:530[B]:ARG:HH11	1:B:530[B]:ARG:CG	1.78	0.96
1:B:530[B]:ARG:HG3	1:B:530[B]:ARG:NH1	1.72	0.93
1:A:530[B]:ARG:HG2	1:A:902:ILE:HD11	1.82	0.61
1:B:535:ARG:NH1	1:B:626:ASP:OD1	2.36	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	447/467 (96%)	434 (97%)	13 (3%)	0	100	100
1	B	444/467 (95%)	432 (97%)	12 (3%)	0	100	100
All	All	891/934 (95%)	866 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	383/398 (96%)	377 (98%)	6 (2%)	55	79
1	B	380/398 (96%)	375 (99%)	5 (1%)	61	81
All	All	763/796 (96%)	752 (99%)	11 (1%)	59	80



5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	728	VAL
1	B	825	ASP
1	B	915	LEU
1	B	829	ASP
1	A	829	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	874	HIS
1	B	613	GLN
1	B	562	GLN
1	B	540	ASN
1	B	592	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å <sup>2</sup> )	Q < 0.9
1	A	448/467 (95%)	-0.45	1 (0%)	91 83	43, 109, 144, 188	1 (0%)
1	B	445/467 (95%)	-0.36	1 (0%)	91 83	47, 120, 160, 201	1 (0%)
All	All	893/934 (95%)	-0.41	2 (0%)	91 83	43, 115, 156, 201	2 (0%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	480	GLY	3.5
1	A	478	GLY	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.