

wwPDB X-ray Structure Validation Summary Report (i)

Oct 31, 2023 – 03:15 PM JST

PDB ID : 8HCD

Title: Crystal structure of mTREX1-DNA product complex (dNMP)

Authors: Hsiao, Y.Y.; Huang, K.W.; Wu, C.Y.

Deposited on : 2022-11-01

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution		
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$		
R_{free}	130704	8085 (2.00-2.00)		
Clashscore	141614	9178 (2.00-2.00)		
Ramachandran outliers	138981	9054 (2.00-2.00)		
Sidechain outliers	138945	9053 (2.00-2.00)		
RSRZ outliers	127900	7900 (2.00-2.00)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	В	276	76%		20%			
1	С	276	79%		18%			
1	D	276	5% 75%	6%	19%			
1	Е	276	76%	6% •	18%			



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 14770 atoms, of which 6972 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Three-prime repair exonuclease 1.

Mol	Chain	Residues		Atoms						AltConf	Trace
1	1 B 220	220	Total	С	Н	N	О	S	0	0	0
1	Ъ	220	3399	1075	1699	299	317	9		0	
1	C	227	Total	С	Н	N	О	S	0	0	0
1		221	3512	1110	1758	309	326	9			
1	D	223	Total	С	Н	N	О	S	0	0	0
1	D	223	3452	1092	1727	303	321	9	U		0
1	1 E	E 227	Total	С	Н	N	О	S	0	0	0
1	E	221	3485	1098	1742	308	328	9			U

There are 136 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	-33	MET	-	initiating methionine	UNP Q91XB0
В	-32	GLY	-	expression tag	UNP Q91XB0
В	-31	SER	-	expression tag	UNP Q91XB0
В	-30	SER	-	expression tag	UNP Q91XB0
В	-29	HIS	-	expression tag	UNP Q91XB0
В	-28	HIS	-	expression tag	UNP Q91XB0
В	-27	HIS	-	expression tag	UNP Q91XB0
В	-26	HIS	-	expression tag	UNP Q91XB0
В	-25	HIS	-	expression tag	UNP Q91XB0
В	-24	HIS	-	expression tag	UNP Q91XB0
В	-23	SER	-	expression tag	UNP Q91XB0
В	-22	SER	-	expression tag	UNP Q91XB0
В	-21	GLY	-	expression tag	UNP Q91XB0
В	-20	LEU	-	expression tag	UNP Q91XB0
В	-19	VAL	-	expression tag	UNP Q91XB0
В	-18	PRO	-	expression tag	UNP Q91XB0
В	-17	ARG	-	expression tag	UNP Q91XB0
В	-16	GLY	-	expression tag	UNP Q91XB0
В	-15	SER	-	expression tag	UNP Q91XB0
В	-14	HIS	-	expression tag	UNP Q91XB0
В	-13	MET	-	expression tag	UNP Q91XB0



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-12	ALA	-	expression tag	UNP Q91XB0
В	-11	SER	-	expression tag	UNP Q91XB0
В	-10	MET	-	expression tag	UNP Q91XB0
В	-9	THR	-	expression tag	UNP Q91XB0
В	-8	GLY	-	expression tag	UNP Q91XB0
В	-7	GLY	-	expression tag	UNP Q91XB0
В	-6	GLN	-	expression tag	UNP Q91XB0
В	-5	GLN	-	expression tag	UNP Q91XB0
В	-4	MET	-	expression tag	UNP Q91XB0
В	-3	GLY	-	expression tag	UNP Q91XB0
В	-2	ARG	-	expression tag	UNP Q91XB0
В	-1	GLY	-	expression tag	UNP Q91XB0
В	0	SER	-	expression tag	UNP Q91XB0
С	-33	MET	-	initiating methionine	UNP Q91XB0
С	-32	GLY	-	expression tag	UNP Q91XB0
С	-31	SER	-	expression tag	UNP Q91XB0
С	-30	SER	-	expression tag	UNP Q91XB0
С	-29	HIS	-	expression tag	UNP Q91XB0
С	-28	HIS	-	expression tag	UNP Q91XB0
С	-27	HIS	-	expression tag	UNP Q91XB0
С	-26	HIS	-	expression tag	UNP Q91XB0
С	-25	HIS	-	expression tag	UNP Q91XB0
С	-24	HIS	-	expression tag	UNP Q91XB0
С	-23	SER	-	expression tag	UNP Q91XB0
С	-22	SER	-	expression tag	UNP Q91XB0
С	-21	GLY	-	expression tag	UNP Q91XB0
С	-20	LEU	-	expression tag	UNP Q91XB0
С	-19	VAL	-	expression tag	UNP Q91XB0
С	-18	PRO	-	expression tag	UNP Q91XB0
С	-17	ARG	-	expression tag	UNP Q91XB0
С	-16	GLY	-	expression tag	UNP Q91XB0
С	-15	SER	-	expression tag	UNP Q91XB0
С	-14	HIS	-	expression tag	UNP Q91XB0
С	-13	MET		expression tag	UNP Q91XB0
С	-12	ALA	-	expression tag	UNP Q91XB0
С	-11	SER		expression tag	UNP Q91XB0
С	-10	MET	-	expression tag	UNP Q91XB0
С	-9	THR		expression tag	UNP Q91XB0
С	-8	GLY		expression tag	UNP Q91XB0
С	-7	GLY	-	expression tag	UNP Q91XB0
С	-6	GLN	-	expression tag	UNP Q91XB0
С	-5	GLN	-	expression tag	UNP Q91XB0



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	-4	MET	-	expression tag	UNP Q91XB0
С	-3	GLY	-	expression tag	UNP Q91XB0
С	-2	ARG	-	expression tag	UNP Q91XB0
С	-1	GLY	-	expression tag	UNP Q91XB0
С	0	SER	-	expression tag	UNP Q91XB0
D	-33	MET	-	initiating methionine	UNP Q91XB0
D	-32	GLY	-	expression tag	UNP Q91XB0
D	-31	SER	-	expression tag	UNP Q91XB0
D	-30	SER	-	expression tag	UNP Q91XB0
D	-29	HIS	-	expression tag	UNP Q91XB0
D	-28	HIS	-	expression tag	UNP Q91XB0
D	-27	HIS	-	expression tag	UNP Q91XB0
D	-26	HIS	-	expression tag	UNP Q91XB0
D	-25	HIS	-	expression tag	UNP Q91XB0
D	-24	HIS	-	expression tag	UNP Q91XB0
D	-23	SER	-	expression tag	UNP Q91XB0
D	-22	SER	-	expression tag	UNP Q91XB0
D	-21	GLY	-	expression tag	UNP Q91XB0
D	-20	LEU	-	expression tag	UNP Q91XB0
D	-19	VAL	-	expression tag	UNP Q91XB0
D	-18	PRO	-	expression tag	UNP Q91XB0
D	-17	ARG	-	expression tag	UNP Q91XB0
D	-16	GLY	-	expression tag	UNP Q91XB0
D	-15	SER	-	expression tag	UNP Q91XB0
D	-14	HIS	-	expression tag	UNP Q91XB0
D	-13	MET	_	expression tag	UNP Q91XB0
D	-12	ALA	-	expression tag	UNP Q91XB0
D	-11	SER	-	expression tag	UNP Q91XB0
D	-10	MET	_	expression tag	UNP Q91XB0
D	-9	THR	-	expression tag	UNP Q91XB0
D	-8	GLY	-	expression tag	UNP Q91XB0
D	-7	GLY	_	expression tag	UNP Q91XB0
D	-6	GLN	-	expression tag	UNP Q91XB0
D	-5	GLN	_	expression tag	UNP Q91XB0
D	-4	MET	-	expression tag	UNP Q91XB0
D	-3	GLY	-	expression tag	UNP Q91XB0
D	-2	ARG	-	expression tag	UNP Q91XB0
D	-1	GLY	-	expression tag	UNP Q91XB0
D	0	SER	-	expression tag	UNP Q91XB0
Е	-33	MET	-	initiating methionine	UNP Q91XB0
Е	-32	GLY	-	expression tag	UNP Q91XB0
Е	-31	SER	-	expression tag	UNP Q91XB0



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-30	SER	-	expression tag	UNP Q91XB0
Е	-29	HIS	-	expression tag	UNP Q91XB0
Е	-28	HIS	-	expression tag	UNP Q91XB0
Е	-27	HIS	-	expression tag	UNP Q91XB0
Е	-26	HIS	-	expression tag	UNP Q91XB0
Е	-25	HIS	-	expression tag	UNP Q91XB0
E	-24	HIS	-	expression tag	UNP Q91XB0
E	-23	SER	-	expression tag	UNP Q91XB0
Е	-22	SER	-	expression tag	UNP Q91XB0
E	-21	GLY	-	expression tag	UNP Q91XB0
E	-20	LEU	-	expression tag	UNP Q91XB0
E	-19	VAL	-	expression tag	UNP Q91XB0
E	-18	PRO	-	expression tag	UNP Q91XB0
Е	-17	ARG	-	expression tag	UNP Q91XB0
E	-16	GLY	-	expression tag	UNP Q91XB0
Е	-15	SER	-	expression tag	UNP Q91XB0
E	-14	HIS	-	expression tag	UNP Q91XB0
E	-13	MET	-	expression tag	UNP Q91XB0
E	-12	ALA	-	expression tag	UNP Q91XB0
E	-11	SER	-	expression tag	UNP Q91XB0
Е	-10	MET	-	expression tag	UNP Q91XB0
E	-9	THR	-	expression tag	UNP Q91XB0
E	-8	GLY	-	expression tag	UNP Q91XB0
Ε	-7	GLY	-	expression tag	UNP Q91XB0
Ε	-6	GLN	-	expression tag	UNP Q91XB0
E	-5	GLN	-	expression tag	UNP Q91XB0
Е	-4	MET	-	expression tag	UNP Q91XB0
Е	-3	GLY	-	expression tag	UNP Q91XB0
Е	-2	ARG	-	expression tag	UNP Q91XB0
Е	-1	GLY	-	expression tag	UNP Q91XB0
Е	0	SER	-	expression tag	UNP Q91XB0

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	2	Total Mg 2 2	0	0
2	С	2	Total Mg 2 2	0	0
2	D	2	Total Mg 2 2	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	E	2	Total Mg 2 2	0	0


• Molecule 3 is 2'-DEOXYCYTIDINE-5'-MONOPHOSPHATE (three-letter code: DCM) (formula: $C_9H_{14}N_3O_7P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	В	1	Total C H N O I 32 9 12 3 7	0	0
3	D	1	Total C H N O I 31 9 11 3 7	1 ()	0
3	Е	1	Total C H N O I 31 9 11 3 7	0	0

• Molecule 4 is 2'-DEOXYGUANOSINE-5'-MONOPHOSPHATE (three-letter code: DGP) (formula: $C_{10}H_{14}N_5O_7P$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
1	С	1	Total	С	Н	N	О	Р	0	0
4		1	35	10	12	5	7	1	0	

• Molecule 5 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	Е	1	Total (Co 1	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	190	Total O 190 190	0	0
6	С	171	Total O 171 171	0	0
6	D	192	Total O 192 192	0	0
6	E	231	Total O 231 231	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Three-prime repair exonuclease 1











4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	67.68Å 81.28Å 93.21Å	Depositor
a, b, c, α , β , γ	90.00° 103.69° 90.00°	Depositor
Resolution (Å)	29.29 - 2.00	Depositor
rtesolution (A)	29.29 - 2.00	EDS
% Data completeness	99.4 (29.29-2.00)	Depositor
(in resolution range)	99.3 (29.29-2.00)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.57 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.14_3260	Depositor
P. P.	0.195 , 0.240	Depositor
R, R_{free}	0.196 , 0.241	DCC
R_{free} test set	5152 reflections (7.81%)	wwPDB-VP
Wilson B-factor (Å ²)	27.6	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.38, 45.8	EDS
L-test for twinning ²	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14770	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.94% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CO, DGP, MG, DCM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	В	0.24	0/1741	0.45	0/2373
1	С	0.24	0/1798	0.45	0/2454
1	D	0.24	0/1767	0.43	0/2410
1	Е	0.25	0/1785	0.45	0/2434
All	All	0.24	0/7091	0.44	0/9671

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	В	1700	1699	1699	7	0
1	С	1754	1758	1758	8	0
1	D	1725	1727	1727	10	0
1	Е	1743	1742	1742	10	0
2	В	2	0	0	0	0
2	С	2	0	0	0	0
2	D	2	0	0	0	0
2	Е	2	0	0	0	0
3	В	20	12	12	0	0



Continued	trom	mromonie	maaa
Continueu	110111	predidus	puyc

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	20	11	12	0	0
3	Ε	20	11	12	0	0
4	С	23	12	12	1	0
5	Ε	1	0	0	0	0
6	В	190	0	0	0	0
6	С	171	0	0	6	0
6	D	192	0	0	4	0
6	Ε	231	0	0	3	0
All	All	7798	6972	6974	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 34 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:C:113:GLN:NE2	6:C:602:HOH:O	2.18	0.76
1:C:91:GLU:OE2	6:C:601:HOH:O	2.05	0.75
1:B:166:SER:HA	1:B:167:SER:CB	2.21	0.71
1:D:189:GLN:NE2	6:D:403:HOH:O	2.26	0.68
1:B:42:ARG:NH1	1:D:91:GLU:OE1	2.27	0.67

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	В	$216/276\ (78\%)$	211 (98%)	5 (2%)	0	100	100
1	С	$223/276\ (81\%)$	220 (99%)	3 (1%)	0	100	100
1	D	$219/276\ (79\%)$	213 (97%)	6 (3%)	0	100	100



 $Continued\ from\ previous\ page...$

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	E	225/276~(82%)	222 (99%)	3 (1%)	0	100	100
All	All	883/1104 (80%)	866 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	В	189/232 (82%)	187 (99%)	2 (1%)	73	78
1	\mathbf{C}	196/232~(84%)	193 (98%)	3 (2%)	65	69
1	D	192/232 (83%)	190 (99%)	2 (1%)	76	81
1	E	194/232 (84%)	189 (97%)	5 (3%)	46	48
All	All	771/928 (83%)	759 (98%)	12 (2%)	62	67

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Ε	119	CYS
1	Е	164	GLN
1	Е	230	LYS
1	Ε	171	ASN
1	С	59	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	Ε	10	HIS
1	Ε	12	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type	Tuno	e Chain	Res	Link	В	ond leng	gths	Bond angles		
	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	DCM	Е	301	2	21,21,21	3.53	12 (57%)	31,31,31	1.04	1 (3%)
3	DCM	В	302	2	21,21,21	3.54	12 (57%)	31,31,31	0.93	1 (3%)
3	DCM	D	301	2	21,21,21	3.55	12 (57%)	31,31,31	0.97	0
4	DGP	С	301	2	22,25,25	3.61	13 (59%)	26,38,38	1.36	5 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	DCM	Е	301	2	-	1/10/22/22	0/2/2/2
3	DCM	В	302	2	-	3/10/22/22	0/2/2/2
3	DCM	D	301	2	-	2/10/22/22	0/2/2/2
4	DGP	С	301	2	-	1/6/22/22	0/3/3/3



The worst	5	of	49	bond	length	outliers	are	listed	below:
THE WOLDS	\circ	$O_{\mathbf{I}}$	10	DOM	10115 011	Outilities	COL C	mouca	DCIOW.

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$Ideal(\AA)$
3	В	302	DCM	C2'-C3'	-8.92	1.29	1.52
3	D	301	DCM	C2'-C3'	-8.83	1.29	1.52
3	Е	301	DCM	C2'-C3'	-8.82	1.29	1.52
4	С	301	DGP	O4'-C4'	8.21	1.63	1.45
4	С	301	DGP	C3'-C4'	-8.10	1.30	1.53

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
4	С	301	DGP	C5-C6-N1	3.40	119.95	113.95
4	С	301	DGP	C2-N1-C6	-3.04	119.50	125.10
4	С	301	DGP	C8-N7-C5	2.62	107.98	102.99
3	Е	301	DCM	C2'-C1'-N1	-2.45	108.13	113.77
4	С	301	DGP	O6-C6-C5	-2.31	119.87	124.37

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	В	302	DCM	C5'-O5'-P-O2P
3	В	302	DCM	C5'-O5'-P-O3P
3	D	301	DCM	C5'-O5'-P-O1P
3	Е	301	DCM	C5'-O5'-P-O2P
3	В	302	DCM	C5'-O5'-P-O1P

There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	С	301	DGP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.




The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	#RS	\mathbf{RZ}	>2	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q < 0.9
1	В	220/276~(79%)	0.23	9 (4%)	37	36	20, 33, 59, 114	0
1	С	227/276 (82%)	0.35	15 (6%)	18	17	19, 33, 77, 108	0
1	D	223/276 (80%)	0.32	13 (5%)	23	22	19, 35, 66, 103	0
1	Е	227/276 (82%)	0.16	10 (4%)	34	33	17, 30, 70, 119	0
All	All	897/1104 (81%)	0.27	47 (5%)	27	26	17, 33, 70, 119	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	235	THR	12.0
1	С	235	THR	8.0
1	С	5	THR	6.9
1	В	8	HIS	6.1
1	С	236	PRO	5.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	MG	С	303	1/1	0.38	0.20	50,50,50,50	0
2	MG	В	303	1/1	0.61	0.13	47,47,47,47	0
2	MG	D	303	1/1	0.80	0.12	50,50,50,50	0
3	DCM	D	301	20/20	0.81	0.18	35,49,56,60	0
5	CO	E	304	1/1	0.90	0.07	69,69,69,69	0
3	DCM	В	302	20/20	0.92	0.12	31,37,44,46	0
4	DGP	С	301	23/23	0.93	0.13	25,39,49,50	0
3	DCM	Е	301	20/20	0.94	0.11	32,40,48,53	0
2	MG	Е	302	1/1	0.96	0.09	24,24,24,24	0
2	MG	С	302	1/1	0.96	0.24	20,20,20,20	0
2	MG	D	302	1/1	0.97	0.14	29,29,29,29	0
2	MG	В	301	1/1	0.98	0.13	25,25,25,25	0
2	MG	E	303	1/1	1.00	0.05	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


















Electron density around DCM D 301: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)







































6.5 Other polymers (i)

There are no such residues in this entry.

