

wwPDB X-ray Structure Validation Summary Report (i)

Aug 27, 2023 – 06:43 PM EDT

PDB ID	:	3HDY
Title	:	Crystal Structure of UDP-galactopyranose mutase (reduced form) in complex
		with substrate
Authors	:	Partha, S.K.; van Straaten, K.E.; Sanders, D.A.R.
Deposited on	:	2009-05-07
Resolution	:	2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.35
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\;DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain								
1	А	397	2% 7 0%	21% • 8%							
1	В	397	73%	16% • 9%							
1	С	397	58%	27% • 12%							
1	D	397	4% 66%	23% • 9%							
1	Е	397	^{3%} 70%	20% • 9%							



Mol	Chain	Length	Quality of chain			
1	F	397	3%	20%		09/
1	1	001	7070	20%	•	970
1	G	397	73%	16%	•	9%
1	Н	397	71%	18%	•	9%
			2%	20,0		
1	Ι	397	69%	20%	·	9%
1	J	397	73%	17%	·	10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	FDA	В	1385	X	-	-	-
4	FDA	D	1385	Х	-	-	-
4	FDA	G	1385	Х	-	-	-



 $\mathbf{2}$

Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 30890 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
1	Δ	365	Total	С	Ν	Ο	S	0	0	0
1	Л	505	2984	1904	518	554	8	0	0	0
1	В	B 361	Total	С	Ν	Ο	\mathbf{S}	0	0	0
1	D	501	2951	1885	512	546	8	0	0	0
1	С	340	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	19	0	0
1	U	045	2850	1820	493	529	8	12	0	0
1	Л	361	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	Q	0	0
1	D	501	2951	1885	512	546	8	0	0	0
1	1 F	361	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	12	0	0
			2951	1885	512	546	8		0	Ŭ ,
1	F	362	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	10	0	0
1	Ľ		2960	1891	514	547	8	19	0	0
1	G	360	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0	0
1	G	500	2947	1883	511	545	8	0	0	0
1	н	361	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	0	0	0
1	11	501	2956	1889	513	546	8	0	0	0
1	1 T	361	Total	\mathbf{C}	Ν	Ο	\mathbf{S}	1	0	0
	1	501	2951	1885	512	546	8	L	0	0
1	1 T	350	Total	С	Ν	0	S	1	0	0
I J	009	2943	1881	510	544	8		0	U	

• Molecule 1 is a protein called UDP-galactopyranose mutase.

• Molecule 2 is GALACTOSE-URIDINE-5'-DIPHOSPHATE (three-letter code: GDU) (formula: $C_{15}H_{24}N_2O_{17}P_2$).





Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	
9	Λ	1	Total	С	Ν	Ο	Р	0	0	
		I	36	15	2	17	2	0	0	
2	В	1	Total	С	Ν	Ο	Р	0	0	
2	D	I	36	15	2	17	2	0	0	
2	С	1	Total	\mathbf{C}	Ν	Ο	Р	0	0	
2	U	I	36	15	2	17	2	0	0	
2	Л	1	Total	\mathbf{C}	Ν	Ο	Р	0	0	
2	D	I	36	15	2	17	2	0	0	
2	E	1	Total	С	Ν	Ο	Р	0	0	
	Ц	1	36	15	2	17	2	0	Ŭ	
2	F	1	Total	\mathbf{C}	Ν	Ο	Р	0	0	
	1	1	36	15	2	17	2	0	0	
2	G	1	Total	\mathbf{C}	Ν	Ο	Р	0	0	
	ŭ	Ĩ	36	15	2	17	2	0	0	
2	н	1	Total	\mathbf{C}	Ν	Ο	Р	0	0	
	11	I	36	15	2	17	2	0	0	
2	Т	1	Total	\mathbf{C}	Ν	Ο	Р	0	0	
	1	1	36	15	2	17	2	0	0	
2	T	T	1	Total	\mathbf{C}	Ν	Ο	Р	0	0
	J	L	36	15	2	17	2	U		

- Molecule 3 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $\rm C_{27}H_{33}N_9O_{15}P_2).$





Mol	Chain	Residues		Ate	oms			ZeroOcc	AltConf	
3	Δ	1	Total	С	Ν	Ο	Р	0	0	
0	Л	T	53	27	9	15	2	0	0	
3	С	1	Total	С	Ν	Ο	Р	0	0	
0	U	T	53	27	9	15	2	0	0	
3	F	1	Total	С	Ν	Ο	Р	0	0	
0	Ľ	1	53	27	9	15	2	0	0	
3	F	F 1	Total	С	Ν	Ο	Р	0	0	
0	T,		53	27	9	15	2	0	0	
3	н	1	Total	С	Ν	Ο	Р	0	0	
0	11	T	53	27	9	15	2	0	0	
3	т	1	Total	С	Ν	Ο	Р	0	0	
0	I	T	53	27	9	15	2	0	0	
3	T	1	Total	С	Ν	0	Р	0	0	
J	J	L	53	27	9	15	2	0	0	

• Molecule 4 is DIHYDROFLAVINE-ADENINE DINUCLEOTIDE (three-letter code: FDA) (formula: $C_{27}H_{35}N_9O_{15}P_2$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	D	1	Total	С	Ν	Ο	Р	0	0	
4 B		53	27	9	15	2	0	0		
4	Л	1	Total	С	Ν	Ο	Р	0	0	
4	4 D	1	53	27	9	15	2			
4	С	1	Total	С	Ν	Ο	Р	0	0	
4	G		53	27	9	15	2	0	U	

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	70	Total O 70 70	0	0
5	В	51	Total O 51 51	0	0
5	С	41	Total O 41 41	0	0
5	D	48	Total O 48 48	0	0
5	Е	44	Total O 44 44	0	0
5	F	60	Total O 60 60	0	0
5	G	47	$\begin{array}{cc} \text{Total} & \text{O} \\ 47 & 47 \end{array}$	0	0
5	Н	75	Total O 75 75	0	0
5	Ι	53	Total O 53 53	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	J	67	Total O 67 67	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: UDP-galactopyranose mutase













Q388 GLN GLN GLN GLU GLU GLV ASN ASN GLU

• Molecule 1: UDP-galactopyranose mutase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	132.82Å 174.63Å 218.19Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Bosolution(Å)	39.94 - 2.40	Depositor
Resolution (A)	39.94 - 2.40	EDS
% Data completeness	95.8 (39.94-2.40)	Depositor
(in resolution range)	95.8 (39.94-2.40)	EDS
R _{merge}	0.19	Depositor
R_{sym}	0.19	Depositor
$< I/\sigma(I) > 1$	$1.74 (at 2.39 \text{\AA})$	Xtriage
Refinement program	PHENIX (phenix.refine)	Depositor
B B.	0.235 , 0.280	Depositor
n, n_{free}	0.236 , 0.277	DCC
R_{free} test set	9506 reflections (5.02%)	wwPDB-VP
Wilson B-factor $(Å^2)$	43.8	Xtriage
Anisotropy	0.198	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38 , 53.5	EDS
L-test for $twinning^2$	$ < L >=0.52, < L^2>=0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	30890	wwPDB-VP
Average B, all atoms $(Å^2)$	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.22% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FDA, FAD, GDU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.29	0/3068	0.43	0/4171
1	В	0.34	0/3035	0.43	0/4128
1	С	0.46	0/2931	0.50	0/3989
1	D	0.31	0/3035	0.43	0/4128
1	Е	0.31	0/3035	0.42	0/4128
1	F	0.32	0/3044	0.43	0/4139
1	G	0.30	0/3031	0.43	0/4123
1	Н	0.30	0/3040	0.42	0/4134
1	Ι	0.26	0/3035	0.44	0/4128
1	J	0.27	1/3027~(0.0%)	0.43	0/4118
All	All	0.32	1/30281~(0.0%)	0.44	0/41186

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	С	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	J	301	TYR	C-N	-5.46	1.21	1.34

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:



Mol	Chain	\mathbf{Res}	Type	Group
1	С	50	SER	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2984	0	2854	53	0
1	В	2951	0	2822	53	0
1	С	2850	0	2728	100	0
1	D	2951	0	2822	67	0
1	Е	2951	0	2822	47	0
1	F	2960	0	2835	57	0
1	G	2947	0	2819	45	0
1	Н	2956	0	2832	50	0
1	Ι	2951	0	2822	51	0
1	J	2943	0	2815	44	0
2	А	36	0	22	4	0
2	В	36	0	22	4	0
2	С	36	0	22	7	0
2	D	36	0	22	6	0
2	Е	36	0	22	2	0
2	F	36	0	22	3	0
2	G	36	0	22	2	0
2	Н	36	0	22	1	0
2	Ι	36	0	22	2	0
2	J	36	0	22	0	0
3	А	53	0	31	4	0
3	С	53	0	31	4	0
3	Е	53	0	31	2	0
3	F	53	0	31	3	0
3	Н	53	0	31	2	0
3	Ι	53	0	31	2	0
3	J	53	0	31	3	0
4	В	53	0	28	2	0
4	D	53	0	29	3	0
4	G	53	0	28	4	0
5	А	70	0	0	1	0
5	В	51	0	0	0	0
5	С	41	0	0	3	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	48	0	0	2	0
5	Ε	44	0	0	0	0
5	F	60	0	0	0	0
5	G	47	0	0	1	0
5	Н	75	0	0	0	0
5	Ι	53	0	0	1	0
5	J	67	0	0	0	0
All	All	30890	0	28693	552	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 552 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:152:LYS:HE2	1:F:152:LYS:HA	1.48	0.96
1:J:152:LYS:HE2	1:J:152:LYS:HA	1.50	0.91
1:I:152:LYS:HE2	1:I:152:LYS:HA	1.52	0.91
1:D:152:LYS:HE2	1:D:152:LYS:HA	1.51	0.89
1:B:152:LYS:HA	1:B:152:LYS:HE2	1.55	0.88

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	363/397~(91%)	347~(96%)	16 (4%)	0	100	100
1	В	359/397~(90%)	347 (97%)	12 (3%)	0	100	100
1	С	347/397~(87%)	314 (90%)	30 (9%)	3 (1%)	17	25
1	D	359/397~(90%)	342 (95%)	17 (5%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	Ε	359/397~(90%)	347~(97%)	12 (3%)	0	100 100
1	F	360/397~(91%)	343~(95%)	15 (4%)	2(1%)	25 36
1	G	358/397~(90%)	342~(96%)	16 (4%)	0	100 100
1	Н	359/397~(90%)	347~(97%)	12 (3%)	0	100 100
1	Ι	359/397~(90%)	343~(96%)	15 (4%)	1 (0%)	41 55
1	J	357/397~(90%)	341 (96%)	16 (4%)	0	100 100
All	All	3580/3970~(90%)	3413 (95%)	161 (4%)	6~(0%)	47 62

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	С	51	SER
1	С	233	PRO
1	F	357	GLN
1	F	370	TYR
1	Ι	370	TYR

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	317/346~(92%)	301~(95%)	16~(5%)	24	40
1	В	313/346~(90%)	299~(96%)	14 (4%)	27	44
1	С	304/346~(88%)	281 (92%)	23~(8%)	13	20
1	D	313/346~(90%)	300~(96%)	13~(4%)	30	47
1	Ε	313/346~(90%)	300~(96%)	13~(4%)	30	47
1	F	314/346~(91%)	296 (94%)	18 (6%)	20	33
1	G	313/346~(90%)	299~(96%)	14 (4%)	27	44
1	Η	314/346~(91%)	299~(95%)	15~(5%)	25	41
1	Ι	313/346~(90%)	297 (95%)	16 (5%)	24	39
1	J	313/346~(90%)	303~(97%)	10 (3%)	39	59



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	3127/3460~(90%)	2975~(95%)	152~(5%)	25	40

5 of 152 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Н	120	LEU
1	J	120	LEU
1	Н	186	LEU
1	Ι	120	LEU
1	J	370	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	Ε	357	GLN
1	Н	227	GLN
1	Н	220	HIS
1	Н	357	GLN
1	С	220	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

20 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Tune	Chain	Dec	Tink	В	ond leng	gths	B	ond ang	gles
INIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	GDU	G	500	-	35,38,38	2.68	4 (11%)	$53,\!58,\!58$	<mark>3.62</mark>	16 (30%)
3	FAD	А	450	-	53,58,58	1.30	5 (9%)	68,89,89	1.41	9 (13%)
2	GDU	J	500	-	35,38,38	2.67	4 (11%)	53,58,58	<mark>3.61</mark>	14 (26%)
4	FDA	D	1385	-	52,58,58	2.69	17 (32%)	60,89,89	2.18	19 (31%)
2	GDU	Е	500	-	35,38,38	2.66	3 (8%)	53,58,58	<mark>3.63</mark>	17 (32%)
3	FAD	С	450	-	53,58,58	1.30	5 (9%)	68,89,89	1.41	9 (13%)
2	GDU	В	500	-	35,38,38	2.64	3 (8%)	53,58,58	<mark>3.67</mark>	18 (33%)
2	GDU	D	500	-	35,38,38	2.66	4 (11%)	53,58,58	<mark>3.60</mark>	15 (28%)
2	GDU	С	500	-	35,38,38	2.66	4 (11%)	53,58,58	<mark>3.65</mark>	17 (32%)
3	FAD	Ι	450	-	53,58,58	1.30	5 (9%)	68,89,89	1.41	9 (13%)
2	GDU	Н	500	-	35,38,38	2.67	4 (11%)	53,58,58	<mark>3.58</mark>	16 (30%)
2	GDU	Ι	500	-	35,38,38	2.67	4 (11%)	53,58,58	<mark>3.60</mark>	14 (26%)
3	FAD	F	450	-	53,58,58	1.30	5 (9%)	68,89,89	1.41	9 (13%)
3	FAD	J	450	-	53,58,58	1.30	5 (9%)	68,89,89	1.41	9 (13%)
2	GDU	А	500	-	35,38,38	2.71	4 (11%)	53,58,58	<mark>3.63</mark>	19 (35%)
4	FDA	G	1385	-	52,58,58	2.98	25 (48%)	60,89,89	2.17	21 (35%)
2	GDU	F	500	-	35,38,38	2.66	4 (11%)	53,58,58	3.61	15 (28%)
3	FAD	Е	450	-	$53,\!58,\!58$	1.30	5 (9%)	68,89,89	1.41	9 (13%)
4	FDA	В	1385	-	52,58,58	2.92	24 (46%)	60,89,89	2.21	19 (31%)
3	FAD	Н	450	-	$53,\!58,\!58$	1.30	5 (9%)	68,89,89	1.40	9 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	\mathbf{Res}	Link	Chirals	Torsions	Rings
2	GDU	G	500	-	-	10/23/59/59	0/3/3/3
3	FAD	А	450	-	-	1/30/50/50	0/6/6/6
2	GDU	J	500	-	-	8/23/59/59	0/3/3/3
4	FDA	D	1385	-	4/4/9/9	5/30/50/50	0/6/6/6
2	GDU	Е	500	-	-	6/23/59/59	0/3/3/3



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FAD	С	450	-	-	1/30/50/50	0/6/6/6
2	GDU	В	500	-	-	9/23/59/59	0/3/3/3
2	GDU	D	500	-	-	10/23/59/59	0/3/3/3
2	GDU	С	500	-	-	11/23/59/59	0/3/3/3
3	FAD	Ι	450	-	-	1/30/50/50	0/6/6/6
2	GDU	Н	500	-	-	8/23/59/59	0/3/3/3
2	GDU	Ι	500	-	-	8/23/59/59	0/3/3/3
3	FAD	F	450	-	-	1/30/50/50	0/6/6/6
3	FAD	J	450	-	-	1/30/50/50	0/6/6/6
2	GDU	А	500	-	-	4/23/59/59	0/3/3/3
4	FDA	G	1385	-	2/2/9/9	6/30/50/50	0/6/6/6
2	GDU	F	500	-	-	9/23/59/59	0/3/3/3
3	FAD	Е	450	-	-	1/30/50/50	0/6/6/6
4	FDA	В	1385	-	2/2/9/9	2/30/50/50	0/6/6/6
3	FAD	Н	450	-	-	1/30/50/50	0/6/6/6

The worst 5 of 139 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms		Observed(Å)	Ideal(Å)
2	D	500	GDU	O2-C2	11.23	1.43	1.23
2	F	500	GDU	O2-C2	11.22	1.43	1.23
2	Н	500	GDU	O2-C2	11.22	1.43	1.23
2	А	500	GDU	O2-C2	11.19	1.43	1.23
2	В	500	GDU	O2-C2	11.18	1.43	1.23

The worst 5 of 283 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	500	GDU	N3-C2-N1	16.32	136.55	114.89
2	J	500	GDU	N3-C2-N1	16.03	136.18	114.89
2	Ι	500	GDU	N3-C2-N1	15.95	136.07	114.89
2	В	500	GDU	N3-C2-N1	15.94	136.05	114.89
2	С	500	GDU	N3-C2-N1	15.93	136.03	114.89

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	В	1385	FDA	C2B
4	В	1385	FDA	C1B
4	D	1385	FDA	C4B



Continued from previous page...

Mol	Chain	Res	Type	Atom
4	D	1385	FDA	C2B
4	D	1385	FDA	C3B

5 of 103 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	500	GDU	O5'-C1'-O3B-PB
2	В	500	GDU	C1'-O3B-PB-O2B
2	В	500	GDU	O5'-C1'-O3B-PB
2	С	500	GDU	O4D-C4D-C5D-O5D
2	С	500	GDU	C5D-O5D-PA-O1A

There are no ring outliers.

19 monomers are involved in 48 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	500	GDU	2	0
3	А	450	FAD	4	0
4	D	1385	FDA	3	0
2	Е	500	GDU	2	0
3	С	450	FAD	4	0
2	В	500	GDU	4	0
2	D	500	GDU	6	0
2	С	500	GDU	7	0
3	Ι	450	FAD	2	0
2	Н	500	GDU	1	0
2	Ι	500	GDU	2	0
3	F	450	FAD	3	0
3	J	450	FAD	3	0
2	А	500	GDU	4	0
4	G	1385	FDA	4	0
2	F	500	GDU	3	0
3	Е	450	FAD	2	0
4	В	1385	FDA	2	0
3	Н	450	FAD	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be



highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	365/397~(91%)	-0.12	6 (1%) 72 70	35, 45, 65, 109	1 (0%)
1	В	361/397~(90%)	-0.01	13 (3%) 42 42	36, 48, 66, 98	0
1	С	348/397~(87%)	0.40	26 (7%) 14 13	20, 55, 89, 122	1 (0%)
1	D	360/397~(90%)	0.18	15 (4%) 36 35	37, 53, 77, 102	1 (0%)
1	Е	360/397~(90%)	0.07	13 (3%) 42 42	40, 55, 76, 101	1 (0%)
1	F	360/397~(90%)	0.00	13 (3%) 42 42	37, 48, 73, 97	1 (0%)
1	G	360/397~(90%)	-0.06	1 (0%) 94 93	35, 50, 69, 102	0
1	Н	361/397~(90%)	-0.10	2 (0%) 89 88	37, 50, 70, 89	1 (0%)
1	Ι	361/397~(90%)	-0.09	6 (1%) 70 68	38, 47, 65, 92	1 (0%)
1	J	359/397~(90%)	0.03	8 (2%) 62 60	36, 50, 74, 95	1 (0%)
All	All	3595/3970 (90%)	0.03	103 (2%) 51 50	20, 50, 75, 122	8 (0%)

The worst 5 of 103 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	С	359	VAL	5.3
1	D	154	GLU	5.3
1	F	247	ALA	5.0
1	G	154	GLU	5.0
1	F	154	GLU	4.8

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B -factors($Å^2$)	Q<0.9
3	FAD	С	450	53/53	0.92	0.17	20,20,20,20	0
3	FAD	Ι	450	53/53	0.92	0.14	20,20,20,20	0
3	FAD	F	450	53/53	0.93	0.14	20,20,20,20	0
3	FAD	Н	450	53/53	0.93	0.14	20,20,20,20	0
2	GDU	С	500	36/36	0.93	0.18	36,48,61,68	0
4	FDA	D	1385	53/53	0.93	0.20	42,53,66,71	0
2	GDU	Н	500	36/36	0.94	0.17	39,49,64,70	0
3	FAD	J	450	53/53	0.94	0.15	20,20,20,20	0
3	FAD	Е	450	53/53	0.94	0.16	20,20,20,20	0
2	GDU	J	500	36/36	0.95	0.19	40,49,61,62	0
2	GDU	Е	500	36/36	0.95	0.19	45,56,64,67	0
2	GDU	А	500	36/36	0.95	0.16	35,47,58,66	0
2	GDU	Ι	500	36/36	0.95	0.18	40,49,64,65	0
2	GDU	F	500	36/36	0.96	0.14	39,48,58,61	0
3	FAD	А	450	53/53	0.96	0.10	20,20,20,20	0
4	FDA	В	1385	53/53	0.96	0.16	42,49,55,57	0
2	GDU	G	500	36/36	0.96	0.15	41,49,61,69	0
4	FDA	G	1385	53/53	0.96	0.14	41,53,59,60	0
2	GDU	В	500	36/36	0.97	0.15	43,50,59,64	0
2	GDU	D	500	36/36	0.97	0.15	42,50,56,60	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

















































6.5 Other polymers (i)

There are no such residues in this entry.

