



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 20, 2023 – 03:42 PM EDT

PDB ID : 5HF6  
Title : Crystal structure of human acetylcholinesterase in complex with paraoxon in the aged state  
Authors : Franklin, M.F.; Rudolph, M.J.; Ginter, C.; Cassidy, M.S.; Cheung, J.  
Deposited on : 2016-01-06  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	542	 90% 7% 3% 0% 0%
1	B	542	 90% 9% 1% 1% 0%
2	C	3	 33% 67%
2	D	3	 33% 33% 33%

## 2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 9023 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Acetylcholinesterase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	528	Total	C	N	O	S	0	5	0
			4158	2668	730	747	13			
1	B	538	Total	C	N	O	S	0	3	0
			4209	2699	739	758	13			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	3	Total	C	N	O	0	0	0
			38	22	2	14			
2	D	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 3 is ETHYL DIHYDROGEN PHOSPHATE (three-letter code: EFS) (formula: C<sub>2</sub>H<sub>7</sub>O<sub>4</sub>P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	O	P	0	0
			6	2	3	1		
3	B	1	Total	C	O	P	0	0
			6	2	3	1		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	296	Total	O	0	0
			296	296		
6	B	242	Total	O	0	0
			242	242		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Acetylcholinesterase




- Molecule 1: Acetylcholinesterase



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[alpha-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	105.90Å 105.90Å 323.08Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	47.52 – 2.30 47.52 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (47.52-2.30) 99.9 (47.52-2.30)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.42 (at 2.29Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.174 , 0.206 0.179 , 0.208	Depositor DCC
$R_{free}$ test set	4741 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.0	Xtrriage
Anisotropy	0.562	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 39.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	9023	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.20% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, EFS, FUC, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/4287	0.57	0/5858
1	B	0.47	0/4341	0.60	0/5934
All	All	0.46	0/8628	0.59	0/11792

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	257	CYS	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4158	0	4035	29	0
1	B	4209	0	4085	29	0
2	C	38	0	34	1	0
2	D	38	0	34	1	0
3	A	6	0	5	0	0
3	B	6	0	5	0	0
4	A	8	0	12	0	0
4	B	8	0	12	0	0
5	B	14	0	13	0	0
6	A	296	0	0	7	0
6	B	242	0	0	8	0
All	All	9023	0	8235	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:527:GLN:NE2	6:A:701:HOH:O	2.10	0.84
1:A:18[A]:ARG:NH2	6:A:702:HOH:O	2.18	0.74
1:B:4:GLU:HG2	1:B:9:LEU:HD11	1.71	0.72
1:B:527:GLN:NE2	6:B:704:HOH:O	2.25	0.69
1:A:18[A]:ARG:NH1	6:A:703:HOH:O	2.18	0.68
1:B:304:ASP:OD1	6:B:701:HOH:O	2.13	0.66
1:A:286:TRP:HB3	1:A:296:ARG:NH2	2.12	0.65
1:A:5:ASP:N	6:A:706:HOH:O	2.31	0.63
1:A:493:ARG:NH2	6:A:707:HOH:O	2.33	0.61
1:A:408:VAL:HG11	1:A:525:ARG:HG3	1.85	0.57
1:B:113:PRO:HG2	1:B:485:ARG:HG2	1.85	0.57
1:A:30:SER:HB2	1:A:103:THR:HG22	1.89	0.55
1:B:107:ARG:NE	6:B:705:HOH:O	2.33	0.53
1:A:200:PHE:CB	1:A:226:VAL:HB	2.39	0.53
1:B:328:VAL:O	1:B:427:ALA:HA	2.09	0.52
1:A:380:LEU:HD22	1:B:535:PHE:CB	2.40	0.52
1:B:306:ASP:HB3	6:B:769:HOH:O	2.09	0.52
1:B:373:LEU:HG	1:B:539:LEU:HD11	1.94	0.50
1:B:491:GLU:HB2	1:B:494:ASP:HB3	1.94	0.50
1:A:227:LEU:HB2	1:A:328:VAL:HG12	1.93	0.49
1:B:408:VAL:HG11	1:B:525:ARG:HG3	1.94	0.49
1:B:468:GLU:HG3	6:B:842:HOH:O	2.12	0.49
1:B:200:PHE:CB	1:B:226:VAL:HB	2.43	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:291:GLN:OE1	1:A:369:GLN:NE2	2.46	0.48
1:A:286:TRP:HB3	1:A:296:ARG:CZ	2.43	0.48
1:B:332:LYS:HE3	6:B:855:HOH:O	2.13	0.47
1:A:328:VAL:O	1:A:427:ALA:HA	2.15	0.47
1:B:68:VAL:HG23	1:B:90:ARG:HB2	1.96	0.47
1:B:329:GLY:HA3	1:B:428:TYR:CZ	2.50	0.47
1:A:291:GLN:HG2	1:A:292:GLU:N	2.29	0.46
1:A:329:GLY:HA3	1:A:428:TYR:CE2	2.52	0.45
1:A:433:ARG:CZ	1:A:437:LEU:HD23	2.47	0.45
1:A:113:PRO:HG3	1:A:485:ARG:HG2	1.98	0.45
1:A:124:TYR:HE2	1:A:297:PHE:CE1	2.35	0.45
1:B:490:ASN:O	6:B:702:HOH:O	2.21	0.45
1:B:458:PRO:HA	1:B:465:TYR:CD1	2.51	0.44
1:B:293:SER:HA	1:B:365:VAL:HG13	1.99	0.44
1:A:143:ARG:NH1	6:A:718:HOH:O	2.46	0.44
1:A:380:LEU:HD22	1:B:535:PHE:HB2	1.99	0.44
1:A:304:ASP:OD1	6:A:704:HOH:O	2.21	0.43
1:B:492:PRO:HG3	6:B:702:HOH:O	2.17	0.43
1:B:200:PHE:HB3	1:B:226:VAL:HB	2.01	0.43
1:A:347:SER:HB2	2:C:1:NAG:H62	2.00	0.43
1:B:24:THR:OG1	1:B:140:GLN:HG3	2.19	0.43
1:B:30:SER:HB2	1:B:103:THR:HG22	2.01	0.43
1:A:329:GLY:HA3	1:A:428:TYR:CZ	2.54	0.42
1:A:200:PHE:HB3	1:A:226:VAL:HB	2.00	0.42
1:A:17:LEU:HD23	1:A:60:VAL:HB	2.01	0.41
1:B:347:SER:HB2	2:D:1:NAG:H62	2.01	0.41
1:A:536:LEU:HD12	1:A:536:LEU:HA	1.89	0.41
1:B:339:LEU:HD23	1:B:339:LEU:HA	1.87	0.41
1:B:297:PHE:CD1	1:B:297:PHE:N	2.88	0.41
1:A:366:GLY:HA3	1:A:402:VAL:HG22	2.04	0.40
1:B:339:LEU:HD13	1:B:346:PHE:CZ	2.56	0.40
1:A:296:ARG:C	1:A:297:PHE:HD1	2.25	0.40
1:B:329:GLY:HA3	1:B:428:TYR:CE2	2.57	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	527/542 (97%)	514 (98%)	13 (2%)	0	100	100
1	B	537/542 (99%)	516 (96%)	18 (3%)	3 (1%)	25	31
All	All	1064/1084 (98%)	1030 (97%)	31 (3%)	3 (0%)	41	50

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	258	PRO
1	B	296	ARG
1	B	495	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	434/437 (99%)	429 (99%)	5 (1%)	71	84
1	B	439/437 (100%)	431 (98%)	8 (2%)	59	75
All	All	873/874 (100%)	860 (98%)	13 (2%)	65	79

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	LEU
1	A	165	ARG
1	A	200	PHE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	288	VAL
1	A	396	GLU
1	B	146	LEU
1	B	200	PHE
1	B	258	PRO
1	B	288	VAL
1	B	356	ARG
1	B	396	GLU
1	B	493	ARG
1	B	540	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	291	GLN
1	A	369	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	NAG	C	1	2,1	14,14,15	0.37	0	17,19,21	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	2	2	14,14,15	0.24	0	17,19,21	0.39	0
2	FUC	C	3	2	10,10,11	1.31	1 (10%)	14,14,16	1.06	1 (7%)
2	NAG	D	1	2,1	14,14,15	0.28	0	17,19,21	1.00	1 (5%)
2	NAG	D	2	2	14,14,15	0.40	0	17,19,21	0.37	0
2	FUC	D	3	2	10,10,11	1.27	2 (20%)	14,14,16	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	FUC	C	3	2	-	-	0/1/1/1
2	NAG	D	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	0/6/23/26	0/1/1/1
2	FUC	D	3	2	-	-	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	FUC	C2-C3	2.66	1.56	1.52
2	D	3	FUC	C2-C3	2.50	1.56	1.52
2	D	3	FUC	O5-C5	2.14	1.48	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	NAG	C1-O5-C5	3.41	116.81	112.19
2	C	3	FUC	C1-O5-C5	2.11	117.56	112.78

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	D	1	NAG	C8-C7-N2-C2

*Continued on next page...*

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	NAG	1	0
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EFS	A	604	1	2,5,6	3.94	1 (50%)	1,5,8	0.03	0
3	EFS	B	605	1	2,5,6	4.00	1 (50%)	1,5,8	0.11	0
4	EDO	A	605	-	3,3,3	0.48	0	2,2,2	0.36	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	EDO	A	606	-	3,3,3	0.55	0	2,2,2	0.16	0
4	EDO	B	606	-	3,3,3	0.58	0	2,2,2	0.15	0
5	NAG	B	604	1	14,14,15	0.49	0	17,19,21	0.62	0
4	EDO	B	607	-	3,3,3	0.57	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EFS	A	604	1	-	0/1/3/4	-
3	EFS	B	605	1	-	0/1/3/4	-
4	EDO	A	605	-	-	0/1/1/1	-
4	EDO	A	606	-	-	0/1/1/1	-
4	EDO	B	606	-	-	0/1/1/1	-
5	NAG	B	604	1	-	2/6/23/26	0/1/1/1
4	EDO	B	607	-	-	1/1/1/1	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	605	EFS	O2-C11	-5.58	1.31	1.44
3	A	604	EFS	O2-C11	-5.46	1.31	1.44

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	604	NAG	C4-C5-C6-O6
5	B	604	NAG	O5-C5-C6-O6
4	B	607	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.



## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	528/542 (97%)	-0.15	8 (1%) 73 79	33, 50, 76, 101	0
1	B	538/542 (99%)	-0.08	15 (2%) 53 60	33, 52, 83, 123	0
All	All	1066/1084 (98%)	-0.12	23 (2%) 62 69	33, 51, 80, 123	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	493	ARG	10.1
1	B	262	THR	5.4
1	B	496	LYS	5.3
1	B	495	PRO	5.3
1	B	497	ALA	4.8
1	B	540	LEU	4.6
1	A	497	ALA	4.5
1	B	494	ASP	4.3
1	B	286	TRP	3.6
1	B	290	PRO	3.6
1	A	290	PRO	3.4
1	B	541	SER	3.2
1	B	492	PRO	3.2
1	B	291	GLN	3.1
1	A	540	LEU	2.9
1	A	539	LEU	2.5
1	B	537	PRO	2.5
1	A	373	LEU	2.4
1	B	165	ARG	2.2
1	B	259	PRO	2.2
1	A	286	TRP	2.1
1	A	165	ARG	2.1
1	A	541	SER	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	C	2	14/15	0.83	0.40	82,108,118,122	0
2	FUC	C	3	10/11	0.88	0.37	71,99,105,109	0
2	NAG	D	1	14/15	0.90	0.26	80,92,103,107	0
2	NAG	D	2	14/15	0.90	0.33	83,101,115,120	0
2	NAG	C	1	14/15	0.92	0.28	70,83,100,107	0
2	FUC	D	3	10/11	0.94	0.38	85,97,105,113	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	NAG	B	604	14/15	0.83	0.27	82,99,110,111	0
4	EDO	B	607	4/4	0.96	0.12	49,50,51,56	0
4	EDO	A	606	4/4	0.96	0.12	48,52,55,61	0
4	EDO	B	606	4/4	0.97	0.15	51,53,53,54	0
3	EFS	B	605	6/7	0.98	0.23	39,48,57,62	0
4	EDO	A	605	4/4	0.98	0.15	48,49,52,57	0
3	EFS	A	604	6/7	0.98	0.21	44,50,59,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.