

Full wwPDB X-ray Structure Validation Report (i)

Jan 7, 2024 - 02:54 am GMT

PDB ID : 6HGQ

Title: Crystal Structure of Human APRT wild type in complex with Hypoxanthine,

PRPP and Mg2+

Authors: Nioche, P.; Huyet, J.; Ozeir, M.

Deposited on : 2018-08-23

Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	179	82%	18%
1	В	179	78%	19% •
1	С	179	80%	19% •
1	D	179	88%	11% •



2 Entry composition (i)


There are 6 unique types of molecules in this entry. The entry contains 6094 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Adenine phosphoribosyltransferase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	179	Total	С	N	О	S	0	9	0
1	A	119	1409	908	238	258	5	0	9	0
1	С	179	Total	С	N	О	S	0	10	0
1		119	1421	914	240	262	5	0	10	0
1	D	177	Total	С	N	О	S	0	8	0
1	ע	111	1394	897	237	255	5	0	8	0
1	D	174	Total	С	N	О	S	0	10	0
1	Б	174	1387	893	234	255	5	U	10	U

• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0

• Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Mg 1 1	0	0
3	С	1	Total Mg 1 1	0	0
3	D	1	Total Mg 1 1	0	0
3	В	1	$\begin{array}{cc} \text{Total} & \text{Mg} \\ 1 & 1 \end{array}$	0	0


• Molecule 4 is 1-O-pyrophosphono-5-O-phosphono-alpha-D-ribofuranose (three-letter code: PRP) (formula: $C_5H_{13}O_{14}P_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	Δ	1	Total	С	О	Р	0	0	
4	11	1	22	5	14	3	O	O	
1	C	1	Total (al C O P		0			
4		1	22	5	14	3	U		
1	D	1	Total	\overline{C}	О	Р	0	0	
4	D	1	22	5	14	3	U	U	
1	D	1	Total	$\overline{\mathrm{C}}$	О	Р	0	0	
4	Б	1	22	5	14	3	U	U	

• Molecule 5 is HYPOXANTHINE (three-letter code: HPA) (formula: $C_5H_4N_4O$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N O 10 5 4 1	0	0
5	С	1	Total C N O 10 5 4 1	0	0
5	D	1	Total C N O 10 5 4 1	0	0
5	В	1	Total C N O 10 5 4 1	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	105	Total O 105 105	0	0
6	С	100	Total O 100 100	0	0
6	D	79	Total O 79 79	0	0
6	В	61	Total O 61 61	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Adenine phosphoribosyltransferase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	49.06Å 49.80Å 71.81Å	Depositor
a, b, c, α , β , γ	90.08° 93.22° 102.31°	Depositor
Resolution (Å)	71.69 - 1.90	Depositor
Resolution (A)	40.91 - 1.90	EDS
% Data completeness	90.0 (71.69-1.90)	Depositor
(in resolution range)	90.0 (40.91-1.90)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) > 1$	4.29 (at 1.89Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
Ρ. Р.	0.175 , 0.214	Depositor
R, R_{free}	0.186 , 0.222	DCC
R_{free} test set	2388 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	17.1	Xtriage
Anisotropy	0.967	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.41 , 54.4	EDS
L-test for twinning ²	$< L >=0.48, < L^2>=0.30$	Xtriage
Estimated twinning fraction	0.022 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6094	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 7.70% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: PRP, HPA, GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
MIOI		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.30	0/1469	0.50	0/1988
1	В	0.28	0/1446	0.49	0/1953
1	С	0.30	0/1478	0.51	0/2000
1	D	0.28	0/1448	0.49	0/1959
All	All	0.29	0/5841	0.50	0/7900

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1409	0	1460	23	0
1	В	1387	0	1429	23	0
1	С	1421	0	1464	26	0
1	D	1394	0	1439	12	0
2	A	6	0	8	2	0
3	A	1	0	0	0	0
3	В	1	0	0	0	0
3	С	1	0	0	0	0
3	D	1	0	0	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
4	A	22	0	6	0	0
4	В	22	0	6	0	0
4	С	22	0	6	1	0
4	D	22	0	6	1	0
5	A	10	0	4	1	0
5	В	10	0	4	1	0
5	С	10	0	4	1	0
5	D	10	0	4	1	0
6	A	105	0	0	2	0
6	В	61	0	0	1	0
6	С	100	0	0	0	0
6	D	79	0	0	2	0
All	All	6094	0	5840	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance $(Å)$	Clash overlap (Å)
1:A:27:ARG:H	5:A:204:HPA:HN1	1.28	0.81
1:A:130:LEU:HD13	1:A:136[A]:MET:SD	2.26	0.75
1:C:107:LYS:C	1:C:107:LYS:HD3	2.10	0.72
1:B:30[B]:SER:OG	1:B:31:PRO:HD3	1.89	0.71
1:D:30[B]:SER:OG	1:D:31:PRO:HD3	1.93	0.68
1:C:75:LEU:HD23	1:C:126[B]:VAL:HG21	1.75	0.68
1:C:27:ARG:H	5:C:202:HPA:HN1	1.41	0.67
1:D:57:ARG:O	1:D:122:ARG:NE	2.26	0.66
1:A:75:LEU:HD23	1:A:126[B]:VAL:HG21	1.78	0.66
1:D:27:ARG:H	5:D:202:HPA:HN1	1.43	0.66
1:B:75:LEU:HD23	1:B:126[B]:VAL:HG21	1.79	0.64
1:C:14:ARG:HG3	1:C:31:PRO:HG2	1.83	0.61
1:A:78:GLU:OE1	6:A:301:HOH:O	2.16	0.61
1:C:30[B]:SER:OG	1:C:31:PRO:HD3	2.02	0.60
1:A:14:ARG:HG3	1:A:31:PRO:HG2	1.83	0.60
1:D:75:LEU:HD23	1:D:126[B]:VAL:HG21	1.84	0.59
1:B:92:LEU:HD12	1:B:111[B]:GLU:HG3	1.85	0.59
1:A:6[B]:LEU:HD23	1:A:45:LEU:HD13	1.86	0.57
1:C:130:LEU:HD13	1:C:136[A]:MET:SD	2.44	0.57
1:A:30[B]:SER:OG	1:A:31:PRO:HD3	2.06	0.56
1:C:97:LEU:HD22	1:C:114:LYS:HG2	1.88	0.55



 $Continued\ from\ previous\ page...$

Atom 1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	$\text{overlap } (\mathring{\mathbf{A}})$
1:B:72:GLY:HA2	1:B:126[A]:VAL:HG11	1.89	0.55
1:C:107:LYS:C	1:C:107:LYS:CD	2.74	0.55
1:B:101:TYR:CE1	1:B:110:LEU:HG	2.41	0.55
1:B:27:ARG:H	5:B:202:HPA:HN1	1.54	0.54
1:C:107:LYS:HE2	1:C:108:ALA:N	2.23	0.53
1:C:89:ARG:HD3	1:C:109:GLU:OE2	2.10	0.52
1:C:161:SER:HB3	1:C:180:GLU:HG3	1.90	0.52
1:A:101:TYR:CE1	1:A:110:LEU:HG	2.45	0.51
1:B:101:TYR:HE1	1:B:110:LEU:HG	1.77	0.50
1:B:54:HIS:NE2	1:B:152:GLU:OE1	2.34	0.49
1:A:127:ASP:O	1:A:155:SER:HA	2.13	0.49
1:C:107:LYS:HD3	1:C:107:LYS:O	2.12	0.48
1:B:96:THR:HB	1:B:111[A]:GLU:HG2	1.95	0.48
1:C:130:LEU:HD11	1:C:133:GLY:HA2	1.96	0.48
1:D:136[A]:MET:HG3	6:D:326:HOH:O	2.13	0.47
1:C:97:LEU:CD2	1:C:114:LYS:HG2	2.43	0.47
1:B:126[A]:VAL:HG23	1:B:154:VAL:HG23	1.95	0.47
1:C:88[B]:LYS:NZ	4:C:201:PRP:O1B	2.38	0.47
1:C:18:ASP:N	1:C:24:VAL:O	2.48	0.46
1:B:88[B]:LYS:HD2	6:B:317:HOH:O	2.14	0.46
1:A:98:TRP:CH2	1:C:3:ASP:HB2	2.50	0.46
1:C:127:ASP:O	1:C:155:SER:HA	2.16	0.46
1:D:161:SER:HB3	1:D:180:GLU:HG2	1.98	0.46
1:B:65:ASP:HA	1:B:66[B]:SER:HA	1.70	0.45
1:A:65:ASP:HA	1:A:66[A]:SER:HA	1.68	0.45
1:A:159:LEU:HD13	2:A:201:GOL:H12	1.98	0.45
1:C:130:LEU:CD1	1:C:133:GLY:HA2	2.47	0.45
1:C:96:THR:HB	1:C:111:GLU:HG2	1.98	0.45
1:A:97:LEU:HD21	1:A:114:LYS:HE2	1.99	0.45
1:C:65:ASP:HA	1:C:66[B]:SER:HA	1.69	0.44
1:B:67:ARG:HD2	1:B:128:ASP:OD2	2.17	0.44
1:B:96:THR:OG1	1:B:111[A]:GLU:OE1	2.35	0.44
1:C:101:TYR:CE1	1:C:110:LEU:HG	2.53	0.44
1:B:130:LEU:HD13	1:B:136[A]:MET:SD	2.57	0.44
1:C:102:SER:O	1:C:134:GLY:HA3	2.18	0.43
1:A:65:ASP:HB2	1:A:87:ARG:HA	2.01	0.42
1:D:54:HIS:HE2	1:D:152:GLU:HB2	1.84	0.42
1:A:180:GLU:O	2:A:201:GOL:C3	2.68	0.42
1:D:101:TYR:CE1	1:D:110:LEU:HG	2.53	0.42
1:B:127:ASP:O	1:B:155:SER:HA	2.19	0.42
1:B:18:ASP:N	1:B:24:VAL:O	2.43	0.42



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:SER:O	1:A:134:GLY:HA3	2.20	0.42
1:D:88[B]:LYS:NZ	4:D:201:PRP:O1B	2.37	0.42
1:A:167:LYS:HE2	6:A:316:HOH:O	2.20	0.41
1:D:97:LEU:HD22	6:D:349:HOH:O	2.19	0.41
1:B:65:ASP:HB2	1:B:87:ARG:HA	2.01	0.41
1:A:72:GLY:HA2	1:A:126[A]:VAL:HG11	2.02	0.41
1:C:65:ASP:HB2	1:C:87:ARG:HA	2.03	0.41
1:D:65:ASP:HA	1:D:66[A]:SER:HA	1.72	0.41
1:B:167:LYS:HA	1:B:167:LYS:HD3	1.94	0.40
1:A:88:LYS:HD3	1:A:88:LYS:HA	1.91	0.40
1:B:76:ALA:HB1	1:B:81:LEU:O	2.22	0.40
1:B:126[B]:VAL:HG22	1:B:154:VAL:CG2	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	186/179 (104%)	184 (99%)	2 (1%)	0	100	100
1	В	180/179 (101%)	177 (98%)	3 (2%)	0	100	100
1	С	187/179 (104%)	185 (99%)	2 (1%)	0	100	100
1	D	183/179 (102%)	182 (100%)	1 (0%)	0	100	100
All	All	736/716 (103%)	728 (99%)	8 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar



resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	155/146 (106%)	151 (97%)	4 (3%)	46 39
1	В	153/146 (105%)	150 (98%)	3 (2%)	55 51
1	\mathbf{C}	156/146 (107%)	154 (99%)	2 (1%)	69 68
1	D	153/146 (105%)	152 (99%)	1 (1%)	84 84
All	All	617/584 (106%)	607 (98%)	10 (2%)	62 60

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91[A]	LYS
1	A	91[B]	LYS
1	A	103	LEU
1	A	107	LYS
1	С	103	LEU
1	С	107	LYS
1	D	103	LEU
1	В	20	PRO
1	В	103	LEU
1	В	122	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trno	Chain	Res	Link	Вс	ond leng	ths	В	ond ang	gles
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	201	-	5,5,5	0.25	0	5,5,5	0.17	0
4	PRP	D	201	3	19,22,22	0.69	0	33,35,35	1.14	1 (3%)
5	HPA	D	202	-	8,11,11	1.93	3 (37%)	5,15,15	3.66	5 (100%)
5	HPA	С	202	-	8,11,11	1.94	2 (25%)	5,15,15	3.77	5 (100%)
5	HPA	A	204	-	8,11,11	1.81	2 (25%)	5,15,15	3.96	5 (100%)
4	PRP	В	201	3	19,22,22	0.70	0	33,35,35	1.16	2 (6%)
5	HPA	В	202	-	8,11,11	1.99	2 (25%)	5,15,15	3.82	5 (100%)
4	PRP	С	201	3	19,22,22	0.65	0	33,35,35	1.04	0
4	PRP	A	203	3	19,22,22	0.69	0	33,35,35	1.05	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	201	_	-	2/4/4/4	-
4	PRP	D	201	3	-	4/16/33/33	0/1/1/1
5	HPA	D	202	-	-	-	0/2/2/2
5	HPA	С	202	-	-	-	0/2/2/2
5	HPA	A	204	_	-	-	0/2/2/2
4	PRP	В	201	3	-	4/16/33/33	0/1/1/1
5	HPA	В	202	-	-	-	0/2/2/2
4	PRP	С	201	3	-	5/16/33/33	0/1/1/1
4	PRP	A	203	3	-	5/16/33/33	0/1/1/1

All (9) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
5	В	202	HPA	C5-C6	4.33	1.48	1.41
5	С	202	HPA	C5-C6	4.19	1.48	1.41
5	D	202	HPA	C5-C6	4.05	1.48	1.41
5	A	204	HPA	C5-C6	3.88	1.48	1.41
5	С	202	HPA	C5-C4	2.20	1.46	1.40
5	D	202	HPA	C5-C4	2.17	1.46	1.40
5	В	202	HPA	C5-C4	2.13	1.46	1.40
5	A	204	HPA	C5-C4	2.12	1.46	1.40
5	D	202	HPA	C2-N3	2.05	1.35	1.32

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
5	A	204	HPA	C4-C5-C6	-5.46	115.58	120.80
5	С	202	HPA	C4-C5-C6	-5.17	115.86	120.80
5	В	202	HPA	C4-C5-C6	-4.92	116.11	120.80
5	D	202	HPA	C4-C5-C6	-4.76	116.25	120.80
5	A	204	HPA	C2-N1-C6	4.67	123.70	115.88
5	D	202	HPA	C2-N1-C6	4.29	123.07	115.88
5	В	202	HPA	C2-N1-C6	4.27	123.02	115.88
5	С	202	HPA	C2-N1-C6	4.14	122.81	115.88
5	В	202	HPA	C4-C5-N7	-3.70	105.55	109.40
5	A	204	HPA	N3-C2-N1	-3.69	122.91	128.68
5	В	202	HPA	N3-C2-N1	-3.42	123.34	128.68
5	С	202	HPA	N3-C2-N1	-3.35	123.44	128.68
5	С	202	HPA	C4-C5-N7	-3.31	105.95	109.40
5	D	202	HPA	N3-C2-N1	-3.30	123.52	128.68
4	D	201	PRP	O3A-PA-O1	3.28	109.09	102.48
5	D	202	HPA	C4-C5-N7	-3.23	106.03	109.40
4	В	201	PRP	O3A-PA-O1	3.05	108.63	102.48
5	A	204	HPA	C4-C5-N7	-2.93	106.34	109.40
4	В	201	PRP	O4-C1-O1	2.36	113.86	109.18
4	A	203	PRP	O3A-PA-O1	2.33	107.18	102.48
5	С	202	HPA	C2-N3-C4	2.27	118.77	113.45
5	В	202	HPA	C2-N3-C4	2.27	118.77	113.45
5	A	204	HPA	C2-N3-C4	2.13	118.45	113.45
5	D	202	HPA	C2-N3-C4	2.09	118.35	113.45

There are no chirality outliers.

All (20) torsion outliers are listed below:

\mathbf{Mol}	Chain	Res	Type	Atoms
2	A	201	GOL	C1-C2-C3-O3



Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	203	PRP	PB-O3A-PA-O1
4	D	201	PRP	PB-O3A-PA-O1
4	В	201	PRP	PB-O3A-PA-O1
4	D	201	PRP	C1-O1-PA-O3A
4	В	201	PRP	C1-O1-PA-O3A
2	A	201	GOL	O2-C2-C3-O3
4	С	201	PRP	C1-O1-PA-O3A
4	В	201	PRP	C1-O1-PA-O2A
4	A	203	PRP	C4-C5-O5-P
4	С	201	PRP	PB-O3A-PA-O1
4	A	203	PRP	C1-O1-PA-O3A
4	A	203	PRP	C1-O1-PA-O1A
4	С	201	PRP	C1-O1-PA-O1A
4	В	201	PRP	C4-C5-O5-P
4	С	201	PRP	C4-C5-O5-P
4	D	201	PRP	C1-O1-PA-O2A
4	D	201	PRP	C4-C5-O5-P
4	A	203	PRP	C1-O1-PA-O2A
4	С	201	PRP	C1-O1-PA-O2A

There are no ring outliers.


7 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	GOL	2	0
4	D	201	PRP	1	0
5	D	202	HPA	1	0
5	С	202	HPA	1	0
5	A	204	HPA	1	0
5	В	202	HPA	1	0
4	С	201	PRP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.




The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q<0.9
1	A	179/179 (100%)	0.02	0 100 100	9, 18, 31, 46	0
1	В	174/179 (97%)	0.24	8 (4%) 32 35	11, 21, 40, 53	0
1	С	179/179 (100%)	0.06	1 (0%) 89 90	10, 17, 33, 47	0
1	D	177/179 (98%)	0.21	4 (2%) 60 63	10, 20, 36, 51	0
All	All	709/716 (99%)	0.13	13 (1%) 68 71	9, 19, 36, 53	0

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	105	TYR	3.6
1	D	180	GLU	3.3
1	В	24	VAL	3.2
1	D	23	GLY	3.0
1	В	107	LYS	2.8
1	В	17	PRO	2.7
1	В	180	GLU	2.6
1	D	4	SER	2.6
1	С	2	ALA	2.5
1	В	103	LEU	2.4
1	D	105	TYR	2.2
1	В	106	GLY	2.1
1	В	57	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	GOL	A	201	6/6	0.84	0.17	26,30,32,35	0
5	HPA	В	202	10/10	0.85	0.25	18,20,21,22	0
5	HPA	A	204	10/10	0.94	0.12	11,12,13,14	0
5	HPA	С	202	10/10	0.94	0.12	13,14,15,15	0
5	HPA	D	202	10/10	0.94	0.15	14,16,17,18	0
3	MG	В	200	1/1	0.94	0.05	14,14,14,14	0
4	PRP	В	201	22/22	0.95	0.11	11,14,16,19	0
4	PRP	С	201	22/22	0.96	0.09	7,9,13,17	0
4	PRP	D	201	22/22	0.97	0.08	10,12,15,19	0
4	PRP	A	203	22/22	0.97	0.08	6,8,14,15	0
3	MG	D	200	1/1	0.97	0.06	10,10,10,10	0
3	MG	A	202	1/1	0.98	0.04	9,9,9,9	0
3	MG	С	200	1/1	0.99	0.04	11,11,11,11	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

























Electron density around PRP B 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive) Electron density around PRP C 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o ext{-}{ m DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around PRP D 201: $2 {\rm mF}_o\text{-}{\rm DF}_c \ ({\rm at}\ 0.7\ {\rm rmsd}) \ {\rm in}\ {\rm gray}$

 ${
m mF}_o{
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around PRP A 203:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















6.5 Other polymers (i)

There are no such residues in this entry.

