

Full wwPDB X-ray Structure Validation Report (i)

Sep 9, 2023 – 06:23 PM EDT

PDB ID : 4HJV

Title: Crystal structure of E. coli MltE with bound bulgecin and murodipeptide

Authors: Fibriansah, G.; Gliubich, F.I.; Thunnissen, A.-M.W.H.

Deposited on : 2012-10-14

Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY\ DIFFRACTION$

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\mathring{A})}) \end{array}$
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	203	5% 85%	6%	9%
1	В	203	87%	•	9%
1	С	203	83%	7%	9%
1	D	203	8%	5%	10%
1	Е	203	7% 85%	6%	9%



 $Continued\ from\ previous\ page...$

Mol	Chain	Length	Quality of chain
2	F	2	100%
2	G	2	100%
2	Н	2	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	В	302	-	-	X	-



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 7814 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Endo-type membrane-bound lytic murein transglycosylase A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	184	Total	С	N	О	S	0	1	0
1	A	104	1415	891	250	267	7	0	1	U
1	В	185	Total	С	N	О	S	0	1	0
1	1 D	100	1425	898	253	268	6	0	1	U
1	С	C 185	Total	С	N	О	S	0	1	0
1			1422	895	252	269	6	U	1	
1	D	182	Total	С	N	О	S	0	1	0
1		102	1397	881	246	263	7	0	1	U
1	Е	185	Total	С	N	О	S	0	0	0
	<u> 1</u> 2	185	1421	895	252	268	6	0	0	U

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P0C960
A	2	ARG	-	expression tag	UNP P0C960
A	3	GLY	-	expression tag	UNP P0C960
A	4	SER	-	expression tag	UNP P0C960
A	5	HIS	-	expression tag	UNP P0C960
A	6	HIS	-	expression tag	UNP P0C960
A	7	HIS	-	expression tag	UNP P0C960
A	8	HIS	-	expression tag	UNP P0C960
A	9	HIS	-	expression tag	UNP P0C960
A	10	HIS	-	expression tag	UNP P0C960
A	11	GLY	-	expression tag	UNP P0C960
A	12	SER	-	expression tag	UNP P0C960
A	13	ALA	-	expression tag	UNP P0C960
A	14	CYS	-	expression tag	UNP P0C960
A	15	GLU	-	expression tag	UNP P0C960
A	16	LEU		expression tag	UNP P0C960
В	1	MET	-	expression tag	UNP P0C960
В	2	ARG	-	expression tag	UNP P0C960
В	3	GLY	-	expression tag	UNP P0C960



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
В	4	SER	-	expression tag	UNP P0C960
В	5	HIS	-	expression tag	UNP P0C960
В	6	HIS	_	expression tag	UNP P0C960
В	7	HIS	-	expression tag	UNP P0C960
В	8	HIS	_	expression tag	UNP P0C960
В	9	HIS	-	expression tag	UNP P0C960
В	10	HIS	-	expression tag	UNP P0C960
В	11	GLY	-	expression tag	UNP P0C960
В	12	SER	-	expression tag	UNP P0C960
В	13	ALA	-	expression tag	UNP P0C960
В	14	CYS	-	expression tag	UNP P0C960
В	15	GLU	-	expression tag	UNP P0C960
В	16	LEU	-	expression tag	UNP P0C960
С	1	MET	-	expression tag	UNP P0C960
С	2	ARG	-	expression tag	UNP P0C960
С	3	GLY	-	expression tag	UNP P0C960
С	4	SER	-	expression tag	UNP P0C960
С	5	HIS	-	expression tag	UNP P0C960
С	6	HIS	-	expression tag	UNP P0C960
С	7	HIS	-	expression tag	UNP P0C960
С	8	HIS	-	expression tag	UNP P0C960
С	9	HIS	-	expression tag	UNP P0C960
С	10	HIS	-	expression tag	UNP P0C960
С	11	GLY	-	expression tag	UNP P0C960
С	12	SER	-	expression tag	UNP P0C960
С	13	ALA	-	expression tag	UNP P0C960
С	14	CYS	-	expression tag	UNP P0C960
С	15	GLU	-	expression tag	UNP P0C960
С	16	LEU	-	expression tag	UNP P0C960
D	1	MET	-	expression tag	UNP P0C960
D	2	ARG	-	expression tag	UNP P0C960
D	3	GLY	-	expression tag	UNP P0C960
D	4	SER	-	expression tag	UNP P0C960
D	5	HIS	-	expression tag	UNP P0C960
D	6	HIS	-	expression tag	UNP P0C960
D	7	HIS	-	expression tag	UNP P0C960
D	8	HIS	-	expression tag	UNP P0C960
D	9	HIS		expression tag	UNP P0C960
D	10	HIS	-	expression tag	UNP P0C960
D	11	GLY	-	expression tag	UNP P0C960
D	12	SER	-	expression tag	UNP P0C960
D	13	ALA		expression tag	UNP P0C960



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	14	CYS	-	expression tag	UNP P0C960
D	15	GLU	-	expression tag	UNP P0C960
D	16	LEU	-	expression tag	UNP P0C960
Е	1	MET	-	expression tag	UNP P0C960
E	2	ARG	-	expression tag	UNP P0C960
Е	3	GLY	-	expression tag	UNP P0C960
E	4	SER	-	expression tag	UNP P0C960
Е	5	HIS	-	expression tag	UNP P0C960
Е	6	HIS	-	expression tag	UNP P0C960
Е	7	HIS	-	expression tag	UNP P0C960
E	8	HIS	-	expression tag	UNP P0C960
Е	9	HIS	-	expression tag	UNP P0C960
Е	10	HIS	-	expression tag	UNP P0C960
E	11	GLY	-	expression tag	UNP P0C960
E	12	SER	-	expression tag	UNP P0C960
Е	13	ALA	-	expression tag	UNP P0C960
Е	14	CYS	-	expression tag	UNP P0C960
Е	15	GLU	-	expression tag	UNP P0C960
Е	16	LEU	-	expression tag	UNP P0C960


 \bullet Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-N-acetyl-beta-muramic acid.



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	Trace		
2	F	9	Total C N O		0	0	0		
	I.	2	34	19	2	13			
9		C 2	Total	С	N	O	0	0	0
2	G		34	19	2	13			
9	П	9	Total	С	N	О	0	0	0
2	П	$1 \mid 2 \mid$	34	19	2	13	0	0	U


• Molecule 3 is 4-O-(4-O-SULFONYL-N-ACETYLGLUCOSAMININYL)-5-METHYLHYDR OXY-L-PROLINE-TAURINE (three-letter code: BLG) (formula: $C_{16}H_{30}N_3O_{14}S_2$).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
2	3 A	1	Total	С	N	О	S	0	0	
3			35	16	3	14	2	0		
3	В	1	Total	С	N	О	S	0	0	
3	3 B	1	35	16	3	14	2	U		
3	C	1	Total	С	N	О	S	0	0	
3		1	35	16	3	14	2	U	0	
3	D	1	Total	С	N	О	S	0	0	
3	ש	1	35	16	3	14	2	U	U	

 \bullet Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	В	1	Total O S 5 4 1	0	0
4	С	1	Total O S 5 4 1	0	0
4	E	1	Total O S 5 4 1	0	0

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	98	Total O 98 98	0	0
5	В	113	Total O 113 113	0	0
5	С	114	Total O 114 114	0	0
5	D	47	Total O 47 47	0	0
5	E	105	Total O 105 105	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Endo-type membrane-bound lytic murein transglycosylase A









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	78.78Å 94.85Å 162.02Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.76 - 2.30	Depositor
Resolution (A)	22.92 - 2.30	EDS
% Data completeness	99.7 (22.76-2.30)	Depositor
(in resolution range)	99.7 (22.92-2.30)	EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$< I/\sigma(I) > 1$	3.44 (at 2.31Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
D D	0.163 , 0.203	Depositor
R, R_{free}	0.184 , 0.215	DCC
R_{free} test set	5516 reflections (10.12%)	wwPDB-VP
Wilson B-factor (Å ²)	31.6	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37 , 41.7	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7814	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 3.37% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, AMU, BLG, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.43	0/1452	0.55	0/1972
1	В	0.45	0/1462	0.55	0/1984
1	С	0.42	0/1459	0.56	0/1981
1	D	0.38	0/1433	0.55	0/1946
1	Е	0.46	0/1453	0.55	0/1973
All	All	0.43	0/7259	0.55	0/9856

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1415	0	1409	6	0
1	В	1425	0	1426	8	0
1	С	1422	0	1418	11	0
1	D	1397	0	1398	5	0
1	Е	1421	0	1417	7	0
2	F	34	0	30	0	0
2	G	34	0	30	1	0
2	Н	34	0	30	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	35	0	28	1	0
3	В	35	0	29	3	0
3	С	35	0	29	0	0
3	D	35	0	28	0	0
4	В	5	0	0	2	0
4	С	5	0	0	0	0
4	Е	5	0	0	0	0
5	A	98	0	0	0	0
5	В	113	0	0	2	0
5	С	114	0	0	1	0
5	D	47	0	0	0	0
5	Е	105	0	0	0	0
All	All	7814	0	7272	39	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (39) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:302:SO4:S	5:B:493:HOH:O	2.38	0.80
1:C:154:LEU:HD13	1:C:161:ARG:HG3	1.67	0.76
1:C:75:SER:OG	1:C:84:LYS:HE2	2.03	0.58
4:B:302:SO4:O1	5:B:493:HOH:O	2.17	0.58
1:B:50:TRP:HH2	1:B:96:MET:HE3	1.67	0.58
1:D:160:ASP:HB3	1:D:163:LYS:HD2	1.85	0.58
1:B:50:TRP:CH2	1:B:96:MET:HE3	2.39	0.58
1:C:96:MET:SD	1:D:96[A]:MET:SD	3.05	0.55
1:B:154:LEU:HD13	1:B:161:ARG:HG3	1.91	0.53
1:E:107:LEU:HD23	1:E:116:MET:CE	2.40	0.51
1:C:154:LEU:HD13	1:C:161:ARG:CG	2.39	0.51
1:B:73:SER:HB2	3:B:301:BLG:H122	1.92	0.50
1:E:107:LEU:HD23	1:E:116:MET:HE3	1.93	0.50
1:E:84:LYS:HD2	1:E:87:THR:OG1	2.14	0.48
1:A:92:VAL:HG12	1:A:96[B]:MET:CE	2.44	0.47
1:D:112:ARG:HD3	1:D:116:MET:CE	2.45	0.47
2:G:1:AMU:H111	2:G:2:NAG:O5	2.15	0.47
1:A:73:SER:HB3	1:A:82:GLN:OE1	2.16	0.46
1:E:107:LEU:CD2	1:E:116:MET:HE3	2.45	0.45
1:C:101:GLU:HB2	1:C:102:PRO:HD2	1.98	0.45
1:A:84:LYS:HE3	3:A:600:BLG:HG1	1.98	0.45



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
1:A:31:VAL:HB	1:A:32:PRO:HD3	1.99	0.45
1:B:102:PRO:HA	1:B:106:GLU:OE1	2.16	0.44
1:E:31:VAL:HB	1:E:32:PRO:HD3	1.99	0.44
1:C:31:VAL:HB	1:C:32:PRO:HD3	2.00	0.43
1:C:103:THR:HG22	1:C:106:GLU:OE2	2.18	0.43
1:C:19:LYS:N	5:C:493:HOH:O	2.51	0.43
1:A:101:GLU:HB2	1:A:102:PRO:HD2	2.00	0.42
1:B:146:TYR:CE1	3:B:301:BLG:H82	2.55	0.42
1:C:64:GLU:HB3	1:C:82:GLN:CG	2.50	0.42
1:A:163:LYS:O	1:A:167:LYS:HG3	2.19	0.41
1:B:146:TYR:CZ	3:B:301:BLG:H82	2.55	0.41
1:B:101:GLU:HB2	1:B:102:PRO:HD2	2.03	0.41
1:C:24:ASN:OD1	1:C:25:PRO:HA	2.21	0.41
1:E:103:THR:HG22	1:E:106:GLU:HG3	2.03	0.41
1:E:30:LYS:O	1:E:34:GLN:HG3	2.21	0.41
1:D:31:VAL:HB	1:D:32:PRO:HD3	2.02	0.40
1:D:157:PHE:CZ	1:D:179:HIS:CD2	3.09	0.40
1:C:103:THR:HG22	1:C:106:GLU:HG3	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	183/203 (90%)	182 (100%)	1 (0%)	0	100	100
1	В	184/203 (91%)	182 (99%)	2 (1%)	0	100	100
1	С	184/203 (91%)	183 (100%)	1 (0%)	0	100	100
1	D	181/203 (89%)	179 (99%)	2 (1%)	0	100	100
1	Е	183/203 (90%)	181 (99%)	2 (1%)	0	100	100
All	All	915/1015 (90%)	907 (99%)	8 (1%)	0	100	100



There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	A	148/163 (91%)	147 (99%)	1 (1%)	84	92
1	В	149/163 (91%)	149 (100%)	0	100	100
1	\mathbf{C}	149/163 (91%)	148 (99%)	1 (1%)	84	92
1	D	146/163 (90%)	144 (99%)	2 (1%)	67	81
1	E	148/163 (91%)	145 (98%)	3 (2%)	55	72
All	All	740/815 (91%)	733 (99%)	7 (1%)	76	89

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLN
1	С	75	SER
1	D	23	THR
1	D	24	ASN
1	Е	76	ASN
1	Е	99	SER
1	Е	192	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	183	ASN
1	В	20	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Type Chain Re		Res Link		Bond lengths			Bond angles		
MIOI	Type	Chain	Res	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	AMU	F	1	2	20,20,20	0.82	0	24,28,28	1.53	2 (8%)
2	NAG	F	2	2	14,14,15	0.51	0	17,19,21	1.20	3 (17%)
2	AMU	G	1	2	20,20,20	0.80	0	24,28,28	1.32	2 (8%)
2	NAG	G	2	2	14,14,15	0.54	0	17,19,21	1.06	1 (5%)
2	AMU	Н	1	2	20,20,20	0.69	0	24,28,28	1.32	2 (8%)
2	NAG	Н	2	2	14,14,15	0.45	0	17,19,21	1.40	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMU	F	1	2	-	2/14/34/34	0/1/1/1
2	NAG	F	2	2	-	0/6/23/26	0/1/1/1
2	AMU	G	1	2	-	2/14/34/34	0/1/1/1
2	NAG	G	2	2	-	0/6/23/26	0/1/1/1
2	AMU	Н	1	2	_	3/14/34/34	0/1/1/1
2	NAG	Н	2	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (12) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}({}^o)$	$\operatorname{Ideal}({}^{o})$
2	F	1	AMU	O5-C1-C2	4.84	114.38	109.52
2	G	2	NAG	C1-O5-C5	3.67	117.17	112.19
2	G	1	AMU	O5-C1-C2	3.65	113.18	109.52
2	Н	2	NAG	C1-O5-C5	3.35	116.73	112.19
2	Н	1	AMU	O5-C1-C2	2.69	112.22	109.52
2	F	2	NAG	C1-O5-C5	2.39	115.43	112.19
2	Н	2	NAG	O5-C5-C6	2.32	110.84	107.20
2	F	1	AMU	C3-C2-N2	-2.28	107.13	110.91
2	F	2	NAG	O5-C5-C6	2.23	110.70	107.20
2	Н	1	AMU	C1-C2-C3	2.15	113.33	110.25
2	G	1	AMU	C8-C7-N2	2.07	119.61	116.10
2	F	2	NAG	C4-C3-C2	-2.01	108.07	111.02

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	1	AMU	O10-C10-C9-O3
2	F	1	AMU	O11-C10-C9-O3
2	G	1	AMU	O10-C10-C9-O3
2	G	1	AMU	O11-C10-C9-O3
2	Н	1	AMU	O10-C10-C9-O3
2	Н	1	AMU	O11-C10-C9-O3
2	Н	1	AMU	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	1	AMU	1	0
2	G	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.














5.6 Ligand geometry (i)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Bond lengths			Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	BLG	D	600	-	36,36,36	1.80	9 (25%)	41,53,53	1.58	10 (24%)
3	BLG	С	301	-	36,36,36	1.72	10 (27%)	41,53,53	1.57	9 (21%)
3	BLG	A	600	-	36,36,36	1.78	8 (22%)	41,53,53	1.61	9 (21%)
3	BLG	В	301	-	36,36,36	1.82	9 (25%)	41,53,53	2.07	16 (39%)
4	SO4	Е	301	-	4,4,4	0.16	0	6,6,6	0.14	0
4	SO4	В	302	-	4,4,4	0.27	0	6,6,6	0.23	0



Mol	Type	Chain	Pos	Link	В	ond leng	gths	Bond angles		
MIOI	$\operatorname{cl} \mid \operatorname{Type} \mid \operatorname{Chain} \mid \operatorname{Res} \mid$		LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
4	SO4	С	304	-	4,4,4	0.31	0	6,6,6	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BLG	A	600	-	-	1/28/60/60	0/2/2/2
3	BLG	В	301	-	-	9/28/60/60	0/2/2/2
3	BLG	С	301	-	-	3/28/60/60	0/2/2/2
3	BLG	D	600	-	-	5/28/60/60	0/2/2/2

All (36) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(\mathbf{\mathring{A}})$	Ideal(Å)
3	В	301	BLG	C10-N3	5.61	1.45	1.33
3	D	600	BLG	C10-N3	5.44	1.45	1.33
3	D	600	BLG	C1-C2	5.31	1.61	1.53
3	С	301	BLG	C10-N3	5.03	1.44	1.33
3	В	301	BLG	C1-C2	5.02	1.61	1.53
3	A	600	BLG	C10-N3	4.67	1.43	1.33
3	A	600	BLG	C1-C2	4.61	1.60	1.53
3	С	301	BLG	C1-C2	4.28	1.60	1.53
3	A	600	BLG	CA-N	-3.44	1.43	1.48
3	D	600	BLG	C7-N2	3.27	1.45	1.34
3	В	301	BLG	C7-N2	3.23	1.45	1.34
3	A	600	BLG	C7-N2	3.15	1.45	1.34
3	С	301	BLG	C7-N2	3.02	1.44	1.34
3	В	301	BLG	O4-S4	-2.92	1.48	1.57
3	С	301	BLG	O4-S4	-2.80	1.49	1.57
3	В	301	BLG	O5-C1	2.77	1.48	1.41
3	D	600	BLG	O4-S4	-2.74	1.49	1.57
3	С	301	BLG	CA-N	-2.69	1.44	1.48
3	A	600	BLG	O4-S4	-2.65	1.49	1.57
3	С	301	BLG	O4-C4	-2.61	1.41	1.46
3	A	600	BLG	CG-CB	2.60	1.58	1.52
3	В	301	BLG	CA-N	-2.59	1.44	1.48
3	С	301	BLG	CG-CB	2.45	1.58	1.52
3	D	600	BLG	CG-CB	2.43	1.58	1.52
3	В	301	BLG	O4-C4	-2.43	1.41	1.46
3	D	600	BLG	O5-C1	2.39	1.47	1.41



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	$[Ideal(\AA)]$
3	A	600	BLG	O5-C1	2.36	1.47	1.41
3	С	301	BLG	O5-C1	2.32	1.47	1.41
3	D	600	BLG	O1-CB	2.31	1.49	1.44
3	С	301	BLG	O5-C5	-2.31	1.38	1.44
3	В	301	BLG	CG-CB	2.30	1.57	1.52
3	С	301	BLG	O3-C3	-2.25	1.37	1.43
3	A	600	BLG	O5-C5	-2.24	1.38	1.44
3	В	301	BLG	O5-C5	-2.21	1.39	1.44
3	D	600	BLG	CA-N	-2.16	1.45	1.48
3	D	600	BLG	O4-C4	-2.16	1.41	1.46

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
3	A	600	BLG	OS1-S-C12	4.96	112.89	106.92
3	В	301	BLG	O3-C3-C2	4.41	118.57	109.66
3	В	301	BLG	OS1-S-C12	4.37	112.18	106.92
3	В	301	BLG	OS2-S-OS1	-4.06	99.89	113.95
3	A	600	BLG	CG-CD-N	3.79	112.70	106.74
3	В	301	BLG	C4-C3-C2	3.68	119.20	110.62
3	В	301	BLG	O3-C3-C4	3.61	119.51	109.94
3	В	301	BLG	C1-O1-CB	-3.60	109.77	115.33
3	С	301	BLG	O1-C1-C2	-3.60	102.03	108.24
3	D	600	BLG	OS2-S-OS1	-3.57	101.60	113.95
3	С	301	BLG	CG-CD-N	3.52	112.28	106.74
3	С	301	BLG	C1-O1-CB	-3.51	109.91	115.33
3	A	600	BLG	OS2-S-OS1	-3.35	102.36	113.95
3	С	301	BLG	OS1-S-C12	3.26	110.84	106.92
3	D	600	BLG	OS1-S-C12	3.24	110.81	106.92
3	D	600	BLG	OS2-S-C12	3.21	110.78	106.92
3	A	600	BLG	C1-C2-N2	-3.17	105.55	111.00
3	D	600	BLG	CG-CD-N	3.12	111.65	106.74
3	D	600	BLG	C1-C2-N2	-3.05	105.75	111.00
3	В	301	BLG	OS2-S-C12	2.97	110.49	106.92
3	В	301	BLG	C3-C2-N2	2.93	116.15	110.62
3	С	301	BLG	OS2-S-OS1	-2.90	103.89	113.95
3	В	301	BLG	O1-C1-C2	-2.87	103.28	108.24
3	В	301	BLG	CG-CD-N	2.81	111.16	106.74
3	С	301	BLG	C1-O5-C5	-2.80	108.19	113.69
3	D	600	BLG	O42-S4-O41	-2.79	101.03	112.22
3	D	600	BLG	C1-O1-CB	-2.77	111.05	115.33
3	В	301	BLG	OS3-S-C12	2.74	110.20	105.77



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	D	600	BLG	OS3-S-C12	2.69	110.11	105.77
3	С	301	BLG	OS2-S-C12	2.61	110.06	106.92
3	В	301	BLG	C1-O5-C5	-2.52	108.75	113.69
3	В	301	BLG	C1-C2-N2	-2.49	106.72	111.00
3	A	600	BLG	C1-O5-C5	-2.46	108.86	113.69
3	A	600	BLG	OS2-S-C12	2.42	109.83	106.92
3	D	600	BLG	CG-CD-C10	-2.34	107.70	111.19
3	A	600	BLG	C12-C11-N3	-2.33	104.47	111.30
3	В	301	BLG	C2-N2-C7	-2.33	117.51	123.18
3	С	301	BLG	C1-C2-N2	-2.28	107.09	111.00
3	D	600	BLG	C1-O5-C5	-2.25	109.28	113.69
3	С	301	BLG	CG-CB-CA	2.13	106.84	103.80
3	В	301	BLG	O42-S4-O41	-2.13	103.67	112.22
3	В	301	BLG	C9-CA-N	2.12	115.26	111.46
3	A	600	BLG	O1-C1-C2	-2.10	104.62	108.24
3	A	600	BLG	C1-O1-CB	-2.03	112.19	115.33

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	600	BLG	C4-O4-S4-O43
3	В	301	BLG	C12-C11-N3-C10
3	В	301	BLG	C11-C12-S-OS1
3	В	301	BLG	C11-C12-S-OS2
3	В	301	BLG	C11-C12-S-OS3
3	С	301	BLG	C4-O4-S4-O41
3	С	301	BLG	C4-O4-S4-O43
3	D	600	BLG	C4-O4-S4-O41
3	D	600	BLG	C4-O4-S4-O43
3	В	301	BLG	C4-O4-S4-O41
3	В	301	BLG	C4-O4-S4-O42
3	С	301	BLG	C4-O4-S4-O42
3	D	600	BLG	C4-O4-S4-O42
3	В	301	BLG	C4-O4-S4-O43
3	В	301	BLG	O10-C10-CD-N
3	В	301	BLG	N3-C10-CD-N
3	D	600	BLG	N3-C11-C12-S
3	D	600	BLG	C11-C12-S-OS2


There are no ring outliers.

3 monomers are involved in 6 short contacts:





Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	BLG	1	0
3	В	301	BLG	3	0
4	В	302	SO4	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q<0.9
1	A	184/203 (90%)	0.15	10 (5%) 25 32	22, 32, 50, 84	0
1	В	185/203~(91%)	-0.12	5 (2%) 54 62	20, 29, 46, 81	0
1	С	185/203 (91%)	0.05	4 (2%) 62 69	21, 35, 54, 63	0
1	D	182/203 (89%)	0.40	16 (8%) 10 13	29, 44, 70, 86	0
1	E	185/203 (91%)	0.28	15 (8%) 12 16	22, 37, 73, 85	0
All	All	921/1015 (90%)	0.15	50 (5%) 25 32	20, 35, 64, 86	0

All (50) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	90	ARG	5.4
1	Е	19	LYS	5.2
1	Е	96	MET	4.3
1	Е	98	TRP	4.3
1	A	20	HIS	4.1
1	В	19	LYS	4.0
1	С	60	ILE	3.7
1	D	182	ARG	3.5
1	A	182	ARG	3.5
1	D	24	ASN	3.3
1	Е	90	ARG	3.1
1	Е	97	GLY	3.1
1	Е	112	ARG	3.1
1	Е	95	ARG	3.1
1	Е	21	ASP	3.0
1	D	141	ALA	3.0
1	С	74	LYS	2.9
1	В	20	HIS	2.9
1	D	26	PRO	2.8
1	В	21	ASP	2.7



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	143	VAL	2.7
1	С	143	VAL	2.7
1	D	94	ARG	2.7
1	D	163	LYS	2.7
1	Е	99	SER	2.7
1	A	141	ALA	2.7
1	D	178	GLU	2.6
1	D	170	ASP	2.6
1	Е	86	SER	2.6
1	В	143	VAL	2.6
1	Е	85	ALA	2.5
1	A	193	ILE	2.4
1	A	117	GLY	2.4
1	D	144	VAL	2.4
1	D	25	PRO	2.4
1	Е	93	TYR	2.3
1	D	160	ASP	2.3
1	A	60	ILE	2.3
1	D	168	ILE	2.3
1	D	185	PRO	2.2
1	A	74	LYS	2.2
1	Е	100	GLY	2.1
1	A	176	PHE	2.1
1	Е	20	HIS	2.1
1	В	60	ILE	2.1
1	D	167	LYS	2.1
1	С	193	ILE	2.1
1	Е	75	SER	2.1
1	D	143	VAL	2.0
1	A	140	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	AMU	F	1	20/20	0.90	0.15	36,40,58,61	0
2	AMU	Н	1	20/20	0.90	0.17	43,57,68,68	0
2	AMU	G	1	20/20	0.91	0.16	34,41,58,60	0
2	NAG	Н	2	14/15	0.92	0.13	35,38,45,49	0
2	NAG	G	2	14/15	0.93	0.11	34,37,39,39	0
2	NAG	F	2	14/15	0.94	0.12	33,36,43,46	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.











6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$ extbf{B-factors}(extbf{A}^2)$	Q<0.9
3	BLG	В	301	35/35	0.70	0.33	35,54,144,146	35
3	BLG	D	600	35/35	0.92	0.15	28,41,95,95	0
3	BLG	С	301	35/35	0.93	0.14	24,32,61,64	0
3	BLG	A	600	35/35	0.94	0.15	28,33,63,64	0
4	SO4	В	302	5/5	0.94	0.12	62,63,64,67	0
4	SO4	Е	301	5/5	0.94	0.21	63,68,70,71	0
4	SO4	С	304	5/5	0.97	0.18	42,43,46,47	0

The following is a graphical depiction of the model fit to experimental electron density of all




instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around BLG D 600:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around BLG C 301:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

