



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 10, 2023 – 02:04 AM EDT

PDB ID : 4HK6  
Title : Crystal structure of Cordyceps militaris IDCase in complex with 5-nitro-uracil  
Authors : Xu, S.; Zhu, J.; Ding, J.  
Deposited on : 2012-10-15  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	387	 6% 89% 6% • 5%
1	B	387	 6% 87% 8% • 5%
1	C	387	 4% 87% 9% • •
1	D	387	 5% 87% 8% 5%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 12008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Uracil-5-carboxylate decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	369	2819	1815	482	508	14	0	0	0
1	B	368	2810	1809	480	507	14	0	0	0
1	C	374	2860	1840	489	517	14	0	0	0
1	D	369	2822	1817	482	509	14	0	1	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP G3J531
A	-1	GLY	-	expression tag	UNP G3J531
A	0	SER	-	expression tag	UNP G3J531
A	377	LEU	-	expression tag	UNP G3J531
A	378	GLU	-	expression tag	UNP G3J531
A	379	HIS	-	expression tag	UNP G3J531
A	380	HIS	-	expression tag	UNP G3J531
A	381	HIS	-	expression tag	UNP G3J531
A	382	HIS	-	expression tag	UNP G3J531
A	383	HIS	-	expression tag	UNP G3J531
A	384	HIS	-	expression tag	UNP G3J531
B	-2	MET	-	expression tag	UNP G3J531
B	-1	GLY	-	expression tag	UNP G3J531
B	0	SER	-	expression tag	UNP G3J531
B	377	LEU	-	expression tag	UNP G3J531
B	378	GLU	-	expression tag	UNP G3J531
B	379	HIS	-	expression tag	UNP G3J531
B	380	HIS	-	expression tag	UNP G3J531
B	381	HIS	-	expression tag	UNP G3J531
B	382	HIS	-	expression tag	UNP G3J531
B	383	HIS	-	expression tag	UNP G3J531

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	384	HIS	-	expression tag	UNP G3J531
C	-2	MET	-	expression tag	UNP G3J531
C	-1	GLY	-	expression tag	UNP G3J531
C	0	SER	-	expression tag	UNP G3J531
C	377	LEU	-	expression tag	UNP G3J531
C	378	GLU	-	expression tag	UNP G3J531
C	379	HIS	-	expression tag	UNP G3J531
C	380	HIS	-	expression tag	UNP G3J531
C	381	HIS	-	expression tag	UNP G3J531
C	382	HIS	-	expression tag	UNP G3J531
C	383	HIS	-	expression tag	UNP G3J531
C	384	HIS	-	expression tag	UNP G3J531
D	-2	MET	-	expression tag	UNP G3J531
D	-1	GLY	-	expression tag	UNP G3J531
D	0	SER	-	expression tag	UNP G3J531
D	377	LEU	-	expression tag	UNP G3J531
D	378	GLU	-	expression tag	UNP G3J531
D	379	HIS	-	expression tag	UNP G3J531
D	380	HIS	-	expression tag	UNP G3J531
D	381	HIS	-	expression tag	UNP G3J531
D	382	HIS	-	expression tag	UNP G3J531
D	383	HIS	-	expression tag	UNP G3J531
D	384	HIS	-	expression tag	UNP G3J531

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is 5-nitouracil (three-letter code: 5NU) (formula: C<sub>4</sub>H<sub>3</sub>N<sub>3</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	11	4	3	4	0	0
3	B	1	11	4	3	4	0	0
3	C	1	11	4	3	4	0	0
3	D	1	11	4	3	4	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	139	139	139	0	0
4	B	156	156	156	0	0
4	C	183	183	183	0	0
4	D	171	171	171	0	0





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	77.26Å 145.28Å 77.51Å 90.00° 95.84° 90.00°	Depositor
Resolution (Å)	77.10 – 2.30 45.05 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (77.10-2.30) 99.4 (45.05-2.30)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.79 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.153 , 0.202 0.153 , 0.201	Depositor DCC
$R_{free}$ test set	3808 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	20.6	Xtrriage
Anisotropy	0.352	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 46.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.019 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	12008	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.46% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5NU, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.45	2/2891 (0.1%)	0.53	0/3937
1	B	0.45	2/2882 (0.1%)	0.53	0/3926
1	C	0.48	3/2933 (0.1%)	0.54	0/3994
1	D	0.45	2/2897 (0.1%)	0.54	0/3945
All	All	0.46	9/11603 (0.1%)	0.54	0/15802

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	100	TRP	NE1-CE2	-5.59	1.30	1.37
1	B	287	TRP	CD2-CE2	5.33	1.47	1.41
1	D	338	TRP	CD2-CE2	5.20	1.47	1.41
1	D	287	TRP	CD2-CE2	5.15	1.47	1.41
1	C	338	TRP	CD2-CE2	5.15	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2819	0	2838	10	0
1	B	2810	0	2825	14	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	2860	0	2873	16	0
1	D	2822	0	2843	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	11	0	3	1	0
3	B	11	0	3	1	0
3	C	11	0	3	1	0
3	D	11	0	3	1	0
4	A	139	0	0	0	0
4	B	156	0	0	0	0
4	C	183	0	0	1	0
4	D	171	0	0	3	0
All	All	12008	0	11391	57	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 57 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:GLN:HA	1:A:79:GLN:HE21	1.55	0.71
1:A:7:VAL:HG23	1:A:90:ARG:HG2	1.77	0.65
1:D:334:VAL:O	1:D:338:TRP:NE1	2.38	0.57
1:A:25:GLU:HG3	1:A:33:VAL:O	2.06	0.55
1:C:226:THR:O	1:C:230:VAL:HG23	2.08	0.53

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	367/387 (95%)	359 (98%)	6 (2%)	2 (0%)	29	35
1	B	366/387 (95%)	357 (98%)	7 (2%)	2 (0%)	29	35
1	C	372/387 (96%)	359 (96%)	11 (3%)	2 (0%)	29	35
1	D	368/387 (95%)	359 (98%)	7 (2%)	2 (0%)	29	35
All	All	1473/1548 (95%)	1434 (97%)	31 (2%)	8 (0%)	29	35

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	197	HIS
1	B	197	HIS
1	D	197	HIS
1	A	197	HIS
1	B	364	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/313 (96%)	290 (97%)	9 (3%)	41	57
1	B	298/313 (95%)	290 (97%)	8 (3%)	44	61
1	C	303/313 (97%)	292 (96%)	11 (4%)	35	49
1	D	300/313 (96%)	293 (98%)	7 (2%)	50	67
All	All	1200/1252 (96%)	1165 (97%)	35 (3%)	42	58

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	51	LEU
1	D	114	ASP
1	D	323	ASP
1	B	114	ASP
1	B	90	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	202	ASN
1	D	73	HIS
1	D	335	GLN
1	D	79	GLN
1	B	177	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	5NU	C	402	2	9,11,11	3.13	2 (22%)	11,15,15	4.49	7 (63%)
3	5NU	D	402	2	9,11,11	3.14	4 (44%)	11,15,15	4.70	7 (63%)
3	5NU	B	402	2	9,11,11	3.17	4 (44%)	11,15,15	4.58	6 (54%)
3	5NU	A	402	2	9,11,11	3.53	4 (44%)	11,15,15	4.59	6 (54%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	5NU	C	402	2	-	0/2/4/4	0/1/1/1
3	5NU	D	402	2	-	0/2/4/4	0/1/1/1
3	5NU	B	402	2	-	0/2/4/4	0/1/1/1
3	5NU	A	402	2	-	0/2/4/4	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	402	5NU	C5-C4	-8.69	1.39	1.47
3	A	402	5NU	C5-C4	-8.61	1.39	1.47
3	D	402	5NU	C5-C4	-8.56	1.39	1.47
3	B	402	5NU	C5-C4	-8.03	1.40	1.47
3	A	402	5NU	O53-N51	4.63	1.30	1.22

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	402	5NU	C5-C4-N3	12.52	118.44	112.19
3	A	402	5NU	C5-C4-N3	12.44	118.40	112.19
3	B	402	5NU	C5-C4-N3	12.36	118.36	112.19
3	C	402	5NU	C5-C4-N3	12.17	118.26	112.19
3	D	402	5NU	N1-C2-N3	5.14	120.96	115.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	402	5NU	1	0
3	D	402	5NU	1	0
3	B	402	5NU	1	0
3	A	402	5NU	1	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	369/387 (95%)	-0.01	23 (6%) 20 26	9, 22, 58, 85	0
1	B	368/387 (95%)	-0.14	22 (5%) 21 28	10, 21, 53, 81	0
1	C	374/387 (96%)	-0.22	14 (3%) 41 48	10, 20, 50, 72	0
1	D	369/387 (95%)	-0.17	18 (4%) 29 36	10, 20, 52, 71	0
All	All	1480/1548 (95%)	-0.14	77 (5%) 27 34	9, 21, 53, 85	0

The worst 5 of 77 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	375	ALA	8.7
1	D	334	VAL	7.0
1	D	336	GLY	6.6
1	C	378	GLU	6.5
1	A	40	ASP	6.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	5NU	A	402	11/11	0.94	0.14	18,21,24,29	0
3	5NU	B	402	11/11	0.94	0.14	18,20,24,28	0
3	5NU	C	402	11/11	0.94	0.13	14,18,24,26	0
3	5NU	D	402	11/11	0.94	0.13	18,19,22,28	0
2	ZN	C	401	1/1	0.99	0.08	21,21,21,21	0
2	ZN	A	401	1/1	1.00	0.07	25,25,25,25	0
2	ZN	D	401	1/1	1.00	0.06	21,21,21,21	0
2	ZN	B	401	1/1	1.00	0.09	22,22,22,22	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.