

# Full wwPDB X-ray Structure Validation Report (i)

#### May 13, 2020 - 05:44 pm BST

PDB ID	:	1HNH
$\operatorname{Title}$	:	CRYSTAL STRUCTURE OF BETA-KETOACYL-ACP SYNTHASE III +
		DEGRADED FORM OF ACETYL-COA
Authors	:	Qiu, X.; Janson, C.A.; Smith, W.W.; Head, M.; Lonsdale, J.; Konstantinidis,
		A.K.
Deposited on		
$\operatorname{Resolution}$	:	1.90  Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
$\mathrm{EDS}$	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain		
					_
1	А	317	78%	21%	•



# 2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 2618 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called BETA-KETOACYL-ACYL CARRIER PROTEIN SYNTHASE III.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	А	317	Total 2334	C 1460	N 401	O 460	S 5	${ m Se} 8$	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	1	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
А	25	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
A	54	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
А	65	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
А	97	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
А	112	SCY	CYS	MODIFIED RESIDUE	UNP P0A6R0
А	207	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
А	260	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0
А	262	MSE	MET	MODIFIED RESIDUE	UNP P0A6R0

• Molecule 2 is COENZYME A (three-letter code: COA) (formula: C<sub>21</sub>H<sub>36</sub>N<sub>7</sub>O<sub>16</sub>P<sub>3</sub>S).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	А	1	Total 48	С 21	N 7	O 16	Р 3	${ m S}$ 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	236	Total         O           236         236	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: BETA-KETOACYL-ACYL CARRIER PROTEIN SYNTHASE III





## 4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants	72.45Å $72.45$ Å $102.80$ Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	20.00 - 1.90	Depositor
% Data completeness	(Not available) (20.00-1.90)	Depositor
(in resolution range)	(1101 available) (20.00 1.50)	Depositor
$R_{merge}$	0.08	Depositor
R <sub>sym</sub>	(Not available)	Depositor
Refinement program	CNS	Depositor
$R, R_{free}$	0.221 , $0.274$	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2618	wwPDB-VP
Average B, all atoms $(Å^2)$	23.0	wwPDB-VP



# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: COA, SCY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths			nd angles
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.65	0/2356	0.78	2/3197~(0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	273	GLY	N-CA-C	-5.44	99.51	113.10
1	А	157	PHE	N-CA-C	5.04	124.62	111.00

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2334	0	2294	56	0
2	А	48	0	32	3	0
3	А	236	0	0	6	1
All	All	2618	0	2326	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic	Clash
	Atom-2	distance (Å)	overlap (Å)
1:A:45:ALA:HB2	1:A:54:MSE:HE1	1.33	1.06
1:A:258:LEU:HB3	1:A:260:MSE:HE2	1.56	0.87
1:A:198:ASN:HD22	1:A:198:ASN:N	1.80	0.78
1:A:45:ALA:CB	1:A:54:MSE:HE1	2.12	0.78
1:A:258:LEU:CB	1:A:260:MSE:HE2	2.14	0.77
1:A:7:GLY:HA2	1:A:65:MSE:HE2	1.72	0.71
1:A:245:GLN:HB3	1:A:269:LEU:HB3	1.73	0.70
1:A:11:TYR:CD2	1:A:54:MSE:HG2	2.33	0.64
1:A:198:ASN:N	1:A:198:ASN:ND2	2.46	0.63
1:A:274:ASN:ND2	2:A:350:COA:S1P	2.72	0.62
1:A:110:ALA:HB3	1:A:114:GLY:HA2	1.83	0.60
1:A:87:PHE:HA	1:A:88:PRO:C	2.21	0.59
1:A:65:MSE:HG3	3:A:583:HOH:O	2.04	0.57
1:A:45:ALA:HB2	1:A:54:MSE:CE	2.22	0.57
1:A:11:TYR:HD2	1:A:54:MSE:HG2	1.70	0.56
1:A:2:TYR:CD2	1:A:172:PRO:HD3	2.40	0.56
1:A:293:ILE:HD13	1:A:293:ILE:N	2.23	0.54
1:A:258:LEU:CB	1:A:260:MSE:CE	2.86	0.54
1:A:132:LYS:HE2	1:A:133:TYR:CZ	2.43	0.53
1:A:93:GLN:O	1:A:97:MSE:HG3	2.10	0.52
1:A:212:VAL:HB	2:A:350:COA:H62	1.92	0.52
1:A:45:ALA:CB	1:A:54:MSE:CE	2.84	0.52
1:A:307:GLY:N	1:A:308:PHE:HA	2.25	0.51
1:A:197:VAL:C	1:A:198:ASN:HD22	2.13	0.51
1:A:195:ASP:HB3	1:A:198:ASN:O	2.11	0.51
1:A:176:SER:OG	1:A:178:HIS:HE1	1.94	0.50
1:A:197:VAL:C	1:A:198:ASN:ND2	2.66	0.49
1:A:284:LEU:O	1:A:288:VAL:HG23	2.11	0.49
1:A:129:GLY:HA2	3:A:487:HOH:O	2.12	0.49
1:A:224:VAL:HG11	1:A:258:LEU:HG	1.95	0.48
1:A:212:VAL:CG1	2:A:350:COA:H62	2.43	0.47
1:A:262:MSE:HG2	3:A:578:HOH:O	2.14	0.47
1:A:2:TYR:CE2	1:A:172:PRO:HD3	2.50	0.47
1:A:235:ARG:HG3	3:A:461:HOH:O	2.15	0.46
1:A:54:MSE:HB3	1:A:161:ALA:HB2	1.97	0.46
1:A:32:TRP:CE2	1:A:36:ARG:HG3	2.50	0.46
1:A:304:PHE:HZ	1:A:308:PHE:CD2	2.34	0.46
1:A:261:SER:C	1:A:263:ASP:N	2.68	0.44
1:A:301:LEU:O	1:A:312:SER:HA	2.17	0.44
1:A:224:VAL:O	1:A:228:LEU:HG	2.17	0.44
1:A:8:THR:O	1:A:62:ALA:HA	2.18	0.44

Continued on next page...



Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:12:LEU:HD11	1:A:272:HIS:NE2	2.32	0.44
1:A:213:PHE:O	1:A:217:VAL:HG23	2.17	0.44
1:A:32:TRP:CD2	1:A:36:ARG:HG3	2.53	0.43
1:A:240:TRP:CD1	1:A:293:ILE:HD12	2.54	0.43
1:A:219:GLU:OE1	1:A:219:GLU:HA	2.18	0.43
1:A:31:GLU:HB2	3:A:549:HOH:O	2.18	0.42
1:A:258:LEU:HB2	1:A:260:MSE:CE	2.50	0.42
1:A:146:CYS:O	1:A:148:PRO:HD3	2.20	0.41
1:A:26:VAL:HB	1:A:151:ARG:HB3	2.02	0.41
1:A:93:GLN:NE2	3:A:412:HOH:O	2.53	0.41
1:A:175:ILE:CG2	1:A:231:ASN:ND2	2.84	0.41
1:A:298:LEU:HD12	1:A:316:ARG:HD3	2.02	0.41
1:A:258:LEU:HD23	1:A:260:MSE:HE1	2.03	0.40
1:A:10:SER:OG	1:A:282:CYS:HA	2.21	0.40
1:A:13:PRO:HG3	1:A:54:MSE:HE2	2.03	0.40

Continued from previous page...

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:628:HOH:O	3:A:628:HOH:O[8_665]	2.07	0.13

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles	
1	А	314/317~(99%)	301~(96%)	12~(4%)	1 (0%)	41 31	

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	209	GLY



#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	А	241/238~(101%)	235~(98%)	6(2%)	47 41	

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	$\mathbf{Res}$	Type
1	А	31	GLU
1	А	42	ARG
1	А	71	ASP
1	А	142	LEU
1	А	198	ASN
1	А	258	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	85	HIS
1	А	93	GLN
1	А	124	GLN
1	А	178	HIS
1	А	193	ASN
1	А	198	ASN
1	А	210	ASN
1	А	231	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul



statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
1	SCY	А	112	1	$7,\!8,\!9$	1.10	0	$3,\!9,\!11$	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SCY	А	112	1	-	0/5/7/9	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths		Bond angles			
	туре	Unam	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
2	COA	А	350	-	41,50,50	2.04	9 (21%)	52,75,75	2.11	15 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	COA	А	350	-	-	9/44/64/64	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	А	350	COA	C2A-N3A	5.16	1.40	1.32
2	А	350	COA	O4B-C1B	4.59	1.47	1.41
2	А	350	COA	C8A-N7A	4.18	1.42	1.34
2	А	350	COA	C3P-N4P	3.87	1.55	1.46
2	А	350	COA	P3B-O3B	3.74	1.66	1.59
2	А	350	COA	O9P-C9P	3.61	1.30	1.23
2	А	350	COA	P3B-07A	3.22	1.60	1.50
2	А	350	COA	C5A-N7A	-2.86	1.29	1.39
2	А	350	COA	C2B-C3B	-2.18	1.48	1.52

All (	(15)	bond	angle	outliers	are	listed	below:	
1	( <del>-</del> ~ )			0.0101010		110000		

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	350	COA	O4B-C1B-C2B	-6.41	97.56	106.93
2	А	350	COA	C1B-N9A-C4A	-5.35	117.25	126.64
2	А	350	COA	C3P-N4P-C5P	5.33	132.74	122.84
2	А	350	COA	C7P-N8P-C9P	4.47	130.56	122.59
2	А	350	COA	N3A-C2A-N1A	-4.27	122.00	128.68
2	А	350	COA	CDP-CBP-CAP	3.27	114.50	108.82
2	А	350	COA	C7P-C6P-C5P	-3.06	107.27	112.36
2	А	350	COA	N6A-C6A-N1A	2.85	124.48	118.57
2	А	350	COA	O2B-C2B-C3B	2.79	119.09	111.17
2	А	350	COA	C2A-N1A-C6A	2.56	123.13	118.75
2	А	350	COA	C6P-C5P-N4P	2.38	120.42	116.42
2	А	350	COA	O5P-C5P-N4P	-2.18	118.90	123.01
2	А	350	COA	C6P-C7P-N8P	2.15	116.24	111.90
2	А	350	COA	P2A-O3A-P1A	-2.06	125.75	132.83
2	А	350	COA	O9P-C9P-N8P	2.05	127.39	122.99



There are no chirality outliers.

Mol	Chain	$\mathbf{Res}$	Type	Atoms
2	А	350	COA	C5P-C6P-C7P-N8P
2	А	350	COA	C2P-C3P-N4P-C5P
2	А	350	COA	O5P-C5P-N4P-C3P
2	А	350	COA	C6P-C5P-N4P-C3P
2	А	350	COA	CAP-C9P-N8P-C7P
2	А	350	COA	O9P-C9P-N8P-C7P
2	А	350	COA	P1A-O3A-P2A-O5A
2	А	350	COA	S1P-C2P-C3P-N4P
2	А	350	COA	P1A-O3A-P2A-O4A

All (9) torsion outliers are listed below:


There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	350	COA	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

