



Full wwPDB X-ray Structure Validation Report ⓘ

May 21, 2020 – 10:03 pm BST

PDB ID : 2HRO
Title : Structure of the full-length Enzyme I of the PTS system from *Staphylococcus carnosus*
Authors : Marquez, J.A.; Reinelt, S.; Koch, B.; Engelman, R.; Hengstenberg, W.; Schefzek, K.
Deposited on : 2006-07-20
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	573	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4247 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Phosphoenolpyruvate-protein phosphotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	551	4070	2551	697	803	19	0	0	0

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	124	MET	ILE	SEE REMARK 999	UNP P23533
A	?	-	LYS	SEE REMARK 999	UNP P23533
A	406	LEU	PHE	SEE REMARK 999	UNP P23533
A	408	GLU	LYS	SEE REMARK 999	UNP P23533
A	409	GLU	LYS	SEE REMARK 999	UNP P23533
A	410	GLU	ASN	SEE REMARK 999	UNP P23533
A	411	ARG	VAL	SEE REMARK 999	UNP P23533
A	412	ALA	LEU	SEE REMARK 999	UNP P23533
A	413	ASN	THR	SEE REMARK 999	UNP P23533
A	416	ASN	MET	SEE REMARK 999	UNP P23533
A	417	GLU	LYS	SEE REMARK 999	UNP P23533
A	418	GLY	ALA	SEE REMARK 999	UNP P23533
A	419	TYR	MET	SEE REMARK 999	UNP P23533
A	420	GLU	LYS	SEE REMARK 999	UNP P23533
A	479	ASN	ILE	SEE REMARK 999	UNP P23533
A	480	PRO	SER	SEE REMARK 999	UNP P23533
A	481	ALA	ASN	SEE REMARK 999	UNP P23533
A	482	ILE	PHE	SEE REMARK 999	UNP P23533
A	483	LEU	SER	SEE REMARK 999	UNP P23533
A	484	ARG	PHE	SEE REMARK 999	UNP P23533
A	536	ARG	VAL	SEE REMARK 999	UNP P23533

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	172	172	172	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Phosphoenolpyruvate-protein phosphotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	173.36Å 46.85Å 85.30Å 90.00° 101.46° 90.00°	Depositor
Resolution (Å)	34.86 – 2.50 34.86 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (34.86-2.50) 98.9 (34.86-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 2.39Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.215 , 0.267 0.216 , 0.266	Depositor DCC
R_{free} test set	1321 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	43.0	Xtrriage
Anisotropy	0.972	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 50.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4247	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/4119	0.59	3/5572 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	201	PRO	N-CA-CB	5.74	110.18	103.30
1	A	166	PRO	N-CA-CB	5.62	110.04	103.30
1	A	232	PRO	N-CA-CB	5.59	110.02	103.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4070	0	3828	85	0
2	A	5	0	0	0	0
3	A	172	0	0	0	0
All	All	4247	0	3828	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396:THR:HG22	1:A:398:GLN:H	1.42	0.85
1:A:345:TYR:O	1:A:346:LEU:HB2	1.80	0.79
1:A:558:THR:OG1	1:A:561:GLU:HG3	1.81	0.78
1:A:428:GLY:HA2	1:A:447:VAL:HG13	1.68	0.76
1:A:54:GLU:O	1:A:58:ILE:HG12	1.92	0.69
1:A:547:LYS:O	1:A:551:GLU:HG3	1.94	0.68
1:A:411:ARG:NH2	1:A:423:ASP:HA	2.08	0.68
1:A:65:GLN:HE22	1:A:173:ASN:HA	1.60	0.66
1:A:431:VAL:HG13	1:A:436:THR:CG2	2.26	0.66
1:A:411:ARG:HH21	1:A:423:ASP:HA	1.60	0.65
1:A:246:ASN:HA	1:A:249:LYS:HE3	1.79	0.64
1:A:367:GLN:O	1:A:370:ILE:HG22	1.99	0.62
1:A:100:ASN:C	1:A:100:ASN:HD22	2.04	0.60
1:A:149:ASN:HB2	1:A:150:PRO:HD2	1.86	0.58
1:A:8:ILE:HA	1:A:203:VAL:HA	1.85	0.58
1:A:447:VAL:HG12	1:A:449:PHE:H	1.69	0.58
1:A:345:TYR:HD1	1:A:346:LEU:HD13	1.69	0.57
1:A:428:GLY:HA2	1:A:447:VAL:CG1	2.36	0.56
1:A:308:MET:SD	1:A:370:ILE:HD12	2.46	0.56
1:A:396:THR:HG22	1:A:398:GLN:N	2.16	0.55
1:A:14:VAL:H	1:A:222:ASP:HA	1.72	0.55
1:A:509:ASP:O	1:A:513:ILE:HG13	2.07	0.54
1:A:297:ARG:HG2	1:A:333:ARG:NH2	2.23	0.53
1:A:487:LYS:HB2	1:A:519:LEU:HD22	1.90	0.53
1:A:325:MET:O	1:A:328:LYS:HB2	2.10	0.52
1:A:380:ARG:HD3	1:A:419:TYR:CD2	2.45	0.51
1:A:300:PHE:CE2	1:A:301:LEU:HD22	2.46	0.50
1:A:78:LEU:O	1:A:82:GLU:HG2	2.11	0.50
1:A:345:TYR:CD1	1:A:346:LEU:HD13	2.46	0.50
1:A:150:PRO:HB3	1:A:171:GLN:O	2.10	0.50
1:A:30:ASP:O	1:A:32:GLU:HG3	2.13	0.49
1:A:21:LEU:HA	1:A:160:ILE:O	2.12	0.49
1:A:301:LEU:HG	1:A:314:GLN:HG2	1.94	0.49
1:A:337:ILE:HD13	1:A:343:LEU:HD11	1.93	0.49
1:A:332:VAL:HG11	1:A:378:LEU:HD21	1.95	0.49
1:A:67:GLY:O	1:A:68:ALA:HB3	2.13	0.48
1:A:113:ILE:O	1:A:117:GLU:HG2	2.13	0.48
1:A:237:ILE:O	1:A:241:GLN:HB2	2.14	0.47
1:A:109:SER:O	1:A:113:ILE:HG13	2.14	0.47
1:A:14:VAL:H	1:A:222:ASP:CA	2.26	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:100:ASN:C	1:A:100:ASN:ND2	2.68	0.47
1:A:76:ALA:O	1:A:80:VAL:HG23	2.14	0.46
1:A:245:GLU:O	1:A:249:LYS:HB2	2.15	0.46
1:A:88:GLN:HB2	1:A:89:PRO:HD3	1.97	0.46
1:A:319:LYS:HE2	1:A:384:PHE:CE2	2.51	0.46
1:A:542:ASN:HB3	1:A:545:GLU:HB2	1.97	0.46
1:A:20:TYR:H	1:A:21:LEU:HD12	1.82	0.45
1:A:431:VAL:HG13	1:A:436:THR:HG22	1.97	0.45
1:A:328:LYS:HA	1:A:328:LYS:HE2	1.99	0.45
1:A:459:GLN:OE1	1:A:466:ARG:HG2	2.16	0.45
1:A:248:PHE:O	1:A:252:GLN:HB2	2.17	0.45
1:A:65:GLN:NE2	1:A:173:ASN:HA	2.31	0.45
1:A:15:ALA:HB1	1:A:179:GLY:HA2	1.99	0.44
1:A:244:ARG:C	1:A:246:ASN:H	2.21	0.44
1:A:124:MET:SD	1:A:127:ARG:HD3	2.57	0.44
1:A:194:MET:HA	1:A:197:SER:CB	2.48	0.44
1:A:364:CYS:HB3	1:A:371:PHE:CG	2.52	0.44
1:A:55:LEU:HB2	1:A:78:LEU:HD21	1.99	0.44
1:A:502:MET:HG2	1:A:507:ALA:HB2	1.99	0.44
1:A:360:ALA:HB1	1:A:392:PRO:HG2	2.00	0.44
1:A:263:VAL:HG22	1:A:269:HIS:CD2	2.53	0.43
1:A:48:LEU:CD1	1:A:52:LYS:HE3	2.48	0.43
1:A:48:LEU:HD11	1:A:52:LYS:HE3	2.00	0.43
1:A:16:ILE:HA	1:A:219:ILE:O	2.17	0.43
1:A:323:GLU:HG3	1:A:384:PHE:HB3	2.00	0.43
1:A:27:LEU:HB3	1:A:139:LEU:CD1	2.49	0.42
1:A:447:VAL:HG12	1:A:449:PHE:N	2.32	0.42
1:A:123:TYR:O	1:A:126:GLU:HB3	2.19	0.42
1:A:311:GLU:OE1	1:A:376:ARG:HD3	2.20	0.42
1:A:512:ALA:O	1:A:516:LEU:HG	2.19	0.42
1:A:21:LEU:HD12	1:A:21:LEU:N	2.35	0.42
1:A:107:ASP:O	1:A:111:GLN:HG2	2.20	0.41
1:A:547:LYS:HE3	1:A:547:LYS:HB2	1.87	0.41
1:A:524:PHE:CD1	1:A:524:PHE:N	2.88	0.41
1:A:296:TYR:HB3	1:A:332:VAL:HA	2.01	0.41
1:A:469:GLU:OE1	1:A:469:GLU:HA	2.20	0.41
1:A:249:LYS:O	1:A:253:GLU:HG3	2.20	0.41
1:A:485:LEU:O	1:A:489:VAL:HG23	2.21	0.41
1:A:291:GLU:O	1:A:328:LYS:HG3	2.20	0.41
1:A:9:ALA:O	1:A:202:ALA:HB3	2.21	0.41
1:A:413:ASN:O	1:A:417:GLU:HG3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:284:GLY:HA2	1:A:287:GLU:HG2	2.03	0.40
1:A:375:LEU:HD22	1:A:407:LEU:CD1	2.51	0.40
1:A:410:GLU:HA	1:A:410:GLU:OE1	2.21	0.40
1:A:441:ASP:OD2	1:A:442:ILE:N	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	537/573 (94%)	460 (86%)	64 (12%)	13 (2%)	6 9

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	221	VAL
1	A	342	GLU
1	A	12	ASP
1	A	18	LYS
1	A	20	TYR
1	A	170	ALA
1	A	351	GLU
1	A	14	VAL
1	A	177	VAL
1	A	181	VAL
1	A	16	ILE
1	A	214	GLU
1	A	5	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	391/481 (81%)	372 (95%)	19 (5%)	25 47

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	97	GLU
1	A	100	ASN
1	A	147	LEU
1	A	241	GLN
1	A	242	GLU
1	A	250	ASP
1	A	252	GLN
1	A	295	LEU
1	A	296	TYR
1	A	301	LEU
1	A	322	LEU
1	A	328	LYS
1	A	342	GLU
1	A	347	ASP
1	A	379	LEU
1	A	392	PRO
1	A	436	THR
1	A	532	LEU
1	A	543	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	31	ASN
1	A	65	GLN
1	A	100	ASN
1	A	103	GLN
1	A	121	ASN
1	A	353	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	416	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	900	-	4,4,4	0.22	0	6,6,6	0.10	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/573 (96%)	0.68	77 (13%) 2 2	29, 58, 132, 139	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	15	ALA	8.2
1	A	161	GLY	7.2
1	A	123	TYR	5.7
1	A	191	SER	5.6
1	A	202	ALA	5.6
1	A	203	VAL	5.5
1	A	9	ALA	5.5
1	A	180	PHE	5.4
1	A	5	ILE	4.8
1	A	215	ALA	4.8
1	A	164	LEU	4.6
1	A	192	ALA	4.4
1	A	169	THR	4.4
1	A	10	ALA	4.3
1	A	64	LYS	4.2
1	A	65	GLN	4.1
1	A	7	GLY	4.1
1	A	167	SER	3.9
1	A	165	THR	3.8
1	A	19	ALA	3.8
1	A	31	ASN	3.8
1	A	21	LEU	3.8
1	A	185	GLY	3.7
1	A	246	ASN	3.5
1	A	208	SER	3.5
1	A	431	VAL	3.4
1	A	240	TYR	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	216	GLY	3.2
1	A	239	GLU	3.2
1	A	181	VAL	3.2
1	A	184	ILE	3.2
1	A	58	ILE	3.1
1	A	172	LEU	3.1
1	A	430	MET	3.0
1	A	193	ILE	3.0
1	A	71	ALA	3.0
1	A	238	ALA	3.0
1	A	249	LYS	3.0
1	A	74	PHE	3.0
1	A	66	LEU	2.9
1	A	8	ILE	2.9
1	A	221	VAL	2.8
1	A	453	GLY	2.8
1	A	452	ILE	2.7
1	A	14	VAL	2.7
1	A	68	ALA	2.7
1	A	242	GLU	2.7
1	A	198	LEU	2.6
1	A	205	GLY	2.6
1	A	457	LEU	2.6
1	A	182	THR	2.6
1	A	70	LYS	2.6
1	A	162	ASN	2.6
1	A	183	ASN	2.6
1	A	166	PRO	2.5
1	A	195	SER	2.4
1	A	12	ASP	2.4
1	A	173	ASN	2.4
1	A	120	ASP	2.4
1	A	437	ALA	2.4
1	A	149	ASN	2.3
1	A	160	ILE	2.3
1	A	189	SER	2.3
1	A	502	MET	2.3
1	A	147	LEU	2.3
1	A	98	SER	2.3
1	A	438	ALA	2.3
1	A	150	PRO	2.3
1	A	454	THR	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	176	TYR	2.2
1	A	306	ASP	2.2
1	A	436	THR	2.1
1	A	6	LYS	2.1
1	A	571	LYS	2.1
1	A	33	SER	2.0
1	A	433	ILE	2.0
1	A	525	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	900	5/5	0.96	0.29	63,64,66,66	0

6.5 Other polymers [i](#)

There are no such residues in this entry.