



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 27, 2023 – 02:13 PM EDT

PDB ID : 3HTU
Title : Crystal structure of the human VPS25-VPS20 subcomplex
Authors : Im, Y.J.; Hurley, J.H.
Deposited on : 2009-06-12
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	79	 8% 70% 23% • 5%
1	C	79	 5% 66% 23% 6% 5%
1	E	79	 10% 76% 19% 5%
1	G	79	 10% 75% 18% • 5%
2	B	39	 3% 69% 21% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	39	
2	F	39	
2	H	39	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3870 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vacuolar protein-sorting-associated protein 25.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	75	616	386	107	123	0	0	0
1	C	75	616	386	107	123	0	0	0
1	E	75	616	386	107	123	0	0	0
1	G	75	616	386	107	123	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	98	GLY	-	expression tag	UNP Q9BRG1
A	99	ALA	-	expression tag	UNP Q9BRG1
A	100	MET	-	expression tag	UNP Q9BRG1
A	101	GLY	-	expression tag	UNP Q9BRG1
C	98	GLY	-	expression tag	UNP Q9BRG1
C	99	ALA	-	expression tag	UNP Q9BRG1
C	100	MET	-	expression tag	UNP Q9BRG1
C	101	GLY	-	expression tag	UNP Q9BRG1
E	98	GLY	-	expression tag	UNP Q9BRG1
E	99	ALA	-	expression tag	UNP Q9BRG1
E	100	MET	-	expression tag	UNP Q9BRG1
E	101	GLY	-	expression tag	UNP Q9BRG1
G	98	GLY	-	expression tag	UNP Q9BRG1
G	99	ALA	-	expression tag	UNP Q9BRG1
G	100	MET	-	expression tag	UNP Q9BRG1
G	101	GLY	-	expression tag	UNP Q9BRG1

- Molecule 2 is a protein called Vacuolar protein-sorting-associated protein 20.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	35	Total	C	N	O	0	0	0
			300	182	62	56			
2	D	36	Total	C	N	O	0	0	0
			311	188	66	57			
2	F	32	Total	C	N	O	0	0	0
			271	166	56	49			
2	H	36	Total	C	N	O	0	0	0
			311	188	66	57			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	GLY	-	expression tag	UNP Q96FZ7
D	10	GLY	-	expression tag	UNP Q96FZ7
F	10	GLY	-	expression tag	UNP Q96FZ7
H	10	GLY	-	expression tag	UNP Q96FZ7

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	38	Total	O	0	0
			38	38		
3	B	19	Total	O	0	0
			19	19		
3	C	43	Total	O	0	0
			43	43		
3	D	16	Total	O	0	0
			16	16		
3	E	23	Total	O	0	0
			23	23		
3	F	22	Total	O	0	0
			22	22		
3	G	21	Total	O	0	0
			21	21		
3	H	31	Total	O	0	0
			31	31		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vacuolar protein-sorting-associated protein 25

Chain A: 





- Molecule 1: Vacuolar protein-sorting-associated protein 25

Chain C: 




- Molecule 1: Vacuolar protein-sorting-associated protein 25

Chain E: 



- Molecule 1: Vacuolar protein-sorting-associated protein 25

Chain G: 




- Molecule 2: Vacuolar protein-sorting-associated protein 20


Chain B: 




- Molecule 2: Vacuolar protein-sorting-associated protein 20



- Molecule 2: Vacuolar protein-sorting-associated protein 20



- Molecule 2: Vacuolar protein-sorting-associated protein 20



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.56Å 51.04Å 76.99Å 90.00° 90.39° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 38.48 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.4 (50.00-2.00) 98.4 (38.48-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.70 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.239 , 0.273 0.256 , 0.282	Depositor DCC
R_{free} test set	1544 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	32.2	Xtrriage
Anisotropy	0.619	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3870	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.53	0/628	0.63	0/846
1	C	0.56	0/628	0.66	0/846
1	E	0.52	0/628	0.63	0/846
1	G	0.52	0/628	0.62	0/846
2	B	0.65	0/300	0.71	0/396
2	D	0.62	0/311	0.70	0/410
2	F	0.59	0/271	0.83	1/358 (0.3%)
2	H	0.66	0/311	0.76	0/410
All	All	0.56	0/3705	0.67	1/4958 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	27	ARG	NE-CZ-NH1	5.10	122.85	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	616	0	582	10	0
1	C	616	0	582	16	0
1	E	616	0	582	7	0
1	G	616	0	582	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	300	0	315	5	0
2	D	311	0	328	5	0
2	F	271	0	290	10	0
2	H	311	0	328	12	0
3	A	38	0	0	0	0
3	B	19	0	0	0	0
3	C	43	0	0	2	0
3	D	16	0	0	1	0
3	E	23	0	0	1	0
3	F	22	0	0	0	0
3	G	21	0	0	0	0
3	H	31	0	0	1	0
All	All	3870	0	3589	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:139:GLU:HG2	1:C:141:PHE:CE2	2.10	0.86
2:F:30:LEU:HB3	2:H:34:GLN:HE21	1.46	0.79
1:C:169:ASP:HB3	3:D:51:HOH:O	1.85	0.77
1:G:124:VAL:HG12	2:H:21:LEU:HD23	1.70	0.74
2:F:30:LEU:CB	2:H:34:GLN:HE21	2.07	0.67
1:A:137:GLU:HA	1:A:142:HIS:CD2	2.31	0.65
1:C:102:ARG:HD3	3:C:47:HOH:O	1.97	0.65
2:H:34:GLN:HG3	3:H:113:HOH:O	1.96	0.64
1:A:116:SER:O	1:A:120:GLN:HG3	1.97	0.64
2:F:34:GLN:HE22	2:H:27:ARG:HG3	1.63	0.63
1:C:150:LEU:O	1:C:154:GLN:HB3	2.01	0.61
1:C:111:ILE:HA	1:C:141:PHE:CE2	2.36	0.61
2:F:27:ARG:HH11	2:F:27:ARG:HG2	1.65	0.60
1:A:117:ARG:O	1:A:118:SER:HB3	2.02	0.59
1:C:139:GLU:HG2	1:C:141:PHE:CZ	2.37	0.59
1:G:172:GLY:HA3	2:H:21:LEU:HD11	1.85	0.58
1:G:136:THR:O	1:G:139:GLU:HB2	2.03	0.58
1:A:133:GLY:O	1:A:137:GLU:HG3	2.06	0.55
1:C:103:ARG:H	1:C:103:ARG:HE	1.52	0.55
1:A:136:THR:HG23	1:A:142:HIS:HD2	1.71	0.55
2:D:25:GLN:O	2:D:29:LYS:HG3	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:34:GLN:NE2	2:H:27:ARG:HG3	2.22	0.54
1:C:121:ASN:O	1:C:122:ASN:HB2	2.07	0.54
2:H:26:GLN:HE22	2:H:29:LYS:HD2	1.74	0.53
1:C:103:ARG:H	1:C:103:ARG:NE	2.08	0.52
2:D:37:ILE:HA	2:D:40:GLN:HE21	1.74	0.52
1:C:136:THR:HG23	1:C:142:HIS:HD2	1.76	0.51
1:C:167:VAL:HG22	2:D:21:LEU:HD13	1.91	0.51
2:F:27:ARG:HH11	2:F:27:ARG:CG	2.23	0.51
1:C:157:GLN:HG2	3:C:16:HOH:O	2.10	0.50
1:C:157:GLN:HE22	1:C:164:ILE:H	1.62	0.48
1:A:118:SER:O	1:A:120:GLN:N	2.46	0.47
1:A:141:PHE:O	1:A:144:LEU:HB2	2.13	0.47
1:G:134:GLU:HA	1:G:137:GLU:HG3	1.96	0.47
2:B:31:ARG:O	2:B:35:LYS:HG3	2.15	0.47
2:H:29:LYS:HG2	2:H:33:TYR:CE2	2.50	0.46
1:A:130:LEU:O	1:A:136:THR:HG21	2.16	0.46
2:F:27:ARG:HG2	2:F:27:ARG:NH1	2.31	0.45
1:A:136:THR:HG23	1:A:142:HIS:CD2	2.51	0.45
2:F:22:GLN:O	2:F:26:GLN:HG3	2.16	0.45
2:B:33:TYR:CE1	2:B:37:ILE:HD11	2.53	0.44
1:E:122:ASN:HD22	1:E:176:PHE:HA	1.83	0.44
2:H:36:ARG:O	2:H:39:GLN:HG2	2.18	0.44
1:E:131:THR:O	1:E:142:HIS:CD2	2.71	0.44
1:E:106:GLU:O	1:E:110:LEU:HG	2.19	0.43
2:B:26:GLN:O	2:B:30:LEU:HG	2.19	0.43
1:G:157:GLN:HE21	1:G:163:GLU:HA	1.84	0.42
2:D:42:GLU:HA	2:D:42:GLU:OE1	2.19	0.42
1:E:119:GLY:HA2	3:E:28:HOH:O	2.18	0.42
1:E:137:GLU:HA	1:E:142:HIS:CG	2.55	0.41
2:B:41:LEU:HA	2:B:41:LEU:HD23	1.79	0.41
1:C:113:GLN:O	1:C:117:ARG:HB3	2.20	0.41
1:C:144:LEU:CD2	1:C:148:THR:HB	2.50	0.41
2:F:31:ARG:HG3	2:H:34:GLN:HE22	1.85	0.41
2:B:33:TYR:CD2	2:D:30:LEU:HD11	2.56	0.41
1:E:122:ASN:HA	1:E:175:PHE:O	2.20	0.41
1:G:103:ARG:HB2	1:G:104:PRO:HD2	2.02	0.40
1:C:154:GLN:O	1:C:154:GLN:HG2	2.22	0.40
1:E:165:ILE:HG21	2:F:21:LEU:HD12	2.04	0.40
1:A:102:ARG:HG2	1:A:107:TRP:CE2	2.56	0.40
2:H:26:GLN:NE2	2:H:29:LYS:HD2	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	73/79 (92%)	68 (93%)	2 (3%)	3 (4%)	3	1
1	C	73/79 (92%)	68 (93%)	5 (7%)	0	100	100
1	E	73/79 (92%)	70 (96%)	3 (4%)	0	100	100
1	G	73/79 (92%)	70 (96%)	2 (3%)	1 (1%)	11	5
2	B	33/39 (85%)	33 (100%)	0	0	100	100
2	D	34/39 (87%)	33 (97%)	1 (3%)	0	100	100
2	F	30/39 (77%)	30 (100%)	0	0	100	100
2	H	34/39 (87%)	33 (97%)	0	1 (3%)	4	1
All	All	423/472 (90%)	405 (96%)	13 (3%)	5 (1%)	13	7

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	118	SER
1	G	168	SER
2	H	11	SER
1	A	119	GLY
1	A	170	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	65/66 (98%)	59 (91%)	6 (9%)	9	5

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	65/66 (98%)	56 (86%)	9 (14%)	3	2
1	E	65/66 (98%)	60 (92%)	5 (8%)	13	8
1	G	65/66 (98%)	58 (89%)	7 (11%)	6	3
2	B	32/34 (94%)	31 (97%)	1 (3%)	40	40
2	D	33/34 (97%)	31 (94%)	2 (6%)	18	14
2	F	29/34 (85%)	24 (83%)	5 (17%)	2	1
2	H	33/34 (97%)	29 (88%)	4 (12%)	5	2
All	All	387/400 (97%)	348 (90%)	39 (10%)	7	4

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	123	SER
1	A	139	GLU
1	A	144	LEU
1	A	146	GLU
1	A	153	LEU
1	A	156	LEU
2	B	16	GLN
1	C	103	ARG
1	C	105	GLU
1	C	110	LEU
1	C	117	ARG
1	C	121	ASN
1	C	123	SER
1	C	135	ASP
1	C	144	LEU
1	C	150	LEU
2	D	21	LEU
2	D	44	GLU
1	E	105	GLU
1	E	123	SER
1	E	136	THR
1	E	150	LEU
1	E	151	ARG
2	F	13	VAL
2	F	21	LEU
2	F	22	GLN
2	F	27	ARG
2	F	30	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	103	ARG
1	G	127	LEU
1	G	139	GLU
1	G	144	LEU
1	G	150	LEU
1	G	153	LEU
1	G	156	LEU
2	H	11	SER
2	H	21	LEU
2	H	26	GLN
2	H	36	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	132	ASN
1	C	142	HIS
1	C	157	GLN
2	D	40	GLN
1	E	113	GLN
2	F	22	GLN
2	F	26	GLN
2	F	34	GLN
1	G	157	GLN
2	H	26	GLN
2	H	34	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	75/79 (94%)	0.61	6 (8%) 12 11	18, 30, 58, 60	0
1	C	75/79 (94%)	0.71	4 (5%) 26 25	14, 29, 44, 53	0
1	E	75/79 (94%)	0.84	8 (10%) 6 5	18, 34, 55, 61	0
1	G	75/79 (94%)	0.92	8 (10%) 6 5	18, 31, 57, 70	0
2	B	35/39 (89%)	0.39	1 (2%) 51 50	16, 24, 46, 52	0
2	D	36/39 (92%)	0.55	1 (2%) 53 51	13, 20, 54, 61	0
2	F	32/39 (82%)	0.50	2 (6%) 20 19	20, 27, 41, 42	0
2	H	36/39 (92%)	0.63	2 (5%) 24 23	17, 23, 49, 52	0
All	All	439/472 (93%)	0.69	32 (7%) 15 14	13, 29, 54, 70	0

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	169	ASP	9.0
1	E	122	ASN	6.2
1	A	118	SER	5.4
1	E	118	SER	4.7
1	G	122	ASN	4.5
1	C	138	ASP	4.2
2	F	33	TYR	4.1
1	E	103	ARG	3.6
2	H	10	GLY	3.5
1	G	168	SER	3.5
2	F	39	GLN	3.3
1	A	122	ASN	3.0
1	E	110	LEU	3.0
1	E	121	ASN	2.9
1	E	169	ASP	2.8
1	A	119	GLY	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	112	TYR	2.6
1	E	143	GLY	2.5
1	A	117	ARG	2.4
1	G	104	PRO	2.4
1	C	122	ASN	2.3
2	H	33	TYR	2.3
1	C	141	PHE	2.2
1	G	103	ARG	2.2
1	A	140	GLU	2.2
1	G	107	TRP	2.2
1	G	141	PHE	2.1
2	B	43	ARG	2.1
1	A	121	ASN	2.1
1	G	121	ASN	2.1
2	D	43	ARG	2.1
1	E	168	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.