



wwPDB X-ray Structure Validation Summary Report ⓘ

Jul 13, 2026 – 01:56 pm BST

PDB ID : 9HYL / pdb_00009hyl
Title : Crystal structure of Trypanosoma brucei trypanothione reductase (TbTR)
bound to PROTAC-3
Authors : Exertier, C.; Ilari, C.; Fiorillo, A.; Antonelli, C.
Deposited on : 2025-01-10
Resolution : 2.08 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.015 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	495	87% 11% ..
1	B	495	88% 10% .
1	C	495	89% 9% ..
1	D	495	86% 12% ..

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 16162 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trypanothione reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	489	Total	C	N	O	S	0	6	0
			3761	2396	642	703	20			
1	B	487	Total	C	N	O	S	0	8	0
			3747	2385	639	704	19			
1	C	490	Total	C	N	O	S	0	2	0
			3733	2374	634	705	20			
1	D	490	Total	C	N	O	S	0	5	0
			3748	2384	637	707	20			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q389T8
A	-1	SER	-	expression tag	UNP Q389T8
A	0	HIS	-	expression tag	UNP Q389T8
B	-2	GLY	-	expression tag	UNP Q389T8
B	-1	SER	-	expression tag	UNP Q389T8
B	0	HIS	-	expression tag	UNP Q389T8
C	-2	GLY	-	expression tag	UNP Q389T8
C	-1	SER	-	expression tag	UNP Q389T8
C	0	HIS	-	expression tag	UNP Q389T8
D	-2	GLY	-	expression tag	UNP Q389T8
D	-1	SER	-	expression tag	UNP Q389T8
D	0	HIS	-	expression tag	UNP Q389T8

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	C	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	D	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is N-(2-(2-(2-((S)-6-chloro-2-methyl-4-phenylquinazolin-3(4H)-yl)ethoxy)ethoxy)ethyl)-2-(2-(2-((2-((3 {R})-(2,6-dioxopiperidin-3-yl)-1,3-dioxoisindolin-4-yl)amino)ethoxy)ethoxy)acetamide (CCD ID: A1IYD) (formula: C₄₀H₄₅ClN₆O₉) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 56	C 40	Cl 1	N 6	O 9	0	0
3	B	1	Total 56	C 40	Cl 1	N 6	O 9	0	0
3	C	1	Total 56	C 40	Cl 1	N 6	O 9	0	0
3	D	1	Total 56	C 40	Cl 1	N 6	O 9	0	0

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	A	1	Total C O 7 4 3	0	0
4	C	1	Total C O 7 4 3	0	0
4	D	1	Total C O 7 4 3	0	0
4	D	1	Total C O 7 4 3	0	0

- Molecule 5 is IMIDAZOLE (CCD ID: IMD) (formula: $C_3H_5N_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N 5 3 2	0	0
5	C	1	Total C N 5 3 2	0	0

- Molecule 6 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 7 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Na	0	0
			2	2		
7	B	3	Total	Na	0	0
			3	3		
7	C	1	Total	Na	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	186	Total	O	0	0
			186	186		
8	B	143	Total	O	0	0
			143	143		
8	C	157	Total	O	0	0
			157	157		
8	D	175	Total	O	0	0
			175	175		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Trypanothione reductase




• Molecule 1: Trypanothione reductase



• Molecule 1: Trypanothione reductase



• Molecule 1: Trypanothione reductase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	101.32Å 63.61Å 169.27Å 90.00° 98.23° 90.00°	Depositor
Resolution (Å)	46.29 – 2.08 46.29 – 2.08	Depositor EDS
% Data completeness (in resolution range)	98.0 (46.29-2.08) 98.2 (46.29-2.08)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.178 , 0.224 0.189 , 0.229	Depositor DCC
R_{free} test set	6307 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å ²)	35.4	Xtriage
Anisotropy	0.072	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 26.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	16162	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FAD, PEG, IMD, A1IYD, NA, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.02	0/3859	1.35	4/5231 (0.1%)
1	B	1.02	0/3849	1.36	8/5219 (0.2%)
1	C	1.03	1/3818 (0.0%)	1.33	1/5180 (0.0%)
1	D	1.05	0/3838	1.36	5/5204 (0.1%)
All	All	1.03	1/15364 (0.0%)	1.35	18/20834 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	423	THR	C-O	6.28	1.31	1.23

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	458	ILE	CA-C-N	6.70	126.46	120.10
1	A	458	ILE	C-N-CA	6.70	126.46	120.10
1	B	263	ASP	CA-CB-CG	6.24	118.84	112.60
1	D	485	GLU	N-CA-C	-6.01	106.20	112.93
1	D	42	PRO	O-C-N	-5.75	118.67	121.31

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	228	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3761	0	3794	38	0
1	B	3747	0	3779	33	0
1	C	3733	0	3744	26	0
1	D	3748	0	3767	38	0
2	A	53	0	31	1	0
2	B	53	0	31	0	0
2	C	53	0	31	0	0
2	D	53	0	31	0	0
3	A	56	0	0	0	0
3	B	56	0	0	0	0
3	C	56	0	0	0	0
3	D	56	0	0	0	0
4	A	35	0	50	3	0
4	C	7	0	10	1	0
4	D	14	0	20	4	0
5	A	5	0	5	0	0
5	C	5	0	5	0	0
6	A	4	0	6	1	0
7	A	2	0	0	0	0
7	B	3	0	0	0	0
7	C	1	0	0	0	0
8	A	186	0	0	1	0
8	B	143	0	0	1	0
8	C	157	0	0	0	0
8	D	175	0	0	2	0
All	All	16162	0	15304	131	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 131 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:82:GLU:OE1	1:B:89:LYS:HE3	1.45	1.13
1:B:320:PRO:O	1:B:321[B]:ASN:ND2	1.97	0.96
1:B:331:ARG:NH1	1:B:359:HIS:HB2	1.87	0.89
1:A:82:GLU:OE1	1:B:89:LYS:CE	2.29	0.78
1:B:130:GLU:HB2	1:B:136:VAL:HG23	1.64	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	493/495 (100%)	482 (98%)	11 (2%)	0	100	100
1	B	493/495 (100%)	473 (96%)	20 (4%)	0	100	100
1	C	490/495 (99%)	474 (97%)	16 (3%)	0	100	100
1	D	493/495 (100%)	481 (98%)	12 (2%)	0	100	100
All	All	1969/1980 (99%)	1910 (97%)	59 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	408/407 (100%)	399 (98%)	9 (2%)	45	51
1	B	408/407 (100%)	394 (97%)	14 (3%)	32	34

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	405/407 (100%)	390 (96%)	15 (4%)	30	31
1	D	407/407 (100%)	384 (94%)	23 (6%)	18	16
All	All	1628/1628 (100%)	1567 (96%)	61 (4%)	32	31

5 of 61 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	262	THR
1	D	353	LYS
1	C	384	LYS
1	D	335	THR
1	D	485	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 24 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	245	ASN
1	C	292	ASN
1	C	267	HIS
1	C	318	ASN
1	B	223	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 6 are monoatomic - leaving 19 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	A1IYD	B	502	-	61,61,61	0.70	1 (1%)	78,83,83	0.78	3 (3%)
4	PEG	A	505	-	6,6,6	0.37	0	5,5,5	0.43	0
4	PEG	C	504	-	6,6,6	0.35	0	5,5,5	0.20	0
2	FAD	D	501	-	56,58,58	0.61	0	81,89,89	0.87	3 (3%)
3	A1IYD	A	502	-	61,61,61	0.94	2 (3%)	78,83,83	1.11	4 (5%)
2	FAD	C	501	-	56,58,58	0.54	0	81,89,89	0.73	2 (2%)
2	FAD	B	501	-	56,58,58	0.53	0	81,89,89	0.78	1 (1%)
4	PEG	D	503	-	6,6,6	0.38	0	5,5,5	0.49	0
4	PEG	A	503	-	6,6,6	0.22	0	5,5,5	0.17	0
5	IMD	C	503	-	5,5,5	0.16	0	5,5,5	0.49	0
4	PEG	A	508	-	6,6,6	0.26	0	5,5,5	0.26	0
3	A1IYD	C	502	-	61,61,61	0.69	1 (1%)	78,83,83	0.79	4 (5%)
4	PEG	A	509	-	6,6,6	0.25	0	5,5,5	0.24	0
6	DMS	A	507	-	3,3,3	0.38	0	3,3,3	0.34	0
4	PEG	A	506	-	6,6,6	0.12	0	5,5,5	0.13	0
2	FAD	A	501	-	56,58,58	0.69	1 (1%)	81,89,89	0.79	3 (3%)
4	PEG	D	504	-	6,6,6	0.33	0	5,5,5	0.30	0
3	A1IYD	D	502	-	61,61,61	0.63	1 (1%)	78,83,83	0.84	4 (5%)
5	IMD	A	504	-	5,5,5	0.23	0	5,5,5	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1IYD	B	502	-	-	16/31/76/76	0/6/6/6
4	PEG	A	506	-	-	3/4/4/4	-
4	PEG	A	505	-	-	0/4/4/4	-
4	PEG	D	503	-	-	3/4/4/4	-
4	PEG	C	504	-	-	4/4/4/4	-
4	PEG	A	503	-	-	0/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	508	-	-	2/4/4/4	-
5	IMD	C	503	-	-	-	0/1/1/1
2	FAD	D	501	-	-	4/34/50/50	0/6/6/6
2	FAD	A	501	-	-	5/34/50/50	0/6/6/6
3	A1IYD	C	502	-	-	19/31/76/76	0/6/6/6
3	A1IYD	A	502	-	-	16/31/76/76	0/6/6/6
4	PEG	A	509	-	-	4/4/4/4	-
4	PEG	D	504	-	-	3/4/4/4	-
3	A1IYD	D	502	-	-	14/31/76/76	0/6/6/6
5	IMD	A	504	-	-	-	0/1/1/1
2	FAD	C	501	-	-	5/34/50/50	0/6/6/6
2	FAD	B	501	-	-	3/34/50/50	0/6/6/6

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	A1IYD	C26-N6	4.69	1.35	1.29
3	B	502	A1IYD	C26-N6	4.52	1.35	1.29
3	A	502	A1IYD	O9-C23	4.50	1.61	1.42
3	C	502	A1IYD	C26-N6	4.33	1.35	1.29
3	D	502	A1IYD	C26-N6	3.90	1.34	1.29

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	A1IYD	C14-C13-N2	6.28	114.81	109.08
3	D	502	A1IYD	C14-C13-N2	3.72	112.47	109.08
3	B	502	A1IYD	C11-C18-C7	-3.58	119.44	121.91
3	A	502	A1IYD	C27-C26-N6	-3.46	114.87	119.34
3	D	502	A1IYD	C11-C18-C7	-3.45	119.53	121.91

There are no chirality outliers.

5 of 101 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	501	FAD	PA-O3P-P-O5'
3	A	502	A1IYD	C2-C1-N4-C20
3	A	502	A1IYD	C5-C6-N1-C7
3	A	502	A1IYD	C8-C7-N1-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	502	A1IYD	C18-C7-N1-C6


There are no ring outliers.

7 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	505	PEG	1	0
4	C	504	PEG	1	0
4	D	503	PEG	2	0
4	A	509	PEG	2	0
6	A	507	DMS	1	0
2	A	501	FAD	1	0
4	D	504	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1IYD B 502




Ligand FAD D 501






Ligand A1IYD A 502



Ligand FAD C 501



Ligand FAD B 501**Ligand A1IYD C 502**



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	489/495 (98%)	-0.13	3 (0%) 85 87	17, 33, 50, 78	6 (1%)
1	B	487/495 (98%)	-0.02	6 (1%) 76 78	17, 35, 54, 82	8 (1%)
1	C	490/495 (98%)	-0.00	12 (2%) 59 62	23, 35, 57, 99	2 (0%)
1	D	490/495 (98%)	-0.14	3 (0%) 85 87	16, 33, 52, 91	5 (1%)
All	All	1956/1980 (98%)	-0.08	24 (1%) 76 78	16, 34, 54, 99	21 (1%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	489	ASP	4.8
1	B	331	ARG	4.5
1	D	487	LEU	4.1
1	D	486	LYS	3.8
1	C	352	ASN	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	DMS	A	507	4/4	0.62	0.39	43,48,49,54	4
4	PEG	D	504	7/7	0.66	0.23	63,68,73,76	0
4	PEG	A	508	7/7	0.66	0.19	71,78,79,82	0
4	PEG	A	505	7/7	0.75	0.16	60,67,72,74	0
4	PEG	D	503	7/7	0.76	0.20	65,68,71,74	0
5	IMD	A	504	5/5	0.78	0.23	73,73,77,80	0
5	IMD	C	503	5/5	0.81	0.20	75,75,76,81	0
3	A1IYD	C	502	56/56	0.81	0.29	41,141,175,178	35
3	A1IYD	A	502	56/56	0.82	0.30	33,121,168,178	35
4	PEG	A	509	7/7	0.82	0.17	53,60,66,67	0
4	PEG	C	504	7/7	0.82	0.17	58,61,64,67	0
4	PEG	A	506	7/7	0.82	0.26	42,42,43,44	7
3	A1IYD	D	502	56/56	0.83	0.28	30,132,185,189	35
3	A1IYD	B	502	56/56	0.84	0.29	34,130,177,182	35
4	PEG	A	503	7/7	0.84	0.19	51,65,81,83	0
7	NA	A	511	1/1	0.87	0.19	49,49,49,49	0
7	NA	B	503	1/1	0.93	0.30	51,51,51,51	0
2	FAD	C	501	53/53	0.97	0.05	25,30,34,36	0
2	FAD	B	501	53/53	0.97	0.06	25,29,33,34	0
7	NA	B	504	1/1	0.97	0.08	31,31,31,31	0
7	NA	C	505	1/1	0.97	0.12	31,31,31,31	0
2	FAD	A	501	53/53	0.98	0.05	22,26,31,32	0
7	NA	A	510	1/1	0.98	0.04	24,24,24,24	0
7	NA	B	505	1/1	0.98	0.05	34,34,34,34	0
2	FAD	D	501	53/53	0.98	0.04	20,25,28,29	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1IYD C 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1IYD A 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1IYD D 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1IYD B 502:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD C 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FAD B 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FAD A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FAD D 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.