



# Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 04:16 am BST

PDB ID : 4I0O  
Title : Nucleoporin ELYS (aa1-494), *Mus musculus*  
Authors : Bilokapic, S.; Schwartz, T.U.  
Deposited on : 2012-11-17  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	497	

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3798 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein ELYS.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	463	3629	2304	618	689	14	4	0	7	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PRO	-	EXPRESSION TAG	UNP Q8CJF7
A	-1	GLY	-	EXPRESSION TAG	UNP Q8CJF7
A	0	SER	-	EXPRESSION TAG	UNP Q8CJF7


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	169	Total	O	0	0
			169	169		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein ELYS



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	52.60Å 78.12Å 58.85Å 90.00° 106.59° 90.00°	Depositor
Resolution (Å)	26.52 – 1.90 26.52 – 1.90	Depositor EDS
% Data completeness (in resolution range)	98.5 (26.52-1.90) 98.8 (26.52-1.90)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.45 (at 1.91Å)	Xtrriage
Refinement program	PHENIX 1.8_1063	Depositor
R, $R_{free}$	0.180 , 0.214 0.181 , 0.215	Depositor DCC
$R_{free}$ test set	1783 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.8	Xtrriage
Anisotropy	0.266	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3798	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	2/3722 (0.1%)	0.68	1/5056 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	354	CYS	CB-SG	-7.06	1.70	1.82
1	A	107	CYS	CB-SG	-5.66	1.72	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	30	LEU	CA-CB-CG	6.03	129.18	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3629	0	3591	42	0
2	A	169	0	0	5	0
All	All	3798	0	3591	42	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (42) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:27:GLU:HA	1:A:30:LEU:HD13	1.58	0.85
1:A:417:ARG:NH2	2:A:613:HOH:O	2.17	0.75
1:A:274[B]:GLU:OE2	2:A:637:HOH:O	2.05	0.75
1:A:314:LEU:HD11	1:A:320[B]:LEU:HG	1.75	0.68
1:A:30:LEU:HD12	1:A:31:GLU:HA	1.77	0.67
1:A:390:TYR:O	2:A:643:HOH:O	2.14	0.65
1:A:352:LEU:O	2:A:582:HOH:O	2.15	0.64
1:A:384[A]:THR:HG21	1:A:448:ILE:HD12	1.79	0.63
1:A:30:LEU:HA	1:A:32:SER:H	1.65	0.61
1:A:352:LEU:HB2	1:A:384[B]:THR:HG23	1.87	0.56
1:A:256:TYR:OH	1:A:258:THR:OG1	2.24	0.56
1:A:26:ASP:HB2	1:A:50:CYS:SG	2.47	0.55
1:A:30:LEU:CG	1:A:31:GLU:HA	2.37	0.54
1:A:36:GLY:O	1:A:37:LYS:HD3	2.08	0.54
1:A:29:THR:O	1:A:32:SER:OG	2.25	0.54
1:A:314:LEU:HD21	1:A:320[B]:LEU:HD12	1.89	0.53
1:A:30:LEU:CD1	1:A:31:GLU:HA	2.39	0.52
1:A:26:ASP:OD1	1:A:27:GLU:HG3	2.11	0.51
1:A:422:LEU:HD22	1:A:428:PHE:CD2	2.46	0.51
1:A:24:GLY:C	1:A:26:ASP:H	2.15	0.50
1:A:30:LEU:HG	1:A:31:GLU:HG2	1.92	0.50
1:A:272:PHE:CE2	1:A:274[B]:GLU:HG3	2.46	0.50
1:A:81[B]:LEU:HG	1:A:99:GLU:HB3	1.93	0.50
1:A:252:MSE:HE3	1:A:252:MSE:HA	1.94	0.48
1:A:17:GLU:O	1:A:21:GLU:HG3	2.13	0.48
1:A:35:ARG:NH2	1:A:80:VAL:O	2.46	0.47
1:A:28:ILE:C	1:A:30:LEU:H	2.18	0.47
1:A:318:GLN:HG3	2:A:668:HOH:O	2.15	0.46
1:A:30:LEU:HG	1:A:31:GLU:CG	2.47	0.45
1:A:88:TRP:CZ2	1:A:91:ARG:HD2	2.51	0.45
1:A:4:LEU:HB3	1:A:431:TRP:CE3	2.51	0.45
1:A:231:ARG:NE	1:A:274[B]:GLU:OE2	2.43	0.45
1:A:359:ARG:HB3	1:A:378:THR:HB	1.98	0.45
1:A:406:ARG:HD2	1:A:406:ARG:HA	1.71	0.45
1:A:132:ILE:HG21	1:A:249:MSE:HE2	1.99	0.44
1:A:294:SER:OG	1:A:295:GLU:N	2.49	0.44
1:A:30:LEU:HG	1:A:31:GLU:HA	1.98	0.44
1:A:27:GLU:HG2	1:A:30:LEU:HD13	2.00	0.44
1:A:29:THR:HG21	1:A:52:PRO:HD2	1.98	0.44
1:A:81[B]:LEU:HD21	1:A:99:GLU:OE1	2.19	0.42
1:A:284:TYR:HB3	1:A:303:LEU:HD11	2.01	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:LEU:HB2	1:A:384[A]:THR:HG22	2.03	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	462/497 (93%)	446 (96%)	16 (4%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	406/420 (97%)	397 (98%)	9 (2%)	52 47

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	26	ASP
1	A	32	SER
1	A	73	VAL
1	A	318	GLN

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	319	ILE
1	A	347	SER
1	A	392	GLN
1	A	418	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	348	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	459/497 (92%)	0.31	42 (9%) <b>9</b> <b>10</b>	14, 31, 65, 101	0

All (42) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	30	LEU	6.8
1	A	374	LEU	6.4
1	A	468	GLN	5.6
1	A	28	ILE	5.4
1	A	102	ASP	5.2
1	A	361	PRO	5.1
1	A	390	TYR	5.1
1	A	101	ALA	5.0
1	A	29	THR	4.9
1	A	74	ASN	4.9
1	A	469	PHE	4.7
1	A	75	GLU	4.4
1	A	177	CYS	4.4
1	A	76	GLN	4.4
1	A	31	GLU	4.2
1	A	419	GLY	4.2
1	A	73	VAL	3.8
1	A	418	SER	3.7
1	A	42	LYS	3.7
1	A	77	PRO	3.6
1	A	27	GLU	3.5
1	A	424	ASN	3.4
1	A	346	THR	3.3
1	A	466	PRO	3.3
1	A	138	ALA	3.2
1	A	176	SER	3.1
1	A	3	ASP	2.9

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	41	GLY	2.7
1	A	90	LYS	2.6
1	A	470	PHE	2.6
1	A	391	GLY	2.6
1	A	196	GLU	2.5
1	A	40	ALA	2.5
1	A	250	LYS	2.2
1	A	389	ILE	2.2
1	A	303	LEU	2.2
1	A	277	ASN	2.2
1	A	199	HIS	2.2
1	A	72	GLY	2.2
1	A	43	ASN	2.2
1	A	393	GLY	2.1
1	A	467	GLU	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.