



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2020 – 12:22 pm BST

PDB ID : 5I0H  
Title : Crystal structure of myosin X motor domain in pre-powerstroke state  
Authors : Isabet, T.; Blanc, F.; Sweeney, H.L.; Houdusse, A.  
Deposited on : 2016-02-04  
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	741	
1	B	741	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	B	805	-	-	X	-
5	EDO	B	806	-	-	-	X

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 13273 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Unconventional myosin-X.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	731	Total	C	N	O	S	0	13	0
			5954	3767	1029	1131	27			
1	B	723	Total	C	N	O	S	0	17	0
			5900	3730	1028	1115	27			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total Mg 1 1	0	0
3	A	1	Total Mg 1 1	0	0

- Molecule 4 is BERYLLIUM TRIFLUORIDE ION (three-letter code: BEF) (formula: BeF<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Be F 4 1 3	0	0
4	B	1	Total Be F 4 1 3	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0
5	B	1	Total C O 4 2 2	0	0

- Molecule 6 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0
6	B	1	Total O S 5 4 1	0	0

- Molecule 7 is water.


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
7	A	648	Total 648	O 648	0	0
7	B	610	Total 610	O 610	0	1




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Unconventional myosin-X



- Molecule 1: Unconventional myosin-X



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	76.48Å 78.30Å 78.67Å 75.69° 86.38° 76.12°	Depositor
Resolution (Å)	26.35 – 1.80 47.63 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.5 (26.35-1.80) 97.5 (47.63-1.80)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.55 (at 1.79Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, $R_{free}$	0.165 , 0.190 0.169 , 0.197	Depositor DCC
$R_{free}$ test set	7760 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.0	Xtriage
Anisotropy	0.100	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 49.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.015 for -l,-k,-h	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	13273	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, BEF, EDO, ADP, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.52	0/6076	0.62	1/8212 (0.0%)
1	B	0.50	0/6016	0.61	0/8128
All	All	0.51	0/12092	0.62	1/16340 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	61	ASN	C-N-CA	6.11	136.99	121.70

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5954	0	5724	17	0
1	B	5900	0	5694	24	0
2	A	27	0	12	0	0
2	B	27	0	12	3	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	4	0	0	0	0
4	B	4	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	32	0	48	0	0
5	B	40	0	60	7	0
6	A	5	0	0	0	0
6	B	20	0	0	0	0
7	A	648	0	0	2	0
7	B	610	0	0	3	0
All	All	13273	0	11550	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:801:ADP:HN61	5:B:805:EDO:H22	1.41	0.85
1:B:590:HIS:HD2	7:B:1411:HOH:O	1.63	0.81
1:B:104:ASN:HA	5:B:805:EDO:H11	1.69	0.74
1:A:581:PHE:CD2	1:A:584:ILE:HG12	2.25	0.71
2:B:801:ADP:N6	5:B:805:EDO:H22	2.07	0.69
1:A:693:LEU:HD11	1:A:709:LEU:HD22	1.76	0.68
1:A:248[B]:ASN:HD21	1:A:252:ARG:NH1	1.95	0.65
1:A:122:SER:HB3	1:A:177:ILE:HD11	1.79	0.64
1:B:693:LEU:HD11	1:B:709:LEU:HD22	1.79	0.64
1:B:537:PRO:HG3	1:B:554:GLN:HE22	1.63	0.63
1:A:537:PRO:HG3	1:A:554:GLN:HE22	1.63	0.63
1:B:122:SER:HB3	1:B:177:ILE:HD11	1.82	0.62
2:B:801:ADP:HN61	5:B:805:EDO:C2	2.14	0.58
1:A:84:PHE:CZ	1:A:88:LYS:HD3	2.40	0.56
1:A:297:ASP:HB3	1:A:300:ILE:HD12	1.87	0.56
1:B:84:PHE:CZ	1:B:88:LYS:HD2	2.41	0.55
1:B:244:LEU:H	5:B:809:EDO:H12	1.74	0.52
5:B:805:EDO:H21	7:B:927:HOH:O	2.08	0.52
1:B:435[B]:ILE:HG12	1:B:458[B]:LEU:CD1	2.40	0.52
1:B:58:HIS:O	1:B:59:PRO:C	2.48	0.51
1:B:62:GLU:O	1:B:63:GLU:HB2	2.12	0.50
1:B:32:VAL:HG21	1:B:49:ILE:HG13	1.94	0.48
1:A:325:GLU:HG2	1:A:581:PHE:HE1	1.78	0.48
1:B:14:ARG:HH11	1:B:19:HIS:CD2	2.32	0.48
1:B:328:ARG:NH1	1:B:581:PHE:CD2	2.83	0.47
1:A:465:HIS:HD2	7:A:1492:HOH:O	1.98	0.46
1:A:507:ASN:ND2	1:A:610:THR:OG1	2.49	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:657:LEU:HD22	1:B:662[B]:MET:SD	2.56	0.45
1:A:283:PRO:HB2	1:A:295:VAL:HG22	1.96	0.45
1:B:567[B]:THR:HG22	7:B:1387:HOH:O	2.15	0.45
1:B:186:LEU:O	1:B:190:THR:HG23	2.17	0.45
1:B:68[B]:MET:HE2	1:B:83:LEU:HD21	1.99	0.44
1:A:287:HIS:HD2	7:A:949:HOH:O	1.99	0.44
1:A:154:LEU:CD1	1:A:461[B]:TYR:CE1	3.01	0.44
1:A:498:LYS:HG3	1:A:499:LYS:O	2.17	0.44
1:A:657:LEU:HD22	1:A:662[A]:MET:SD	2.59	0.43
1:B:498:LYS:HG3	1:B:499:LYS:O	2.17	0.43
1:B:18:GLN:NE2	1:B:650[B]:GLN:HE22	2.17	0.43
1:B:244:LEU:H	5:B:809:EDO:C1	2.32	0.43
1:A:248[B]:ASN:HD21	1:A:252:ARG:HH11	1.68	0.41
1:B:635[B]:ARG:HH21	1:B:660:SER:HB2	1.85	0.41
1:A:462:PHE:CE2	1:A:635:ARG:HD3	2.56	0.41
1:B:154:LEU:CD1	1:B:461:TYR:CE1	3.04	0.41
1:B:464:LYS:HE2	1:B:469:LEU:HD11	2.03	0.40
1:B:7:GLU:HA	1:B:8:GLY:HA2	1.64	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	740/741 (100%)	719 (97%)	18 (2%)	3 (0%)	34	21
1	B	734/741 (99%)	708 (96%)	23 (3%)	3 (0%)	34	21
All	All	1474/1482 (100%)	1427 (97%)	41 (3%)	6 (0%)	34	21

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	59	PRO
1	A	381	ARG
1	B	63	GLU
1	B	381	ARG
1	A	62	GLU
1	A	60	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	640/662 (97%)	633 (99%)	7 (1%)	73	68
1	B	634/662 (96%)	630 (99%)	4 (1%)	86	84
All	All	1274/1324 (96%)	1263 (99%)	11 (1%)	78	75

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	GLU
1	A	142	ARG
1	A	295	VAL
1	A	296	GLU
1	A	436	PHE
1	A	659	TYR
1	A	676	ARG
1	B	32	VAL
1	B	43	THR
1	B	436	PHE
1	B	676	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	19	HIS
1	A	40	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	46	GLN
1	A	85	GLN
1	A	120	GLN
1	A	287	HIS
1	A	507	ASN
1	A	554	GLN
1	B	18	GLN
1	B	19	HIS
1	B	40	GLN
1	B	46	GLN
1	B	226	GLN
1	B	248	ASN
1	B	287	HIS
1	B	465	HIS
1	B	554	GLN
1	B	680	GLN
1	B	708	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 2 are monoatomic - leaving 27 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	EDO	A	808	-	3,3,3	0.59	0	2,2,2	0.29	0
6	SO4	B	811	-	4,4,4	0.26	0	6,6,6	0.32	0
5	EDO	A	804	-	3,3,3	0.53	0	2,2,2	0.36	0
5	EDO	B	815	-	3,3,3	0.56	0	2,2,2	0.41	0
6	SO4	B	812	-	4,4,4	0.18	0	6,6,6	0.12	0
5	EDO	A	805	-	3,3,3	0.72	0	2,2,2	0.11	0
2	ADP	A	801	3,4	24,29,29	0.84	0	29,45,45	0.75	1 (3%)
2	ADP	B	801	3,4	24,29,29	0.81	1 (4%)	29,45,45	0.76	1 (3%)
5	EDO	A	806	-	3,3,3	0.56	0	2,2,2	0.46	0
5	EDO	A	811	-	3,3,3	0.63	0	2,2,2	0.18	0
5	EDO	B	808	-	3,3,3	0.44	0	2,2,2	0.62	0
6	SO4	B	813	-	4,4,4	0.13	0	6,6,6	0.13	0
5	EDO	B	816	-	3,3,3	0.46	0	2,2,2	0.25	0
5	EDO	B	810	-	3,3,3	0.78	0	2,2,2	0.11	0
6	SO4	A	809	-	4,4,4	0.40	0	6,6,6	0.25	0
5	EDO	B	806	-	3,3,3	0.58	0	2,2,2	0.36	0
4	BEF	B	803	2	0,3,3	0.00	-	-	-	-
5	EDO	B	804	-	3,3,3	0.51	0	2,2,2	1.05	0
4	BEF	A	803	2	0,3,3	0.00	-	-	-	-
5	EDO	A	807	-	3,3,3	0.56	0	2,2,2	0.34	0
5	EDO	B	809	-	3,3,3	0.61	0	2,2,2	0.64	0
5	EDO	B	807	-	3,3,3	0.62	0	2,2,2	0.05	0
5	EDO	A	810	-	3,3,3	0.50	0	2,2,2	0.45	0
5	EDO	B	805	-	3,3,3	0.44	0	2,2,2	0.32	0
5	EDO	B	817	-	3,3,3	0.43	0	2,2,2	0.48	0
5	EDO	A	812	-	3,3,3	0.45	0	2,2,2	0.59	0
6	SO4	B	814	-	4,4,4	0.15	0	6,6,6	0.06	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	808	-	-	0/1/1/1	-
5	EDO	B	816	-	-	0/1/1/1	-
5	EDO	A	808	-	-	0/1/1/1	-
5	EDO	B	807	-	-	1/1/1/1	-
5	EDO	A	804	-	-	0/1/1/1	-
5	EDO	B	815	-	-	0/1/1/1	-
5	EDO	B	810	-	-	0/1/1/1	-
5	EDO	B	804	-	-	0/1/1/1	-

*Continued on next page...*



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	B	805	-	-	1/1/1/1	-
5	EDO	A	805	-	-	0/1/1/1	-
2	ADP	A	801	3,4	-	2/12/32/32	0/3/3/3
5	EDO	B	817	-	-	0/1/1/1	-
5	EDO	B	809	-	-	1/1/1/1	-
5	EDO	A	812	-	-	0/1/1/1	-
5	EDO	B	806	-	-	0/1/1/1	-
5	EDO	A	806	-	-	0/1/1/1	-
5	EDO	A	807	-	-	1/1/1/1	-
2	ADP	B	801	3,4	-	2/12/32/32	0/3/3/3
5	EDO	A	811	-	-	0/1/1/1	-
5	EDO	A	810	-	-	0/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	ADP	PB-O3B	-2.03	1.47	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	801	ADP	C5-C6-N6	2.13	123.59	120.35
2	A	801	ADP	C5-C6-N6	2.07	123.50	120.35

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	ADP	PA-O3A-PB-O2B
2	B	801	ADP	PA-O3A-PB-O3B
5	A	807	EDO	O1-C1-C2-O2
5	B	805	EDO	O1-C1-C2-O2
2	A	801	ADP	PA-O3A-PB-O3B
2	B	801	ADP	PA-O3A-PB-O2B
5	B	809	EDO	O1-C1-C2-O2
5	B	807	EDO	O1-C1-C2-O2


There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	ADP	3	0
5	B	809	EDO	2	0
5	B	805	EDO	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	731/741 (98%)	-0.23	19 (2%) 56 51	18, 31, 59, 100	1 (0%)
1	B	723/741 (97%)	-0.12	38 (5%) 26 21	20, 33, 73, 156	8 (1%)
All	All	1454/1482 (98%)	-0.18	57 (3%) 39 33	18, 32, 65, 156	9 (0%)

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	49	ILE	11.1
1	B	33	VAL	10.4
1	B	43	THR	5.5
1	A	381	ARG	5.4
1	B	61	ASN	5.4
1	B	295	VAL	5.2
1	B	46	GLN	5.1
1	B	58	HIS	5.0
1	B	598	ASP	4.9
1	B	39	GLY	4.9
1	B	47	SER	4.8
1	A	182	LEU	4.6
1	B	34	PHE	4.5
1	B	599	THR	4.4
1	B	32	VAL	4.4
1	B	48	THR	4.3
1	B	62	GLU	4.2
1	B	41	VAL	4.1
1	A	61	ASN	4.1
1	A	598	ASP	4.1
1	A	512	PHE	3.9
1	A	60	THR	3.7
1	A	52	GLN	3.7
1	B	44	TYR	3.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	51	HIS	3.4
1	A	184	LEU	3.4
1	B	45	LYS	3.3
1	B	512	PHE	3.3
1	A	597	GLN	3.3
1	B	381	ARG	3.2
1	B	42	PHE	3.2
1	B	59	PRO	3.2
1	B	181	SER	3.0
1	A	59	PRO	2.8
1	A	513	PRO	2.8
1	A	382	GLY	2.7
1	B	294	CYS	2.7
1	A	4	PHE	2.6
1	A	380	LEU	2.6
1	B	54	VAL	2.6
1	B	293	GLY	2.6
1	B	25	ASN	2.6
1	B	50	THR	2.5
1	A	378	MET	2.5
1	B	7	GLU	2.5
1	A	63	GLU	2.4
1	B	380	LEU	2.4
1	A	511	HIS	2.4
1	B	60	THR	2.3
1	A	345	GLY	2.3
1	B	24	VAL	2.3
1	B	513	PRO	2.3
1	B	581	PHE	2.2
1	B	385	ILE	2.1
1	A	183	GLU	2.1
1	B	72	THR	2.0
1	B	692	ASN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.


## 6.4 Ligands i

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	A	808	4/4	0.26	0.40	85,86,87,89	0
5	EDO	B	806	4/4	0.27	0.53	93,94,95,95	0
5	EDO	A	811	4/4	0.74	0.28	53,55,57,59	0
6	SO4	B	814	5/5	0.80	0.29	150,150,150,150	0
5	EDO	A	807	4/4	0.80	0.19	67,68,70,70	0
5	EDO	A	806	4/4	0.81	0.33	68,69,70,70	0
5	EDO	A	804	4/4	0.83	0.20	67,67,67,67	0
5	EDO	B	809	4/4	0.83	0.19	35,36,41,41	0
5	EDO	B	807	4/4	0.83	0.30	44,46,49,50	0
6	SO4	B	813	5/5	0.88	0.16	87,88,90,90	0
5	EDO	B	815	4/4	0.89	0.14	39,41,43,48	0
5	EDO	B	816	4/4	0.92	0.36	48,48,48,49	0
5	EDO	A	805	4/4	0.93	0.12	25,31,31,33	0
5	EDO	B	817	4/4	0.93	0.10	46,48,50,54	0
5	EDO	B	804	4/4	0.94	0.16	32,33,36,40	0
5	EDO	B	805	4/4	0.95	0.17	30,32,37,39	0
5	EDO	A	810	4/4	0.95	0.08	33,34,36,41	0
6	SO4	B	811	5/5	0.96	0.06	42,44,49,53	0
5	EDO	B	810	4/4	0.97	0.08	28,36,37,39	0
5	EDO	B	808	4/4	0.97	0.13	28,29,33,34	0
6	SO4	B	812	5/5	0.97	0.17	82,82,83,83	0
2	ADP	B	801	27/27	0.98	0.07	19,24,31,32	0
6	SO4	A	809	5/5	0.98	0.12	48,52,53,56	0
4	BEF	A	803	4/4	0.98	0.12	18,20,20,21	0
5	EDO	A	812	4/4	0.98	0.07	28,29,29,30	0
3	MG	A	802	1/1	0.99	0.10	20,20,20,20	0
2	ADP	A	801	27/27	0.99	0.11	17,18,20,22	0
4	BEF	B	803	4/4	0.99	0.11	19,21,22,22	0
3	MG	B	802	1/1	1.00	0.10	21,21,21,21	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.5 Other polymers [i](#)

There are no such residues in this entry.