



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2023 – 08:38 AM EDT

PDB ID : 2I17
Title : Human aldose reductase in complex with NADP⁺ and the inhibitor IDD594 at temperature of 60K
Authors : Petrova, T.; Ginell, S.; Mitschler, A.; Hasemann, I.; Schneider, T.; Cousido, A.; Lunin, V.Y.; Joachimiak, A.; Podjarny, A.
Deposited on : 2006-08-13
Resolution : 0.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1152 (1.04-0.60)
Ramachandran outliers	138981	1071 (1.04-0.60)
Sidechain outliers	138945	1072 (1.04-0.60)
RSRZ outliers	127900	1044 (1.04-0.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	316	 88% 9% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3724 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aldose reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	313	2965	1908	498	543	16	5	95	1

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	5	ILE	LEU	conflict	UNP P15121

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: $C_{21}H_{30}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	48	21	7	17	3	0	0

- Molecule 3 is IDD594 (three-letter code: LDT) (formula: $C_{16}H_{12}BrF_2NO_3S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
			Total	Br	C	F	N			O	S
3	A	1	24	1	16	2	1	3	1	0	1

- Molecule 4 is CITRIC ACID (three-letter code: CIT) (formula: C₆H₈O₇).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	C O		
4	A	1	13	6 7	0	1
4	A	1	13	6 7	0	1


- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	610	Total 661	O 661	0	182

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aldose reductase

Chain A:  88% 9% ..



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.21Å 66.70Å 47.31Å 90.00° 92.27° 90.00°	Depositor
Resolution (Å)	50.00 – 0.81 49.18 – 0.81	Depositor EDS
% Data completeness (in resolution range)	96.4 (50.00-0.81) 94.5 (49.18-0.81)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.81 (at 0.81Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.076 , 0.091 0.080 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	5.0	Xtrriage
Anisotropy	0.116	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 73.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.004 for -l,k,h 0.021 for -h,-k,l 0.016 for l,-k,h	Xtrriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	3724	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: LDT, NDP, CIT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	2.66	22/3238 (0.7%)	1.74	45/4385 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	27[A]	GLN	CD-NE2	86.16	3.48	1.32
1	A	27[B]	GLN	CD-NE2	86.16	3.48	1.32
1	A	27[A]	GLN	CD-OE1	47.79	2.29	1.24
1	A	27[B]	GLN	CD-OE1	47.79	2.29	1.24
1	A	151	GLU	CD-OE2	-18.75	1.05	1.25
1	A	151	GLU	CD-OE1	10.25	1.36	1.25
1	A	86[A]	LYS	C-N	-9.75	1.15	1.33
1	A	86[B]	LYS	C-N	-9.75	1.15	1.33
1	A	168	GLU	CD-OE1	-8.25	1.16	1.25
1	A	275	LYS	CD-CE	8.21	1.71	1.51
1	A	232[A]	PRO	C-N	8.03	1.52	1.34
1	A	232[B]	PRO	C-N	8.03	1.52	1.34
1	A	168	GLU	CD-OE2	7.73	1.34	1.25
1	A	275	LYS	CE-NZ	-7.20	1.31	1.49
1	A	140	ASP	CG-OD2	-6.85	1.09	1.25
1	A	225	ASP	C-N	-5.93	1.23	1.34
1	A	147[A]	GLU	CD-OE2	-5.85	1.19	1.25
1	A	147[B]	GLU	CD-OE2	-5.85	1.19	1.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	217	ASP	C-N	-5.19	1.22	1.34
1	A	137[A]	ASN	C-N	5.14	1.45	1.34
1	A	137[B]	ASN	C-N	5.14	1.45	1.34
1	A	217	ASP	CB-CG	5.01	1.62	1.51

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	27[A]	GLN	OE1-CD-NE2	-35.97	39.18	121.90
1	A	27[B]	GLN	OE1-CD-NE2	-35.97	39.18	121.90
1	A	27[A]	GLN	CG-CD-NE2	-32.14	39.57	116.70
1	A	27[B]	GLN	CG-CD-NE2	-32.14	39.57	116.70
1	A	27[A]	GLN	CG-CD-OE1	-30.09	61.43	121.60
1	A	27[B]	GLN	CG-CD-OE1	-30.09	61.43	121.60
1	A	217	ASP	CB-CG-OD2	-14.17	105.54	118.30
1	A	277[A]	PHE	CB-CG-CD1	10.21	127.95	120.80
1	A	277[B]	PHE	CB-CG-CD1	10.21	127.95	120.80
1	A	140	ASP	CB-CG-OD2	9.54	126.89	118.30
1	A	135[A]	ASP	CB-CG-OD2	-8.07	111.03	118.30
1	A	135[B]	ASP	CB-CG-OD2	-8.07	111.03	118.30
1	A	308[A]	LYS	CA-CB-CG	7.89	130.75	113.40
1	A	308[B]	LYS	CA-CB-CG	7.89	130.75	113.40
1	A	85[A]	GLU	CA-C-O	-7.25	104.89	120.10
1	A	85[B]	GLU	CA-C-O	-7.25	104.89	120.10
1	A	297[A]	ARG	NE-CZ-NH2	-6.98	116.81	120.30
1	A	297[B]	ARG	NE-CZ-NH2	-6.98	116.81	120.30
1	A	140	ASP	CB-CG-OD1	-6.43	112.52	118.30
1	A	294[A]	ARG	NE-CZ-NH1	-6.41	117.09	120.30
1	A	294[B]	ARG	NE-CZ-NH1	-6.41	117.09	120.30
1	A	99[A]	ASP	CB-CG-OD2	6.35	124.02	118.30
1	A	99[B]	ASP	CB-CG-OD2	6.35	124.02	118.30
1	A	4	ARG	NE-CZ-NH2	-6.13	117.23	120.30
1	A	1[A]	MET	CG-SD-CE	6.03	109.85	100.20
1	A	299[A]	CYS	O-C-N	6.00	132.30	122.70
1	A	299[B]	CYS	O-C-N	6.00	132.30	122.70
1	A	294[A]	ARG	CA-CB-CG	-5.71	100.84	113.40
1	A	294[B]	ARG	CA-CB-CG	-5.71	100.84	113.40
1	A	86[A]	LYS	CA-C-N	5.67	127.54	116.20
1	A	86[B]	LYS	CA-C-N	5.67	127.54	116.20
1	A	147[A]	GLU	OE1-CD-OE2	5.66	130.09	123.30
1	A	147[B]	GLU	OE1-CD-OE2	5.66	130.09	123.30
1	A	116	PHE	CB-CG-CD2	-5.63	116.86	120.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195[A]	LYS	CD-CE-NZ	5.60	124.59	111.70
1	A	195[B]	LYS	CD-CE-NZ	5.60	124.59	111.70
1	A	86[A]	LYS	CA-C-O	-5.51	108.52	120.10
1	A	86[B]	LYS	CA-C-O	-5.51	108.52	120.10
1	A	99[A]	ASP	OD1-CG-OD2	-5.31	113.21	123.30
1	A	99[B]	ASP	OD1-CG-OD2	-5.31	113.21	123.30
1	A	278[A]	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	A	278[B]	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	A	64[A]	ARG	CD-NE-CZ	5.13	130.78	123.60
1	A	190[A]	TYR	CB-CG-CD1	5.01	124.01	121.00
1	A	190[B]	TYR	CB-CG-CD1	5.01	124.01	121.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	210	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2965	0	0	20	0
2	A	48	0	0	0	0
3	A	24	0	0	0	0
4	A	26	0	0	1	0
5	A	661	0	0	11	0
All	All	3724	0	0	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169[B]:MET:SD	5:A:918[B]:HOH:O	2.03	1.15

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103[B]:ASP:O	5:A:933[B]:HOH:O	1.93	0.87
1:A:308[B]:LYS:CD	5:A:905[B]:HOH:O	2.26	0.83
1:A:94[A]:GLN:OE1	5:A:908[A]:HOH:O	1.99	0.79
1:A:275:LYS:CE	5:A:958[A]:HOH:O	2.40	0.68
1:A:306[B]:SER:OG	5:A:593:HOH:O	2.12	0.68
1:A:30[A]:GLU:OE1	5:A:1019:HOH:O	2.16	0.61
1:A:85[B]:GLU:OE2	5:A:653:HOH:O	2.15	0.58
1:A:137[A]:ASN:ND2	5:A:753:HOH:O	2.36	0.57
1:A:99[A]:ASP:OD2	4:A:451[A]:CIT:O5	2.23	0.56
1:A:135[B]:ASP:OD1	1:A:135[B]:ASP:C	2.54	0.47
1:A:195[A]:LYS:NZ	5:A:953[A]:HOH:O	2.50	0.44
1:A:135[B]:ASP:OD1	1:A:135[B]:ASP:O	2.36	0.43
1:A:85[B]:GLU:CG	5:A:832:HOH:O	2.68	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	391/316 (124%)	382 (98%)	9 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	353/281 (126%)	353 (100%)	0	100 100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIT	A	451[A]	-	12,12,12	1.47	5 (41%)	17,17,17	2.10	2 (11%)
4	CIT	A	401[A]	-	12,12,12	2.52	6 (50%)	17,17,17	2.36	4 (23%)
2	NDP	A	319	-	45,52,52	1.22	3 (6%)	53,80,80	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	A	451[A]	-	-	0/16/16/16	-
4	CIT	A	401[A]	-	-	0/16/16/16	-
2	NDP	A	319	-	-	5/30/77/77	0/5/5/5

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	319	NDP	C4N-C3N	-4.67	1.40	1.49
4	A	401[A]	CIT	C3-C6	4.23	1.57	1.53
2	A	319	NDP	C4N-C5N	-3.92	1.38	1.48
4	A	401[A]	CIT	O5-C6	3.65	1.33	1.22
4	A	401[A]	CIT	O3-C5	3.61	1.34	1.22
4	A	401[A]	CIT	O4-C5	-3.32	1.19	1.30
4	A	401[A]	CIT	O1-C1	3.06	1.32	1.22
2	A	319	NDP	C6N-C5N	3.02	1.38	1.33
4	A	401[A]	CIT	C2-C3	2.38	1.56	1.53
4	A	451[A]	CIT	O1-C1	2.36	1.30	1.22
4	A	451[A]	CIT	O5-C6	2.15	1.29	1.22
4	A	451[A]	CIT	O6-C6	-2.09	1.22	1.30
4	A	451[A]	CIT	O2-C1	-2.09	1.23	1.30
4	A	451[A]	CIT	C2-C3	2.05	1.56	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401[A]	CIT	O5-C6-C3	-7.22	112.04	122.25
4	A	451[A]	CIT	O5-C6-C3	-6.12	113.59	122.25
4	A	451[A]	CIT	O6-C6-C3	5.25	122.17	113.05
4	A	401[A]	CIT	O6-C6-C3	3.34	118.85	113.05
4	A	401[A]	CIT	O7-C3-C2	-2.84	102.75	109.40
4	A	401[A]	CIT	C3-C4-C5	2.77	120.52	113.81

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	319	NDP	O4D-C1D-N1N-C2N
2	A	319	NDP	PA-O3-PN-O5D
2	A	319	NDP	C4D-C5D-O5D-PN
2	A	319	NDP	C2B-O2B-P2B-O2X
2	A	319	NDP	PN-O3-PA-O1A

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	451[A]	CIT	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	4

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	86[A]:LYS	C	87:GLY	N	1.15
1	A	135[B]:ASP	C	136[B]:THR	N	1.05
1	A	232[B]:PRO	C	233:ARG	N	1.00
1	A	64:ARG	C	65[B]:GLU	N	0.85

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/316 (99%)	-0.87	1 (0%) 94 81	3, 5, 12, 35	26 (8%)

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	223	PRO	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CIT	A	401[A]	13/13	0.95	0.13	5,7,10,14	13
4	CIT	A	451[A]	13/13	0.97	0.07	3,5,6,8	13
2	NDP	A	319	48/48	1.00	0.03	2,3,5,6	0
3	LDT	A	321[A]	24/24	1.00	0.03	3,3,4,4	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.