



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2024 – 07:57 AM EDT

PDB ID : 4I1Q  
Title : Crystal Structure of hBRAP1 N-BAR domain  
Authors : Sanchez-Barrena, M.J.  
Deposited on : 2012-11-21  
Resolution : 2.53 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5743 (2.54-2.50)
Clashscore	141614	6463 (2.54-2.50)
Ramachandran outliers	138981	6335 (2.54-2.50)
Sidechain outliers	138945	6337 (2.54-2.50)
RSRZ outliers	127900	5630 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	221	 5% 66% 21% 10%
1	B	221	 9% 58% 29% 10%

## 2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3393 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bridging integrator 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	200	1674	1054	288	327	5	0	0	0
1	B	198	1657	1044	285	323	5	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	18	GLY	-	expression tag	UNP Q9UBW5
A	19	SER	-	expression tag	UNP Q9UBW5
A	48	ASN	SER	SEE REMARK 999	UNP Q9UBW5
B	18	GLY	-	expression tag	UNP Q9UBW5
B	19	SER	-	expression tag	UNP Q9UBW5
B	48	ASN	SER	SEE REMARK 999	UNP Q9UBW5

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	38	Total 38	O 38	0	0
2	B	24	Total 24	O 24	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Bridging integrator 2

Chain A: 



- Molecule 1: Bridging integrator 2

Chain B: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	79.15Å 81.28Å 81.10Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	57.41 – 2.53 57.41 – 2.59	Depositor EDS
% Data completeness (in resolution range)	87.6 (57.41-2.53) 87.6 (57.41-2.59)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.97 (at 2.58Å)	Xtrriage
Refinement program	REFMAC 5.7.0024	Depositor
R, $R_{free}$	0.196 , 0.252 0.196 , 0.252	Depositor DCC
$R_{free}$ test set	802 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	49.0	Xtrriage
Anisotropy	0.115	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 27.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.018 for -h,l,k 0.079 for -k,-h,-l 0.036 for l,-k,h 0.016 for l,h,k 0.016 for k,l,h	Xtrriage
Reported twinning fraction	0.888 for H, K, L 0.112 for K, H, -L	Depositor
Outliers	0 of 15848 reflections	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3393	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.75	0/1704	0.85	2/2288 (0.1%)
1	B	0.74	0/1687	0.87	2/2266 (0.1%)
All	All	0.74	0/3391	0.86	4/4554 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	108	LEU	CA-CB-CG	6.02	129.15	115.30
1	A	227	LEU	CB-CG-CD1	-5.88	101.01	111.00
1	A	117	LEU	CB-CG-CD1	-5.32	101.96	111.00
1	B	187	LEU	CA-CB-CG	5.30	127.50	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	160	GLN	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1674	0	1627	38	0
1	B	1657	0	1610	50	0
2	A	38	0	0	2	0
2	B	24	0	0	0	0
All	All	3393	0	3237	83	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (83) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:ALA:O	1:A:175:GLU:HG2	1.74	0.86
1:A:142:ARG:HB2	1:A:187:LEU:HD22	1.64	0.78
1:B:156:LEU:HD21	1:B:174:GLU:HA	1.68	0.75
1:B:156:LEU:HD11	1:B:173:ALA:O	1.88	0.74
1:A:217:ARG:NH1	1:B:228:ASN:OD1	2.22	0.73
1:A:125:MET:HG2	1:A:209:ILE:HD11	1.71	0.73
1:A:75:HIS:ND1	1:A:114:GLU:OE2	2.23	0.71
1:A:118:ALA:O	1:A:123:ARG:HG3	1.91	0.71
1:A:156:LEU:HB2	1:A:177:PHE:HD2	1.54	0.71
1:B:146:LEU:HB2	1:B:184:PHE:CE1	2.27	0.69
1:A:56:GLU:HG2	1:B:88:ILE:HG22	1.75	0.69
1:B:46:SER:O	1:B:49:ASN:HB2	1.92	0.69
1:B:94:ASP:OD2	1:B:237:LYS:NZ	2.27	0.67
1:A:156:LEU:HB2	1:A:177:PHE:CD2	2.31	0.66
1:B:43:PHE:HE1	1:B:192:LEU:HD13	1.61	0.65
1:A:40:ASP:O	1:A:44:GLU:HG2	1.98	0.63
1:A:149:TYR:HE1	1:A:181:GLN:HB3	1.62	0.62
1:A:125:MET:HG2	1:A:209:ILE:CD1	2.28	0.62
1:B:237:LYS:NZ	1:B:237:LYS:HB3	2.15	0.61
1:A:39:LYS:N	1:A:41:GLU:HG3	2.17	0.60
1:A:52:GLN:NE2	1:A:199:TYR:OH	2.35	0.59
1:A:67:PHE:HA	1:B:77:SER:OG	2.04	0.57
1:B:45:GLN:HA	1:B:48:ASN:HB3	1.86	0.56
1:B:181:GLN:O	1:B:185:GLU:HG2	2.07	0.55
1:B:153:ARG:HG2	1:B:177:PHE:HE2	1.71	0.55
1:B:43:PHE:O	1:B:46:SER:N	2.40	0.54
1:A:181:GLN:HG3	1:A:182:THR:N	2.22	0.54

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:50:PHE:CZ	1:B:135:ILE:HG22	2.43	0.54
1:B:187:LEU:O	1:B:191:LEU:HB2	2.08	0.54
1:B:156:LEU:HG	1:B:173:ALA:HB1	1.90	0.53
1:B:136:LYS:O	1:B:139:ILE:HG13	2.08	0.53
2:A:322:HOH:O	1:B:228:ASN:HB3	2.08	0.52
1:A:145:LYS:HD2	1:A:183:VAL:CG1	2.40	0.52
1:A:149:TYR:CE1	1:A:181:GLN:CB	2.93	0.51
1:A:152:ALA:HB1	1:A:177:PHE:CD1	2.45	0.51
1:A:149:TYR:HE1	1:A:181:GLN:CB	2.24	0.51
1:B:142:ARG:NH2	1:B:188:ASN:OD1	2.44	0.50
1:B:186:ASP:O	1:B:190:GLU:HB2	2.11	0.50
1:A:149:TYR:CE1	1:A:181:GLN:HB3	2.43	0.50
1:B:43:PHE:CE1	1:B:192:LEU:HD13	2.43	0.50
1:A:141:LYS:HG2	1:A:144:ARG:NH2	2.26	0.50
1:B:237:LYS:NZ	1:B:237:LYS:CB	2.74	0.50
1:A:146:LEU:HB2	1:A:184:PHE:CE1	2.47	0.50
1:A:64:LEU:HD12	1:A:125:MET:HG3	1.93	0.49
1:B:45:GLN:O	1:B:48:ASN:HB3	2.12	0.49
1:B:50:PHE:CE1	1:B:135:ILE:HG21	2.48	0.48
1:A:82:SER:HB2	1:A:103:VAL:HG22	1.94	0.48
1:A:163:LYS:HD2	1:A:165:LYS:HG2	1.96	0.48
1:B:79:LYS:HD2	1:B:107:ASP:OD1	2.14	0.47
1:A:125:MET:O	1:A:129:VAL:HG22	2.15	0.47
1:A:141:LYS:HG2	1:A:144:ARG:HH21	1.80	0.47
1:A:141:LYS:CG	1:A:144:ARG:NH2	2.77	0.47
1:B:62:LYS:HA	1:B:62:LYS:HD2	1.81	0.47
1:A:44:GLU:OE1	1:A:44:GLU:HA	2.14	0.46
1:B:144:ARG:O	1:B:147:VAL:N	2.48	0.46
1:B:152:ALA:O	1:B:156:LEU:HD13	2.15	0.46
1:B:146:LEU:HB2	1:B:184:PHE:CD1	2.50	0.46
1:B:75:HIS:ND1	1:B:114:GLU:OE2	2.44	0.46
1:B:50:PHE:CZ	1:B:135:ILE:CG2	2.99	0.46
1:A:214:SER:O	1:B:228:ASN:ND2	2.45	0.46
1:B:54:GLN:HE22	1:B:136:LYS:HD2	1.81	0.45
1:B:212:ASN:O	1:B:216:LEU:HG	2.16	0.45
1:B:54:GLN:HA	1:B:132:PHE:CE2	2.52	0.44
1:A:156:LEU:HG	1:A:160:GLN:OE1	2.16	0.44
1:A:51:TYR:O	1:A:54:GLN:HG3	2.18	0.44
1:B:153:ARG:HG2	1:B:177:PHE:CE2	2.51	0.44
1:A:152:ALA:HB1	1:A:177:PHE:CG	2.52	0.43
1:B:234:VAL:HG12	1:B:235:MET:N	2.33	0.43

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:188:ASN:O	1:B:192:LEU:HB2	2.19	0.43
1:B:185:GLU:O	1:B:189:GLN:CB	2.67	0.43
1:A:141:LYS:HD2	1:A:187:LEU:HD11	2.00	0.43
1:B:85:LEU:HD11	1:B:235:MET:CE	2.49	0.43
1:B:203:ILE:O	1:B:207:VAL:HG23	2.18	0.43
1:B:142:ARG:NE	1:B:188:ASN:OD1	2.51	0.42
1:A:126:GLU:OE2	2:A:338:HOH:O	2.21	0.42
1:B:193:GLU:O	1:B:197:ILE:HG13	2.20	0.42
1:B:82:SER:HB2	1:B:103:VAL:HG22	2.01	0.42
1:B:193:GLU:O	1:B:196:PRO:HD2	2.20	0.42
1:B:109:LEU:HB3	1:B:224:MET:HE2	2.01	0.42
1:A:109:LEU:O	1:A:110:TRP:C	2.56	0.41
1:B:56:GLU:OE2	1:B:202:ARG:HD2	2.19	0.41
1:A:56:GLU:CG	1:B:88:ILE:HG22	2.48	0.40
1:B:64:LEU:HD23	1:B:64:LEU:HA	1.89	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	198/221 (90%)	184 (93%)	14 (7%)	0	100	100
1	B	196/221 (89%)	181 (92%)	13 (7%)	2 (1%)	15	27
All	All	394/442 (89%)	365 (93%)	27 (7%)	2 (0%)	29	47

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	162	ALA
1	B	236	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	181/198 (91%)	164 (91%)	17 (9%)	8	16
1	B	179/198 (90%)	165 (92%)	14 (8%)	12	23
All	All	360/396 (91%)	329 (91%)	31 (9%)	10	19

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	39	LYS
1	A	40	ASP
1	A	41	GLU
1	A	52	GLN
1	A	83	GLU
1	A	141	LYS
1	A	142	ARG
1	A	147	VAL
1	A	155	HIS
1	A	157	GLU
1	A	163	LYS
1	A	166	ASP
1	A	179	LYS
1	A	217	ARG
1	A	225	SER
1	A	234	VAL
1	A	238	LEU
1	B	43	PHE
1	B	69	SER
1	B	82	SER
1	B	91	SER
1	B	107	ASP
1	B	111	GLU
1	B	134	GLU
1	B	146	LEU
1	B	151	SER
1	B	153	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	161	ASN
1	B	163	LYS
1	B	175	GLU
1	B	205	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	A	49	ASN
1	A	52	GLN
1	B	49	ASN
1	B	52	GLN
1	B	54	GLN
1	B	155	HIS
1	B	160	GLN
1	B	229	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	200/221 (90%)	0.18	11 (5%) 25 27	30, 47, 125, 156	0
1	B	198/221 (89%)	0.57	20 (10%) 7 6	31, 50, 119, 154	0
All	All	398/442 (90%)	0.37	31 (7%) 13 13	30, 49, 124, 156	0

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	164	LYS	10.5
1	B	162	ALA	8.2
1	A	161	ASN	6.0
1	B	160	GLN	5.9
1	B	169	LYS	5.4
1	A	162	ALA	5.3
1	B	183	VAL	5.3
1	A	39	LYS	5.0
1	B	170	THR	4.9
1	B	163	LYS	4.0
1	B	168	ALA	3.7
1	A	157	GLU	3.7
1	A	168	ALA	3.6
1	B	159	VAL	3.6
1	B	178	ASN	3.6
1	B	179	LYS	3.5
1	A	150	ASP	3.5
1	A	153	ARG	3.4
1	B	161	ASN	3.2
1	B	174	GLU	3.1
1	B	102	ILE	3.0
1	A	172	LYS	3.0
1	A	156	LEU	2.9
1	B	173	ALA	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	174	GLU	2.6
1	B	158	ALA	2.5
1	A	159	VAL	2.5
1	B	153	ARG	2.5
1	B	41	GLU	2.3
1	B	42	ARG	2.2
1	B	50	PHE	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.