



wwPDB X-ray Structure Validation Summary Report ⓘ

May 17, 2020 – 05:29 am BST

PDB ID : 2I2L
Title : X-ray Crystal Structure of Protein yopX from *Bacillus subtilis*. Northeast Structural Genomics Consortium Target SR411.
Authors : Vorobiev, S.M.; Zhou, W.; Seetharaman, J.; Forouhar, F.; Kuzin, A.A.; Ho, C.K.; Janjua, H.; Cunningham, K.; Ma, L.C.; Xiao, R.; Liu, J.; Acton, T.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Northeast Structural Genomics Consortium (NESG)
Deposited on : 2006-08-16
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	142	 2% 52% 36% 8%
1	B	142	 4% 50% 36% 6% 6%
1	C	142	 % 50% 35% 6% 10%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 3102 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YopX protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	130	981	626	162	189	1	3	0	0	0
1	B	133	995	635	165	192	1	2	0	0	0
1	C	128	992	633	163	192	1	3	0	0	0

There are 33 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	MET	MODIFIED RESIDUE	UNP O34401
A	13	MSE	MET	MODIFIED RESIDUE	UNP O34401
A	55	MSE	MET	MODIFIED RESIDUE	UNP O34401
A	135	LEU	-	CLONING ARTIFACT	UNP O34401
A	136	GLU	-	CLONING ARTIFACT	UNP O34401
A	137	HIS	-	CLONING ARTIFACT	UNP O34401
A	138	HIS	-	CLONING ARTIFACT	UNP O34401
A	139	HIS	-	CLONING ARTIFACT	UNP O34401
A	140	HIS	-	CLONING ARTIFACT	UNP O34401
A	141	HIS	-	CLONING ARTIFACT	UNP O34401
A	142	HIS	-	CLONING ARTIFACT	UNP O34401
B	1	MSE	MET	MODIFIED RESIDUE	UNP O34401
B	13	MSE	MET	MODIFIED RESIDUE	UNP O34401
B	55	MSE	MET	MODIFIED RESIDUE	UNP O34401
B	135	LEU	-	CLONING ARTIFACT	UNP O34401
B	136	GLU	-	CLONING ARTIFACT	UNP O34401
B	137	HIS	-	CLONING ARTIFACT	UNP O34401
B	138	HIS	-	CLONING ARTIFACT	UNP O34401
B	139	HIS	-	CLONING ARTIFACT	UNP O34401
B	140	HIS	-	CLONING ARTIFACT	UNP O34401
B	141	HIS	-	CLONING ARTIFACT	UNP O34401
B	142	HIS	-	CLONING ARTIFACT	UNP O34401
C	1	MSE	MET	MODIFIED RESIDUE	UNP O34401

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	13	MSE	MET	MODIFIED RESIDUE	UNP O34401
C	55	MSE	MET	MODIFIED RESIDUE	UNP O34401
C	135	LEU	-	CLONING ARTIFACT	UNP O34401
C	136	GLU	-	CLONING ARTIFACT	UNP O34401
C	137	HIS	-	CLONING ARTIFACT	UNP O34401
C	138	HIS	-	CLONING ARTIFACT	UNP O34401
C	139	HIS	-	CLONING ARTIFACT	UNP O34401
C	140	HIS	-	CLONING ARTIFACT	UNP O34401
C	141	HIS	-	CLONING ARTIFACT	UNP O34401
C	142	HIS	-	CLONING ARTIFACT	UNP O34401

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	41	Total O 41 41	0	0
2	B	48	Total O 48 48	0	0
2	C	45	Total O 45 45	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.52Å 99.52Å 87.43Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.73 – 2.80 28.73 – 2.79	Depositor EDS
% Data completeness (in resolution range)	83.3 (28.73-2.80) 97.5 (28.73-2.79)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 2.80Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.241 , 0.286 0.263 , 0.318	Depositor DCC
R_{free} test set	958 reflections (4.00%)	wwPDB-VP
Wilson B-factor (Å ²)	48.6	Xtrriage
Anisotropy	0.351	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 65.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.54$, $\langle L^2 \rangle = 0.38$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	3102	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.43 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2011e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/999	0.73	2/1360 (0.1%)
1	B	0.45	0/1013	0.83	3/1382 (0.2%)
1	C	0.42	0/1010	0.65	0/1371
All	All	0.43	0/3022	0.74	5/4113 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	135	LEU	CA-CB-CG	14.02	147.55	115.30
1	B	132	GLY	N-CA-C	-6.30	97.35	113.10
1	A	78	LYS	N-CA-C	-5.59	95.91	111.00
1	B	133	VAL	C-N-CA	-5.53	107.87	121.70
1	A	69	SER	N-CA-C	-5.03	97.41	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	124	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	981	0	896	58	0
1	B	995	0	893	66	0
1	C	992	0	933	41	0
2	A	41	0	0	2	0
2	B	48	0	0	1	0
2	C	45	0	0	3	0
All	All	3102	0	2722	156	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 156 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:ARG:HH21	1:A:55:MSE:HE2	1.32	0.94
1:A:70:ILE:HG22	1:A:119:VAL:HA	1.54	0.89
1:B:131:GLU:O	1:B:132:GLY:O	1.94	0.86
1:B:68:ARG:HB2	1:B:121:GLY:O	1.77	0.85
1:B:130:LEU:C	1:B:131:GLU:OE2	2.15	0.84

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	126/142 (89%)	104 (82%)	20 (16%)	2 (2%)	9 31

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	129/142 (91%)	110 (85%)	16 (12%)	3 (2%)	6	21
1	C	124/142 (87%)	106 (86%)	15 (12%)	3 (2%)	6	20
All	All	379/426 (89%)	320 (84%)	51 (14%)	8 (2%)	7	23

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	76	ASP
1	B	132	GLY
1	C	16	TRP
1	C	47	SER
1	B	98	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	97/123 (79%)	89 (92%)	8 (8%)	11	33
1	B	95/123 (77%)	81 (85%)	14 (15%)	3	9
1	C	103/123 (84%)	91 (88%)	12 (12%)	5	16
All	All	295/369 (80%)	261 (88%)	34 (12%)	5	17

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	41	LEU
1	B	68	ARG
1	C	92	VAL
1	B	54	LEU
1	A	128	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	51	ASN
1	B	2	ASN
1	B	12	GLN
1	B	28	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	127/142 (89%)	-0.08	3 (2%) 59 49	23, 44, 70, 76	0
1	B	130/142 (91%)	-0.02	6 (4%) 32 22	14, 45, 70, 85	0
1	C	125/142 (88%)	-0.16	1 (0%) 86 81	23, 42, 64, 73	0
All	All	382/426 (89%)	-0.09	10 (2%) 56 46	14, 44, 69, 85	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	80	SER	5.7
1	B	79	GLU	2.9
1	B	37	TYR	2.7
1	A	79	GLU	2.7
1	A	63	LYS	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.