



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 11:58 pm BST

PDB ID : 5I2C
Title : Arginine-bound CASTOR1 from Homo sapiens
Authors : Saxton, R.A.; Knockenhauer, K.E.; Schwartz, T.U.
Deposited on : 2016-02-08
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	329	 2% 84% 8% 9%
1	B	329	 81% 5% 14%
1	C	329	 2% 81% 6% 13%
1	D	329	 81% 6% 13%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	A	402	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9872 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GATS-like protein 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	301	Total 2340	C 1516	N 382	O 434	S 8	0	0	0
1	B	283	Total 2215	C 1440	N 361	O 406	S 8	0	0	0
1	C	285	Total 2240	C 1458	N 366	O 409	S 7	0	0	0
1	D	287	Total 2217	C 1446	N 356	O 408	S 7	0	1	0

- Molecule 2 is ARGinine (three-letter code: ARG) (formula: C₆H₁₅N₄O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	Total 12	C 6	N 4	O 2	0	0
2	B	1	Total 12	C 6	N 4	O 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			12	6	4	2		
2	D	1	Total	C	N	O	0	0
			12	6	4	2		

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	235	Total	O	0	0
			235	235		
4	B	204	Total	O	0	0
			204	204		
4	C	196	Total	O	0	0
			196	196		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	161	Total 161	O 161	0	0

E209	SER	F222	T226	E244	K247	F301	R322	R323	GLN	GLU	GLY	LEU	ALA	SER
------	-----	------	------	------	------	------	------	------	-----	-----	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	91.39Å 82.60Å 96.67Å 90.00° 116.23° 90.00°	Depositor
Resolution (Å)	86.71 – 1.80 86.71 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.8 (86.71-1.80) 93.0 (86.71-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.72 (at 1.80Å)	Xtrriage
Refinement program	PHENIX (1.10_2155: ???)	Depositor
R, R_{free}	0.172 , 0.204 0.172 , 0.205	Depositor DCC
R_{free} test set	2002 reflections (1.71%)	wwPDB-VP
Wilson B-factor (Å ²)	29.5	Xtrriage
Anisotropy	0.238	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	9872	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2399	0.57	0/3277
1	B	0.37	0/2266	0.55	0/3090
1	C	0.37	0/2294	0.54	0/3129
1	D	0.33	0/2275	0.53	0/3109
All	All	0.37	0/9234	0.55	0/12605

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2340	0	2325	20	0
1	B	2215	0	2224	11	0
1	C	2240	0	2244	15	0
1	D	2217	0	2203	12	0
2	A	12	0	12	0	0
2	B	12	0	12	0	0
2	C	12	0	12	0	0
2	D	12	0	12	0	0
3	A	4	0	3	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	4	0	3	0	0
3	C	4	0	3	0	0
3	D	4	0	3	1	0
4	A	235	0	0	5	0
4	B	204	0	0	1	0
4	C	196	0	0	4	0
4	D	161	0	0	3	0
All	All	9872	0	9056	56	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:38:ARG:NH2	1:C:58:GLU:OE1	2.06	0.88
1:D:244:GLU:HA	1:D:247:LYS:HE2	1.60	0.83
1:B:252:ASP:HB3	1:C:38:ARG:NH2	1.99	0.76
1:A:234:GLU:OE1	1:D:181:GLN:NE2	2.18	0.76
1:C:248:LYS:NZ	4:C:504:HOH:O	2.27	0.67
1:A:183:ARG:HH21	1:A:262:LEU:HD12	1.60	0.66
1:A:99:ARG:N	1:A:99:ARG:HD2	2.11	0.65
1:A:1:MET:SD	4:A:557:HOH:O	2.55	0.64
1:A:115:LEU:HD11	3:A:402:ACT:H1	1.80	0.64
1:C:249:PHE:O	4:C:501:HOH:O	2.16	0.62
1:D:9:ARG:NH1	1:D:127:GLU:OE1	2.23	0.61
1:B:89:VAL:O	4:B:501:HOH:O	2.17	0.58
1:B:20:LEU:HD22	1:B:51:TYR:CZ	2.39	0.58
1:A:318:GLU:OE1	1:A:322:ARG:NH1	2.40	0.55
1:A:44:LEU:HD13	1:A:53:LEU:CD2	2.37	0.55
1:C:225:ILE:N	4:C:509:HOH:O	2.40	0.54
1:A:280:ILE:O	1:A:283:GLN:HG2	2.08	0.53
1:C:38:ARG:CZ	1:C:38:ARG:HB2	2.38	0.53
1:A:183:ARG:NH2	1:A:262:LEU:HD12	2.26	0.51
1:A:187:LEU:HD22	1:A:256:THR:HG22	1.93	0.51
1:D:115:LEU:HD11	3:D:402:ACT:H1	1.93	0.50
1:A:193:THR:HG21	1:A:253:LEU:HB3	1.93	0.50
1:B:252:ASP:HB3	1:C:38:ARG:HH22	1.75	0.50
1:D:25:HIS:HB3	1:D:26:PRO:HD3	1.93	0.50
1:B:2:GLU:HB2	1:B:82:SER:HB2	1.93	0.50
1:C:145:ARG:NH2	4:C:508:HOH:O	2.37	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:114:MET:HE3	4:D:586:HOH:O	2.12	0.49
1:A:1:MET:SD	1:A:2:GLU:HG3	2.53	0.49
1:C:20:LEU:HD22	1:C:51:TYR:CZ	2.49	0.48
3:A:402:ACT:H2	4:A:513:HOH:O	2.13	0.48
1:A:145:ARG:NH2	4:A:504:HOH:O	2.35	0.48
1:A:315:SER:O	1:A:319:VAL:HG13	2.15	0.47
1:A:264:ARG:NH1	4:A:511:HOH:O	2.47	0.46
1:A:21:TRP:CZ3	1:B:64:PRO:HG3	2.50	0.46
1:D:20:LEU:HD22	1:D:51:TYR:CZ	2.51	0.45
1:C:241:MET:HB2	1:C:241:MET:HE2	1.40	0.45
1:C:228:PHE:HE2	1:C:241:MET:HE3	1.81	0.45
1:C:4:HIS:ND1	1:C:146:GLU:OE2	2.38	0.44
1:B:313:ILE:O	1:B:317:ILE:HG12	2.17	0.44
1:B:14:SER:HB2	1:B:301:PHE:CG	2.54	0.43
1:B:134:ILE:HD13	1:B:154:VAL:HG11	1.99	0.43
1:C:280:ILE:O	1:C:283:GLN:HG2	2.18	0.43
1:D:49:GLU:OE2	4:D:502:HOH:O	2.22	0.42
1:B:47:THR:HB	1:B:48:PRO:HD2	2.02	0.42
1:A:262:LEU:N	4:A:520:HOH:O	2.52	0.42
1:A:44:LEU:HD13	1:A:53:LEU:HD23	2.02	0.42
1:B:92:ALA:HB3	1:B:294:SER:HB3	2.00	0.42
1:C:182:ASN:HB2	1:C:184:PHE:CE2	2.55	0.42
1:C:188:THR:OG1	1:C:189:LEU:N	2.52	0.42
1:D:48:PRO:HD3	1:D:226:THR:O	2.20	0.41
1:D:19:GLY:HA3	1:D:68:PHE:CG	2.55	0.41
1:A:64:PRO:HA	1:A:65:PRO:HD3	1.94	0.41
1:A:14:SER:HB2	1:A:301:PHE:CG	2.56	0.41
1:D:14:SER:HB2	1:D:301:PHE:CG	2.56	0.41
1:D:47:THR:HB	1:D:48:PRO:HD2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	293/329 (89%)	291 (99%)	2 (1%)	0	100	100
1	B	275/329 (84%)	271 (98%)	4 (2%)	0	100	100
1	C	275/329 (84%)	272 (99%)	3 (1%)	0	100	100
1	D	280/329 (85%)	278 (99%)	2 (1%)	0	100	100
All	All	1123/1316 (85%)	1112 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/286 (91%)	259 (100%)	1 (0%)	91	89
1	B	245/286 (86%)	244 (100%)	1 (0%)	91	89
1	C	248/286 (87%)	246 (99%)	2 (1%)	81	78
1	D	244/286 (85%)	241 (99%)	3 (1%)	71	65
All	All	997/1144 (87%)	990 (99%)	7 (1%)	84	81

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	99	ARG
1	B	259	SER
1	C	14	SER
1	C	165	ARG
1	D	68	PHE
1	D	208	SER
1	D	322	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	321	GLN
1	C	167	GLN
1	C	181	GLN
1	D	108	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ACT	B	402	-	1,3,3	1.77	0	0,3,3	0.00	-
3	ACT	D	402	-	1,3,3	1.63	0	0,3,3	0.00	-
3	ACT	A	402	-	1,3,3	1.35	0	0,3,3	0.00	-
3	ACT	C	402	-	1,3,3	2.49	1 (100%)	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	402	ACT	CH3-C	2.49	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	402	ACT	1	0
3	A	402	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/329 (91%)	-0.48	2 (0%) 87 86	19, 31, 54, 79	0
1	B	283/329 (86%)	-0.53	0 100 100	23, 35, 60, 83	0
1	C	285/329 (86%)	-0.39	7 (2%) 57 52	23, 40, 64, 88	0
1	D	287/329 (87%)	-0.37	1 (0%) 94 92	25, 42, 76, 90	0
All	All	1156/1316 (87%)	-0.44	10 (0%) 84 82	19, 37, 68, 90	0

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	253	LEU	3.1
1	A	296	TYR	3.0
1	C	38	ARG	2.8
1	C	149	GLY	2.7
1	D	147	VAL	2.4
1	C	252	ASP	2.3
1	A	169	GLY	2.3
1	C	164	PRO	2.1
1	C	135	HIS	2.1
1	C	150	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	C	402	4/4	0.77	0.22	39,54,61,81	0
3	ACT	B	402	4/4	0.90	0.17	54,68,70,70	0
3	ACT	A	402	4/4	0.91	0.21	47,51,57,58	0
3	ACT	D	402	4/4	0.96	0.13	52,60,66,75	0
2	ARG	B	401	12/12	0.98	0.08	23,27,31,31	0
2	ARG	A	401	12/12	0.98	0.08	19,22,24,24	0
2	ARG	C	401	12/12	0.98	0.08	23,27,30,30	0
2	ARG	D	401	12/12	0.99	0.08	26,28,33,33	0

6.5 Other polymers [i](#)

There are no such residues in this entry.