



Full wwPDB NMR Structure Validation Report ⓘ

Oct 11, 2021 – 07:10 PM EDT

PDB ID : 2I38
Title : Solution structure of the RRM of SRp20
Authors : Hargous, Y.F.; Allain, F.H.T.
Deposited on : 2006-08-17

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.23.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	150	

2 Ensemble composition and analysis i

This entry contains 19 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:56 (55)	0.36	16
2	A:75-A:106, A:111-A:143 (65)	0.49	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 3, 6, 8, 10, 12, 16, 19
2	4, 7, 15, 18
3	1, 11, 13
Single-model clusters	5; 9; 14; 17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2302 atoms, of which 1118 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	150	2302	732	1118	216	230	6	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	cloning artifact	UNP P19909
A	2	GLN	-	cloning artifact	UNP P19909
A	57	GLY	-	cloning artifact	UNP P19909
A	58	SER	-	cloning artifact	UNP P19909
A	59	HIS	-	expression tag	UNP P84103
A	60	HIS	-	expression tag	UNP P84103
A	61	HIS	-	expression tag	UNP P84103
A	62	HIS	-	expression tag	UNP P84103
A	63	HIS	-	expression tag	UNP P84103
A	64	HIS	-	expression tag	UNP P84103
A	129	ASP	GLU	engineered mutation	UNP P84103

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.2 Score per residue for model 2

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.3 Score per residue for model 3

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.4 Score per residue for model 4

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.5 Score per residue for model 5

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.6 Score per residue for model 6

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.7 Score per residue for model 7 (medoid)

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3



4.2.8 Score per residue for model 8


- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3



4.2.9 Score per residue for model 9

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3






4.2.10 Score per residue for model 10

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3


Chain A: 64% 15% 20%



4.2.11 Score per residue for model 11

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3


Chain A: 64% 14% 20%



4.2.12 Score per residue for model 12


- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3

Chain A: 63% 15% 20%




4.2.13 Score per residue for model 13

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3




4.2.14 Score per residue for model 14

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3



4.2.15 Score per residue for model 15


- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3



4.2.16 Score per residue for model 16

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3






4.2.17 Score per residue for model 17

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3


Chain A: 65% 13% 20%



4.2.18 Score per residue for model 18

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3


Chain A: 61% 15% 20%



4.2.19 Score per residue for model 19

- Molecule 1: Fusion protein consists of immunoglobulin G-Binding Protein G and Splicing factor, arginine/serine-rich 3

Chain A: 64% 11% 5% 20%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *SANDER module of AMBER 7*.

Of the 50 calculated structures, 19 were deposited, based on the following criterion: *structures with the least restraint violations*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DYANA	structure solution	
Amber 7	refinement	

No chemical shift data was provided.

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.76±0.01	0±0/957 (0.0± 0.0%)	1.26±0.02	6±2/1298 (0.5± 0.1%)
All	All	0.76	0/18183 (0.0%)	1.26	121/24662 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.0±1.9
All	All	0	57

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	121	ARG	NE-CZ-NH1	10.66	125.63	120.30	5	16
1	A	101	ARG	NE-CZ-NH1	9.97	125.29	120.30	16	15
1	A	141	ARG	NE-CZ-NH1	8.60	124.60	120.30	12	16
1	A	92	ARG	NE-CZ-NH1	7.80	124.20	120.30	1	10
1	A	121	ARG	CD-NE-CZ	7.58	134.21	123.60	5	1
1	A	128	ARG	NE-CZ-NH1	7.49	124.04	120.30	18	18
1	A	115	VAL	CA-CB-CG2	7.43	122.05	110.90	5	14
1	A	133	ARG	NE-CZ-NH1	7.34	123.97	120.30	17	9
1	A	121	ARG	NE-CZ-NH2	-6.82	116.89	120.30	5	1
1	A	139	ARG	NE-CZ-NH1	6.09	123.34	120.30	3	10
1	A	101	ARG	NE-CZ-NH2	-5.86	117.37	120.30	8	1
1	A	140	VAL	CA-CB-CG1	5.76	119.54	110.90	11	1
1	A	54	VAL	CA-CB-CG2	5.74	119.51	110.90	3	1
1	A	96	TYR	CB-CG-CD2	-5.59	117.65	121.00	2	4
1	A	86	ASN	C-N-CA	5.27	134.87	121.70	9	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	121	ARG	NH1-CZ-NH2	-5.14	113.75	119.40	17	1
1	A	92	ARG	NE-CZ-NH2	-5.10	117.75	120.30	17	1
1	A	133	ARG	NE-CZ-NH2	5.07	122.83	120.30	9	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	97	TYR	Sidechain	11
1	A	121	ARG	Sidechain	4
1	A	39	VAL	Peptide	4
1	A	139	ARG	Peptide,Sidechain	4
1	A	42	GLU	Peptide	3
1	A	128	ARG	Sidechain	3
1	A	10	LYS	Peptide	3
1	A	92	ARG	Sidechain	2
1	A	75	LYS	Peptide	2
1	A	141	ARG	Peptide,Sidechain	2
1	A	2	GLN	Peptide	2
1	A	129	ASP	Peptide	2
1	A	101	ARG	Peptide,Sidechain	2
1	A	143	GLU	Peptide	2
1	A	24	ALA	Peptide	1
1	A	84	ASN	Peptide	1
1	A	116	GLU	Peptide	1
1	A	140	VAL	Peptide	1
1	A	126	ALA	Peptide	1
1	A	40	ASP	Peptide	1
1	A	83	ASN	Peptide	1
1	A	13	LYS	Peptide	1
1	A	3	TYR	Sidechain	1
1	A	21	VAL	Peptide	1
1	A	117	PHE	Peptide	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	939	897	897	1±1
All	All	17841	17043	17043	16

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:18:THR:HG22	1:A:30:PHE:CZ	0.45	2.46	3	4
1:A:7:LEU:H	1:A:15:GLU:HA	0.44	1.73	16	1
1:A:31:LYS:HE3	1:A:43:TRP:CE2	0.43	2.49	19	4
1:A:43:TRP:CD2	1:A:54:VAL:CG1	0.43	3.02	13	3
1:A:76:VAL:CG1	1:A:115:VAL:HG22	0.43	2.44	5	3
1:A:43:TRP:CD2	1:A:54:VAL:HG13	0.41	2.51	15	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	120/150 (80%)	94±3 (79±2%)	20±3 (17±3%)	6±2 (5±2%)	4	27
All	All	2280/2850 (80%)	1791 (79%)	383 (17%)	106 (5%)	4	27

All 25 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	43	TRP	18
1	A	17	THR	15
1	A	87	LYS	11
1	A	15	GLU	8
1	A	14	GLY	6
1	A	23	ALA	6
1	A	111	GLY	6
1	A	19	GLU	6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	85	GLY	5
1	A	41	GLY	3
1	A	75	LYS	3
1	A	47	ASP	3
1	A	137	GLY	2
1	A	50	LYS	2
1	A	132	GLY	2
1	A	6	ILE	1
1	A	140	VAL	1
1	A	80	ASN	1
1	A	135	LEU	1
1	A	131	ASP	1
1	A	82	GLY	1
1	A	40	ASP	1
1	A	51	THR	1
1	A	136	CYS	1
1	A	86	ASN	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	96/124 (77%)	83±3 (86±3%)	13±3 (14±3%)	6	46
All	All	1824/2356 (77%)	1569 (86%)	255 (14%)	6	46

All 50 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	105	VAL	15
1	A	81	LEU	14
1	A	130	LEU	12
1	A	15	GLU	11
1	A	40	ASP	11
1	A	4	LYS	10
1	A	75	LYS	10
1	A	55	THR	10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	31	LYS	9
1	A	39	VAL	9
1	A	115	VAL	9
1	A	141	ARG	9
1	A	121	ARG	9
1	A	13	LYS	8
1	A	80	ASN	8
1	A	140	VAL	8
1	A	125	ASP	8
1	A	54	VAL	7
1	A	139	ARG	5
1	A	56	GLU	5
1	A	19	GLU	5
1	A	28	LYS	5
1	A	131	ASP	5
1	A	89	GLU	4
1	A	128	ARG	4
1	A	27	GLU	4
1	A	101	ARG	3
1	A	133	ARG	3
1	A	76	VAL	3
1	A	84	ASN	3
1	A	42	GLU	3
1	A	86	ASN	3
1	A	7	LEU	2
1	A	143	GLU	2
1	A	32	GLN	2
1	A	87	LYS	2
1	A	129	ASP	2
1	A	118	GLU	1
1	A	117	PHE	1
1	A	44	THR	1
1	A	92	ARG	1
1	A	22	ASP	1
1	A	12	LEU	1
1	A	138	CYS	1
1	A	142	VAL	1
1	A	10	LYS	1
1	A	78	VAL	1
1	A	18	THR	1
1	A	104	TRP	1
1	A	135	LEU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided