



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 06:01 AM EDT

PDB ID : 1I9L
Title : CARBONIC ANHYDRASE II (F131V) COMPLEXED WITH 4-(AMINOSULFONYL)-N-[(4-FLUOROPHENYL)METHYL]-BENZAMIDE
Authors : Kim, C.-Y.; Chandra, P.P.; Jain, A.; Christianson, D.W.
Deposited on : 2001-03-20
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	259	 85% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	INV	A	555	-	X	-	-
4	INV	A	666	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 2197 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARBONIC ANHYDRASE II.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	258	2054	1317	355	380	2	0	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	131	VAL	PHE	engineered mutation	UNP P00918


- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Hg	0	0
			1	1		

- Molecule 4 is 4-(AMINOSULFONYL)-N-[(4-FLUOROPHENYL)METHYL]-BENZAMIDE (three-letter code: INV) (formula: C₁₄H₁₃FN₂O₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	N	O			S
4	A	1	21	14	1	2	3	1	0	0
4	A	1	21	14	1	2	3	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	99	Total	O	0	0
			99	99		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: CARBONIC ANHYDRASE II

Chain A:  85% 14%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	43.10Å 42.11Å 73.89Å 90.00° 104.91° 90.00°	Depositor
Resolution (Å)	20.00 – 1.93	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-1.93)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR	Depositor
R, R_{free}	0.223 , 0.280	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	2197	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: INV, HG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.59	0/2115	0.73	0/2871

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2054	0	2009	23	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0
4	A	42	0	26	1	0
5	A	99	0	0	3	0
All	All	2197	0	2035	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:HIS:HB2	5:A:265:HOH:O	1.90	0.72
1:A:146:ILE:HG12	1:A:212:LEU:HD12	1.82	0.61
1:A:161:VAL:HG13	1:A:225:LYS:HD2	1.82	0.60
1:A:213:LYS:HD3	1:A:260:PHE:CZ	2.40	0.57
1:A:96:HIS:CD2	1:A:119:HIS:CE1	2.96	0.54
1:A:161:VAL:CG1	1:A:225:LYS:HD2	2.38	0.54
1:A:30:PRO:HG3	1:A:106:GLU:HB3	1.91	0.52
1:A:4:HIS:HA	5:A:328:HOH:O	2.10	0.52
1:A:128:TYR:CE1	1:A:137:GLN:HG3	2.46	0.50
1:A:220:SER:O	1:A:224:LEU:HG	2.11	0.50
1:A:219:SER:OG	1:A:222:GLN:HG3	2.11	0.50
1:A:147:PHE:HB2	1:A:215:PRO:HB3	1.94	0.49
1:A:128:TYR:CZ	1:A:137:GLN:HG3	2.49	0.47
1:A:29:SER:HB3	1:A:30:PRO:HA	1.97	0.47
1:A:247:PRO:HA	5:A:293:HOH:O	2.16	0.46
1:A:6:GLY:O	1:A:12:GLY:HA2	2.15	0.45
1:A:45:LYS:O	1:A:82:GLY:HA2	2.17	0.45
1:A:112:LYS:HE3	1:A:114:TYR:CZ	2.51	0.45
1:A:13:PRO:HA	1:A:16:TRP:CD2	2.54	0.42
1:A:53:GLN:O	1:A:76:LYS:HB3	2.21	0.41
1:A:88:TYR:HB3	1:A:122:HIS:HB3	2.03	0.41
1:A:69:GLU:HB3	4:A:666:INV:H041	2.03	0.41
1:A:14:GLU:CD	1:A:14:GLU:H	2.25	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	256/259 (99%)	244 (95%)	11 (4%)	1 (0%)	34 24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	223/224 (100%)	221 (99%)	2 (1%)	78 75

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	85	ASP
1	A	131	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	17	HIS
1	A	67	ASN
1	A	230	ASN
1	A	255	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	INV	A	666	-	22,22,22	3.96	16 (72%)	31,31,31	2.87	13 (41%)
4	INV	A	555	2	22,22,22	3.20	13 (59%)	31,31,31	3.00	15 (48%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	INV	A	666	-	-	4/15/15/15	0/2/2/2
4	INV	A	555	2	-	2/15/15/15	0/2/2/2

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	666	INV	C17-C18	6.28	1.49	1.37
4	A	666	INV	C17-C16	5.83	1.49	1.38
4	A	666	INV	C07-N09	5.81	1.46	1.33
4	A	666	INV	S11-NP2	5.46	1.71	1.60
4	A	555	INV	C07-N09	5.43	1.45	1.33
4	A	555	INV	C10-C15	-5.35	1.39	1.51
4	A	666	INV	C02-C03	5.31	1.47	1.38
4	A	666	INV	C19-C18	5.26	1.47	1.37
4	A	555	INV	C17-C18	5.16	1.47	1.37
4	A	555	INV	S11-NP2	5.04	1.70	1.60
4	A	666	INV	C20-C19	4.82	1.47	1.38
4	A	666	INV	C20-C15	4.79	1.49	1.38
4	A	666	INV	C03-S11	4.51	1.84	1.77
4	A	666	INV	C10-C15	-4.45	1.41	1.51
4	A	555	INV	C19-C18	4.38	1.45	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	555	INV	C20-C15	4.30	1.48	1.38
4	A	666	INV	C16-C15	4.03	1.47	1.38
4	A	555	INV	C02-C03	3.98	1.45	1.38
4	A	666	INV	O13-S11	3.96	1.51	1.43
4	A	555	INV	C17-C16	3.85	1.45	1.38
4	A	666	INV	O14-S11	3.35	1.50	1.43
4	A	555	INV	C20-C19	3.18	1.44	1.38
4	A	555	INV	C16-C15	3.11	1.45	1.38
4	A	555	INV	O14-S11	3.02	1.49	1.43
4	A	666	INV	C02-C01	2.36	1.43	1.38
4	A	555	INV	C03-S11	2.35	1.80	1.77
4	A	666	INV	C04-C03	2.06	1.42	1.38
4	A	555	INV	C06-C07	2.04	1.54	1.50
4	A	666	INV	O08-C07	2.02	1.27	1.23

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	555	INV	O14-S11-C03	-7.45	99.04	107.35
4	A	666	INV	O13-S11-C03	-6.57	100.02	107.35
4	A	666	INV	O13-S11-NP2	6.37	116.81	107.36
4	A	555	INV	O13-S11-NP2	6.35	116.78	107.36
4	A	555	INV	O14-S11-O13	-6.00	108.90	118.76
4	A	666	INV	O14-S11-O13	-5.65	109.48	118.76
4	A	555	INV	O13-S11-C03	-5.14	101.61	107.35
4	A	555	INV	O14-S11-NP2	5.12	114.95	107.36
4	A	666	INV	C15-C10-N09	4.84	123.43	113.05
4	A	666	INV	C06-C07-N09	4.09	125.87	117.09
4	A	666	INV	O14-S11-NP2	4.03	113.34	107.36
4	A	555	INV	C04-C03-S11	3.79	125.23	119.73
4	A	666	INV	O08-C07-C06	-3.69	114.36	120.94
4	A	555	INV	C03-S11-NP2	3.54	113.39	108.38
4	A	555	INV	C15-C10-N09	3.44	120.42	113.05
4	A	666	INV	C10-N09-C07	-3.42	113.73	121.81
4	A	666	INV	O14-S11-C03	-3.39	103.57	107.35
4	A	555	INV	C06-C07-N09	2.84	123.18	117.09
4	A	666	INV	C03-S11-NP2	2.60	112.06	108.38
4	A	555	INV	C02-C03-S11	-2.52	116.09	119.73
4	A	666	INV	C04-C03-S11	2.45	123.29	119.73
4	A	555	INV	O08-C07-C06	-2.41	116.64	120.94
4	A	666	INV	C19-C18-C17	-2.40	119.63	122.83
4	A	555	INV	C16-C17-C18	2.25	120.69	118.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	666	INV	C16-C17-C18	2.16	120.60	118.36
4	A	555	INV	C20-C15-C16	2.12	121.49	118.17
4	A	555	INV	C19-C18-C17	-2.08	120.06	122.83
4	A	555	INV	C10-N09-C07	-2.00	117.08	121.81

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	666	INV	O08-C07-N09-C10
4	A	666	INV	C06-C07-N09-C10
4	A	555	INV	C04-C03-S11-O14
4	A	555	INV	C02-C03-S11-O14
4	A	666	INV	C02-C03-S11-O14
4	A	666	INV	C04-C03-S11-O14

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	666	INV	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.