



Full wwPDB X-ray Structure Validation Report ⓘ

May 15, 2020 – 09:10 am BST

PDB ID : 6IAZ
Title : The archaeal Methanocaldococcus infernus Elp3 with N-terminus deletion (1-46)
Authors : Lin, T.Y.; Glatt, S.
Deposited on : 2018-11-28
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	492	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 7020 atoms, of which 3426 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Histone acetyltransferase, ELP3 family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	426	6827	2178	3426	574	622	27	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	43	GLY	-	expression tag	UNP D5VRB9
A	44	ALA	-	expression tag	UNP D5VRB9
A	45	MET	-	expression tag	UNP D5VRB9
A	46	ALA	-	expression tag	UNP D5VRB9

- Molecule 2 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	163	Total	O	0	0
			163	163		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone acetyltransferase, ELP3 family



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	63.35Å 63.35Å 250.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.82 – 1.90 34.82 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (34.82-1.90) 99.8 (34.82-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 1.91Å)	Xtrriage
Refinement program	PHENIX (1.13_2998: ???)	Depositor
R, R_{free}	0.195 , 0.217 0.195 , 0.217	Depositor DCC
R_{free} test set	2065 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.5	Xtrriage
Anisotropy	0.111	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.44 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7020	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3468	0.57	2/4669 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	460	ARG	NE-CZ-NH2	-8.12	116.24	120.30
1	A	460	ARG	NE-CZ-NH1	6.51	123.55	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3401	3426	3383	24	0
2	A	30	0	0	1	0
3	A	163	0	0	3	1
All	All	3594	3426	3383	24	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:386:CYS:SG	3:A:768:HOH:O	2.24	0.95
1:A:385:ARG:NH1	3:A:701:HOH:O	2.14	0.80
1:A:342:MET:O	1:A:385:ARG:CZ	2.40	0.69
1:A:137:ARG:NH1	1:A:140:GLN:OE1	2.28	0.66
1:A:135:ARG:NE	1:A:139[A]:GLU:OE2	2.28	0.64
1:A:457:MET:HG2	1:A:503:LYS:HG3	1.80	0.64
1:A:392:VAL:HG21	1:A:480:TRP:NE1	2.17	0.60
1:A:299:ASP:OD2	1:A:460:ARG:NH2	2.30	0.59
1:A:228:ARG:NH1	1:A:300:MET:SD	2.79	0.54
1:A:342:MET:O	1:A:385:ARG:NH2	2.41	0.54
1:A:457:MET:HG3	1:A:497:ALA:CB	2.39	0.52
1:A:321:GLY:HA2	3:A:721:HOH:O	2.11	0.51
1:A:451:GLU:HG3	1:A:529:TYR:CD1	2.47	0.50
1:A:385:ARG:HG2	1:A:385:ARG:HH11	1.80	0.46
1:A:144:VAL:O	1:A:144:VAL:HG12	2.14	0.46
1:A:385:ARG:NE	2:A:604:SO4:O2	2.31	0.46
1:A:160:PHE:HB3	1:A:161:PRO:HD3	1.98	0.46
1:A:268:SER:HB2	1:A:300:MET:HG2	1.97	0.46
1:A:79:ALA:HB1	1:A:141:LEU:HD13	1.98	0.45
1:A:392:VAL:HG21	1:A:480:TRP:CE2	2.51	0.45
1:A:343:PRO:HA	1:A:385:ARG:HH21	1.81	0.45
1:A:342:MET:C	1:A:385:ARG:NH2	2.70	0.44
1:A:343:PRO:HA	1:A:385:ARG:NH2	2.35	0.42
1:A:300:MET:HB2	1:A:347:ARG:O	2.21	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:725:HOH:O	3:A:735:HOH:O 6_454	2.01	0.19

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	423/492 (86%)	412 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/433 (83%)	355 (99%)	3 (1%)	81	82

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
1	A	471	LEU
1	A	474	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	603	-	4,4,4	0.24	0	6,6,6	0.17	0
2	SO4	A	604	-	4,4,4	0.18	0	6,6,6	0.27	0
2	SO4	A	601	-	4,4,4	0.12	0	6,6,6	0.10	0
2	SO4	A	606	-	4,4,4	0.16	0	6,6,6	0.06	0
2	SO4	A	602	-	4,4,4	0.14	0	6,6,6	0.10	0
2	SO4	A	605	-	4,4,4	0.15	0	6,6,6	0.05	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	604	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	426/492 (86%)	0.98	79 (18%) 1 1	26, 41, 83, 136	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	124	LEU	8.9
1	A	471	LEU	8.9
1	A	467	TRP	8.6
1	A	75	ILE	8.4
1	A	476	LYS	7.3
1	A	470	ALA	6.9
1	A	360	ILE	6.6
1	A	87	PRO	6.2
1	A	358	THR	6.0
1	A	354	ASP	5.9
1	A	474	ASP	5.8
1	A	74	THR	5.7
1	A	472	THR	5.6
1	A	357	ALA	5.5
1	A	359	VAL	5.3
1	A	475	ILE	5.0
1	A	311	GLY	5.0
1	A	309	ILE	4.9
1	A	479	SER	4.7
1	A	143	LYS	4.7
1	A	312	THR	4.4
1	A	361	VAL	4.4
1	A	478	VAL	4.1
1	A	355	ILE	4.1
1	A	144	VAL	3.9
1	A	267	VAL	3.8
1	A	351	ILE	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	356	PRO	3.7
1	A	477	GLU	3.7
1	A	300	MET	3.6
1	A	397	TYR	3.6
1	A	345	TRP	3.6
1	A	315	TYR	3.6
1	A	481	GLN	3.5
1	A	229	VAL	3.5
1	A	480	TRP	3.4
1	A	324	LYS	3.4
1	A	145	GLY	3.3
1	A	125	MET	3.3
1	A	482	HIS	3.3
1	A	231	LEU	3.2
1	A	146	HIS	3.1
1	A	301	VAL	3.1
1	A	348	THR	3.1
1	A	347	ARG	3.0
1	A	349	SER	3.0
1	A	141	LEU	3.0
1	A	353	ARG	2.9
1	A	346	VAL	2.8
1	A	466	GLY	2.8
1	A	205	CYS	2.8
1	A	387	ILE	2.8
1	A	298	PRO	2.8
1	A	269	TYR	2.8
1	A	368	ASN	2.7
1	A	299	ASP	2.7
1	A	441	LEU	2.7
1	A	424	PHE	2.6
1	A	385	ARG	2.6
1	A	367	SER	2.5
1	A	147	PRO	2.5
1	A	469	LYS	2.4
1	A	212	TYR	2.4
1	A	140	GLN	2.3
1	A	352	GLN	2.3
1	A	268	SER	2.3
1	A	203	ALA	2.2
1	A	209	ARG	2.2
1	A	228	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	369	LEU	2.1
1	A	440	ARG	2.1
1	A	188	GLU	2.1
1	A	327	ARG	2.1
1	A	461	GLN	2.1
1	A	265	LEU	2.1
1	A	86	SER	2.1
1	A	450	LYS	2.1
1	A	247	GLY	2.1
1	A	316	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	A	605	5/5	0.85	0.31	138,139,142,143	0
2	SO4	A	604	5/5	0.92	0.19	71,71,74,74	0
2	SO4	A	606	5/5	0.95	0.16	126,126,126,130	0
2	SO4	A	602	5/5	0.96	0.12	69,71,72,74	0
2	SO4	A	601	5/5	0.96	0.13	76,76,76,77	0
2	SO4	A	603	5/5	0.98	0.10	39,39,42,44	0

6.5 Other polymers [i](#)

There are no such residues in this entry.