



wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 12:13 am BST

PDB ID : 4IFD
Title : Crystal structure of an 11-subunit eukaryotic exosome complex bound to RNA
Authors : Makino, D.L.; Conti, E.
Deposited on : 2012-12-14
Resolution : 2.81 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION





The reported resolution of this entry is 2.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)
RNA backbone	3102	1227 (3.10-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	304	 85% 13% ..
2	B	248	 83% 14% .
3	C	393	 71% 15% 14%
4	D	245	 79% 12% 9%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	267	<p>%</p> <p>88% 12%</p>
6	F	250	<p>%</p> <p>72% 14% 15%</p>
7	G	242	<p>2%</p> <p>83% 13%</p>
8	H	361	<p>2%</p> <p>70% 12% 18%</p>
9	I	301	<p>5%</p> <p>67% 9% 24%</p>
10	J	1003	<p>4%</p> <p>80% 14% 5%</p>
11	K	179	<p>3%</p> <p>39% 6% 55%</p>
12	R	45	<p>18%</p> <p>42% 22% 33%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	BR	A	402	-	-	X	-
13	BR	J	1101	-	-	X	-
13	BR	J	1102	-	-	X	-

2 Entry composition

There are 18 unique types of molecules in this entry. The entry contains 26848 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exosome complex component RRP45.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	300	2301	1443	394	448	16	0	0	0

- Molecule 2 is a protein called Exosome complex component SKI6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	242	1882	1177	336	361	8	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	EXPRESSION TAG	UNP P46948
B	0	HIS	-	EXPRESSION TAG	UNP P46948

- Molecule 3 is a protein called Exosome complex component RRP43.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	338	2627	1660	453	503	11	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	102	SER	ALA	ENGINEERED MUTATION	UNP P25359
C	363	MET	VAL	ENGINEERED MUTATION	UNP P25359

- Molecule 4 is a protein called Exosome complex component RRP46.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	223	1702	1070	288	334	10	0	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-21	GLY	-	EXPRESSION TAG	UNP P53256
D	-20	HIS	-	EXPRESSION TAG	UNP P53256
D	-19	GLY	-	EXPRESSION TAG	UNP P53256
D	-18	ASN	-	EXPRESSION TAG	UNP P53256
D	-17	ASN	-	EXPRESSION TAG	UNP P53256
D	-16	LYS	-	EXPRESSION TAG	UNP P53256
D	-15	GLU	-	EXPRESSION TAG	UNP P53256
D	-14	PRO	-	EXPRESSION TAG	UNP P53256
D	-13	ASN	-	EXPRESSION TAG	UNP P53256
D	-12	THR	-	EXPRESSION TAG	UNP P53256
D	-11	LYS	-	EXPRESSION TAG	UNP P53256
D	-10	ASN	-	EXPRESSION TAG	UNP P53256
D	-9	ARG	-	EXPRESSION TAG	UNP P53256
D	-8	LEU	-	EXPRESSION TAG	UNP P53256
D	-7	ASP	-	EXPRESSION TAG	UNP P53256
D	-6	SER	-	EXPRESSION TAG	UNP P53256
D	-5	ALA	-	EXPRESSION TAG	UNP P53256
D	-4	GLU	-	EXPRESSION TAG	UNP P53256
D	-3	LYS	-	EXPRESSION TAG	UNP P53256
D	-2	LYS	-	EXPRESSION TAG	UNP P53256
D	-1	LYS	-	EXPRESSION TAG	UNP P53256
D	0	LYS	-	EXPRESSION TAG	UNP P53256

- Molecule 5 is a protein called Exosome complex component RRP42.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	E	267	2042	1304	338	395	5	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	GLY	-	EXPRESSION TAG	UNP Q12277
E	0	HIS	-	EXPRESSION TAG	UNP Q12277
E	138	ILE	VAL	ENGINEERED MUTATION	UNP Q12277

- Molecule 6 is a protein called Exosome complex component MTR3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	F	213	1640	1023	280	327	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	75	SER	THR	ENGINEERED MUTATION	UNP P48240
F	161	THR	MET	ENGINEERED MUTATION	UNP P48240

- Molecule 7 is a protein called Exosome complex component RRP40.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
7	G	236	1804	1150	296	348	10	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
G	-1	GLY	-	EXPRESSION TAG	UNP Q08285
G	0	HIS	-	EXPRESSION TAG	UNP Q08285

- Molecule 8 is a protein called Exosome complex component RRP4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
8	H	295	2248	1400	405	431	12	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	-1	ARG	-	EXPRESSION TAG	UNP P38792
H	0	SER	-	EXPRESSION TAG	UNP P38792

- Molecule 9 is a protein called Exosome complex component CSL4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
9	I	230	1711	1071	303	329	8	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
I	-8	MET	-	EXPRESSION TAG	UNP P53859
I	-7	LYS	-	EXPRESSION TAG	UNP P53859
I	-6	HIS	-	EXPRESSION TAG	UNP P53859
I	-5	HIS	-	EXPRESSION TAG	UNP P53859

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
I	-4	HIS	-	EXPRESSION TAG	UNP P53859
I	-3	HIS	-	EXPRESSION TAG	UNP P53859
I	-2	HIS	-	EXPRESSION TAG	UNP P53859
I	-1	HIS	-	EXPRESSION TAG	UNP P53859
I	0	PRO	-	EXPRESSION TAG	UNP P53859

- Molecule 10 is a protein called Exosome complex exonuclease DIS3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	J	948	7430	4693	1310	1392	35	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
J	-1	GLY	-	EXPRESSION TAG	UNP Q08162
J	0	ALA	-	EXPRESSION TAG	UNP Q08162
J	171	ASN	ASP	ENGINEERED MUTATION	UNP Q08162
J	551	ASN	ASP	ENGINEERED MUTATION	UNP Q08162

- Molecule 11 is a protein called Exosome complex exonuclease RRP6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	K	81	623	393	110	118	2	0	0	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
K	515	ARG	-	EXPRESSION TAG	UNP Q12149
K	516	SER	-	EXPRESSION TAG	UNP Q12149
K	517	MET	-	EXPRESSION TAG	UNP Q12149


- Molecule 12 is a RNA chain called RNA (45-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
12	R	30	540	237	65	208	30	0	0	0

- Molecule 13 is BROMIDE ION (three-letter code: BR) (formula: Br).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
13	J	4	Total Br 4 4	0	0
13	E	1	Total Br 1 1	0	0
13	H	1	Total Br 1 1	0	0
13	I	1	Total Br 1 1	0	0
13	A	2	Total Br 2 2	0	0
13	F	1	Total Br 1 1	0	0


- Molecule 14 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
14	C	1	Total C O 6 3 3	0	0
14	G	1	Total C O 6 3 3	0	0
14	H	1	Total C O 6 3 3	0	0
14	H	1	Total C O 6 3 3	0	0
14	H	1	Total C O 6 3 3	0	0

- Molecule 15 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES)

(formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
15	G	1	12	6	1	4	1	0	0

- Molecule 16 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
16	J	1	1	1	0	0

- Molecule 17 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
17	J	1	1	1	0	0

- Molecule 18 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
18	A	35	35	35	0	0
18	B	35	35	35	0	0
18	C	20	20	20	0	0
18	D	38	38	38	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
18	E	23	Total 23	O 23	0	0
18	F	15	Total 15	O 15	0	0
18	G	19	Total 19	O 19	0	0
18	H	23	Total 23	O 23	0	0
18	I	6	Total 6	O 6	0	0
18	J	27	Total 27	O 27	0	0
18	K	2	Total 2	O 2	0	0
18	R	1	Total 1	O 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Exosome complex component RRP45




- Molecule 2: Exosome complex component SKI6




- Molecule 3: Exosome complex component RRP43




- Molecule 4: Exosome complex component RRP46




- Molecule 5: Exosome complex component RRP42




- Molecule 6: Exosome complex component MTR3




- Molecule 7: Exosome complex component RRP40




- Molecule 8: Exosome complex component RRP4








● Molecule 9: Exosome complex component CSL4




● Molecule 10: Exosome complex exonuclease DIS3



● Molecule 11: Exosome complex exonuclease RRP6





● Molecule 12: RNA (45-MER)



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	154.09Å 107.44Å 150.46Å 90.00° 110.63° 90.00°	Depositor
Resolution (Å)	49.64 – 2.81 49.64 – 2.80	Depositor EDS
% Data completeness (in resolution range)	87.9 (49.64-2.81) 87.9 (49.64-2.80)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8.1_1168)	Depositor
R, R_{free}	0.182 , 0.224 0.187 , 0.226	Depositor DCC
R_{free} test set	4956 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	54.9	Xtrriage
Anisotropy	0.333	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 46.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.020 for l,-k,h	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	26848	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ZN, MES, BR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.21	0/2337	0.35	0/3158
2	B	0.20	0/1906	0.36	0/2570
3	C	0.20	0/2670	0.38	0/3619
4	D	0.20	0/1720	0.39	0/2335
5	E	0.21	0/2082	0.37	0/2834
6	F	0.21	0/1662	0.37	0/2240
7	G	0.21	0/1841	0.37	0/2503
8	H	0.21	0/2282	0.38	0/3086
9	I	0.20	0/1735	0.38	0/2354
10	J	0.20	0/7575	0.37	0/10290
11	K	0.21	0/629	0.37	0/850
12	R	0.15	0/595	0.75	0/919
All	All	0.20	0/27034	0.39	0/36758

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2301	0	2262	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	1882	0	1917	22	0
3	C	2627	0	2668	44	0
4	D	1702	0	1755	20	0
5	E	2042	0	2057	23	0
6	F	1640	0	1610	27	0
7	G	1804	0	1763	18	0
8	H	2248	0	2207	27	0
9	I	1711	0	1676	21	0
10	J	7430	0	7347	93	0
11	K	623	0	626	11	0
12	R	540	0	273	3	0
13	A	2	0	0	3	0
13	E	1	0	0	0	0
13	F	1	0	0	0	0
13	H	1	0	0	0	0
13	I	1	0	0	0	0
13	J	4	0	0	10	0
14	C	6	0	8	0	0
14	G	6	0	8	0	0
14	H	18	0	24	2	0
15	G	12	0	12	1	0
16	J	1	0	0	0	0
17	J	1	0	0	0	0
18	A	35	0	0	0	0
18	B	35	0	0	0	0
18	C	20	0	0	0	0
18	D	38	0	0	1	0
18	E	23	0	0	1	0
18	F	15	0	0	1	0
18	G	19	0	0	0	0
18	H	23	0	0	0	0
18	I	6	0	0	1	0
18	J	27	0	0	0	0
18	K	2	0	0	0	0
18	R	1	0	0	0	0
All	All	26848	0	26213	300	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:J:606:MET:HG2	13:J:1102:BR:BR	2.24	0.93
1:A:73:PHE:HB2	13:A:402:BR:BR	2.30	0.87
10:J:293:PRO:HG2	13:J:1101:BR:BR	2.41	0.75
6:F:44:GLN:HB3	6:F:236:ARG:HH12	1.51	0.74
8:H:143:LEU:HB3	8:H:144:PRO:HD2	1.71	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	298/304 (98%)	288 (97%)	10 (3%)	0	100	100
2	B	240/248 (97%)	235 (98%)	4 (2%)	1 (0%)	34	66
3	C	333/393 (85%)	313 (94%)	20 (6%)	0	100	100
4	D	221/245 (90%)	215 (97%)	5 (2%)	1 (0%)	29	61
5	E	265/267 (99%)	249 (94%)	16 (6%)	0	100	100
6	F	207/250 (83%)	192 (93%)	15 (7%)	0	100	100
7	G	234/242 (97%)	227 (97%)	7 (3%)	0	100	100
8	H	289/361 (80%)	281 (97%)	7 (2%)	1 (0%)	41	72
9	I	222/301 (74%)	216 (97%)	6 (3%)	0	100	100
10	J	942/1003 (94%)	901 (96%)	40 (4%)	1 (0%)	51	81
11	K	77/179 (43%)	71 (92%)	6 (8%)	0	100	100
All	All	3328/3793 (88%)	3188 (96%)	136 (4%)	4 (0%)	51	81

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	167	ASP
4	D	140	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
8	H	121	ASN
10	J	598	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	253/265 (96%)	251 (99%)	2 (1%)	81	94
2	B	210/219 (96%)	209 (100%)	1 (0%)	88	96
3	C	290/349 (83%)	285 (98%)	5 (2%)	60	87
4	D	196/216 (91%)	196 (100%)	0	100	100
5	E	236/241 (98%)	235 (100%)	1 (0%)	91	97
6	F	182/219 (83%)	181 (100%)	1 (0%)	88	96
7	G	198/210 (94%)	195 (98%)	3 (2%)	65	89
8	H	242/313 (77%)	239 (99%)	3 (1%)	71	92
9	I	177/249 (71%)	175 (99%)	2 (1%)	73	92
10	J	812/901 (90%)	801 (99%)	11 (1%)	67	90
11	K	69/163 (42%)	69 (100%)	0	100	100
All	All	2865/3345 (86%)	2836 (99%)	29 (1%)	76	93

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
8	H	119	VAL
9	I	49	ASN
10	J	834	LEU
8	H	126	VAL
9	I	286	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
9	I	49	ASN
11	K	541	GLN
10	J	171	ASN
6	F	55	ASN
10	J	315	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
12	R	28/45 (62%)	9 (32%)	0

5 of 9 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
12	R	-39	A
12	R	-37	A
12	R	-33	G
12	R	-31	U
12	R	-14	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
14	GOL	G	301	-	5,5,5	0.36	0	5,5,5	0.26	0
14	GOL	H	404	-	5,5,5	0.36	0	5,5,5	0.26	0
14	GOL	H	402	-	5,5,5	0.38	0	5,5,5	0.23	0
14	GOL	C	401	-	5,5,5	0.36	0	5,5,5	0.28	0
15	MES	G	302	-	12,12,12	2.26	1 (8%)	14,16,16	2.03	7 (50%)
14	GOL	H	403	-	5,5,5	0.36	0	5,5,5	0.31	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	GOL	G	301	-	-	2/4/4/4	-
14	GOL	H	404	-	-	2/4/4/4	-
14	GOL	H	402	-	-	2/4/4/4	-
14	GOL	C	401	-	-	2/4/4/4	-
15	MES	G	302	-	-	0/6/14/14	0/1/1/1
14	GOL	H	403	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
15	G	302	MES	C8-S	-7.56	1.66	1.77

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
15	G	302	MES	C5-N4-C3	3.93	117.68	108.83
15	G	302	MES	C6-C5-N4	-3.08	105.44	110.10
15	G	302	MES	O1S-S-C8	2.36	109.75	106.92
15	G	302	MES	C7-N4-C5	2.33	117.20	111.23
15	G	302	MES	O2S-S-C8	2.26	109.63	106.92

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
14	G	301	GOL	O1-C1-C2-C3
14	H	402	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
14	H	403	GOL	O1-C1-C2-C3
14	H	404	GOL	O1-C1-C2-C3
14	G	301	GOL	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	H	402	GOL	1	0
15	G	302	MES	1	0
14	H	403	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	300/304 (98%)	-0.34	2 (0%) 87 84	31, 49, 110, 147	0
2	B	242/248 (97%)	-0.53	1 (0%) 92 91	29, 44, 88, 132	0
3	C	338/393 (86%)	-0.12	10 (2%) 50 40	35, 57, 127, 162	0
4	D	223/245 (91%)	-0.39	1 (0%) 92 91	32, 48, 76, 120	0
5	E	267/267 (100%)	-0.36	2 (0%) 87 84	35, 58, 106, 131	0
6	F	213/250 (85%)	-0.32	3 (1%) 75 70	36, 58, 115, 141	0
7	G	236/242 (97%)	-0.29	4 (1%) 70 63	32, 58, 122, 159	0
8	H	295/361 (81%)	-0.26	7 (2%) 59 49	38, 59, 114, 140	0
9	I	230/301 (76%)	0.10	15 (6%) 18 11	51, 81, 128, 140	0
10	J	948/1003 (94%)	-0.04	41 (4%) 35 25	47, 81, 136, 169	0
11	K	81/179 (45%)	0.16	6 (7%) 14 8	48, 90, 131, 143	0
12	R	30/45 (66%)	1.13	8 (26%) 0 0	72, 163, 201, 210	0
All	All	3403/3838 (88%)	-0.18	100 (2%) 51 41	29, 64, 128, 210	0

The worst 5 of 100 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	182	ASP	6.3
10	J	710	SER	6.1
5	E	265	LEU	6.0
9	I	1	MET	5.9
10	J	206	THR	5.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
13	BR	J	1104	1/1	0.73	0.29	137,137,137,137	0
13	BR	A	402	1/1	0.82	0.25	136,136,136,136	0
14	GOL	H	403	6/6	0.87	0.24	72,74,77,81	0
14	GOL	G	301	6/6	0.89	0.19	81,85,91,92	0
16	MG	J	1105	1/1	0.92	0.11	72,72,72,72	0
14	GOL	H	402	6/6	0.92	0.22	66,72,77,77	0
15	MES	G	302	12/12	0.93	0.26	77,107,110,111	0
13	BR	I	301	1/1	0.93	0.07	90,90,90,90	0
13	BR	H	401	1/1	0.94	0.08	75,75,75,75	0
14	GOL	C	401	6/6	0.94	0.17	79,83,84,84	0
13	BR	J	1103	1/1	0.95	0.10	110,110,110,110	0
14	GOL	H	404	6/6	0.95	0.23	58,64,66,67	0
13	BR	A	401	1/1	0.96	0.08	94,94,94,94	0
13	BR	E	301	1/1	0.96	0.09	101,101,101,101	0
13	BR	J	1102	1/1	0.97	0.11	106,106,106,106	0
17	ZN	J	1106	1/1	0.98	0.13	62,62,62,62	0
13	BR	J	1101	1/1	0.98	0.08	76,76,76,76	0
13	BR	F	301	1/1	0.99	0.05	119,119,119,119	0

6.5 Other polymers [i](#)

There are no such residues in this entry.