

# wwPDB X-ray Structure Validation Summary Report (i)

#### Sep 20, 2023 – 12:18 AM EDT

PDB ID : 5IFY

Title: Crystal structure of Glucose-1-phosphate thymidylyltransferase from

Burkholderia vietnamiensis in complex with 2 -Deoxyuridine-5'-monopho

sphate and 2'-Deoxy-Thymidine-B-L-Rhamnose

Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)

Deposited on : 2016-02-26

Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

## 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\ DIFFRACTION$ 

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	305	90%	6%	
1	В	305	91%	5%	•
1	С	305	89%	6% •	5%
1	D	305	91%	5%	•



Continued from previous page...

Mol	Chain	Length	Quality of chain						
1	E	305	91%	5%	-				
1	F	305	91%	5%	<del>.</del>				



# 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 14509 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glucose-1-phosphate thymidylyltransferase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	293	Total	С	Ν	О	S	0	2	0
1	1 11	290	2271	1450	385	427	9	0	2	0 0 0 0
1	В	293	Total	С	N	О	S	0	3	0
1	I D	290	2260	1441	380	430	9	U	3	U
1	С	289	Total	С	N	О	S	0	0	0
1		209	2195	1401	368	418	8	0	0	
1	D	293	Total	С	N	О	S	0	0	0
1	D	290	2151	1376	358	409	8	U	U	0
1	E	293	Total	С	N	Ο	S	0	1	0
1		290	2251	1435	381	426	9	0	1	
1	F	292	Total	С	N	О	S	0	1	0
1	I F	292	2214	1411	369	425	9		1	

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP A4JC15
A	-6	ALA	-	expression tag	UNP A4JC15
A	-5	HIS	-	expression tag	UNP A4JC15
A	-4	HIS	-	expression tag	UNP A4JC15
A	-3	HIS	_	expression tag	UNP A4JC15
A	-2	HIS	-	expression tag	UNP A4JC15
A	-1	HIS	-	expression tag	UNP A4JC15
A	0	HIS	-	expression tag	UNP A4JC15
В	-7	MET	-	expression tag	UNP A4JC15
В	-6	ALA	-	expression tag	UNP A4JC15
В	-5	HIS	-	expression tag	UNP A4JC15
В	-4	HIS	-	expression tag	UNP A4JC15
В	-3	HIS	-	expression tag	UNP A4JC15
В	-2	HIS	-	expression tag	UNP A4JC15
В	-1	HIS	-	expression tag	UNP A4JC15
В	0	HIS	-	expression tag	UNP A4JC15
С	-7	MET	-	expression tag	UNP A4JC15




 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
С	-6	ALA	-	expression tag	UNP A4JC15
С	-5	HIS	-	expression tag	UNP A4JC15
С	-4	HIS	-	expression tag	UNP A4JC15
С	-3	HIS	-	expression tag	UNP A4JC15
С	-2	HIS	-	expression tag	UNP A4JC15
С	-1	HIS	_	expression tag	UNP A4JC15
С	0	HIS	-	expression tag	UNP A4JC15
D	-7	MET	-	expression tag	UNP A4JC15
D	-6	ALA	_	expression tag	UNP A4JC15
D	-5	HIS	_	expression tag	UNP A4JC15
D	-4	HIS	_	expression tag	UNP A4JC15
D	-3	HIS	-	expression tag	UNP A4JC15
D	-2	HIS	-	expression tag	UNP A4JC15
D	-1	HIS	_	expression tag	UNP A4JC15
D	0	HIS	-	expression tag	UNP A4JC15
Е	-7	MET	_	expression tag	UNP A4JC15
Е	-6	ALA	-	expression tag	UNP A4JC15
E	-5	HIS	_	expression tag	UNP A4JC15
Е	-4	HIS	_	expression tag	UNP A4JC15
E	-3	HIS	_	expression tag	UNP A4JC15
Е	-2	HIS	_	expression tag	UNP A4JC15
Е	-1	HIS	-	expression tag	UNP A4JC15
Е	0	HIS	-	expression tag	UNP A4JC15
F	-7	MET	-	expression tag	UNP A4JC15
F	-6	ALA	-	expression tag	UNP A4JC15
F	-5	HIS	-	expression tag	UNP A4JC15
F	-4	HIS	-	expression tag	UNP A4JC15
F	-3	HIS	-	expression tag	UNP A4JC15
F	-2	HIS	-	expression tag	UNP A4JC15
F	-1	HIS	-	expression tag	UNP A4JC15
F	0	HIS	-	expression tag	UNP A4JC15

 $\bullet$  Molecule 2 is 2'-DEOXY-THYMIDINE-BETA-L-RHAMNOSE (three-letter code: TRH) (formula:  $\rm C_{16}H_{26}N_2O_{15}P_2).$ 






Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
2	A	1	Total	С	N	О	Р	0	0	
2	A	1	35	16	2	15	2	0	0	
2	В	1	Total	С	N	О	Р	0	0	
2	Б	1	35	16	2	15	2	0	U	
2	C	1	Total	С	N	О	Р	0	0	
2		1	35	16	2	15	2	U		
2	D	1	Total	С	N	О	Р	0	0	
2	D	1	35	16	2	15	2	U	0	
2	Е	1	Total	С	N	О	Р	0	0	
	Ľ	1	35	16	2	15	2	U	U	
2	F	1	Total	С	N	О	Р	0	0	
	I'	1	35	16	2	15	2	0	U	

 $\bullet$  Molecule 3 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula:  $\rm C_9H_{13}N_2O_8P).$ 





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf		
3	A	1	Total	С	N	О	Р	0	0		
3	Λ	1	20	9	2	8	1	0	U		
3	В	1	Total	С	N	Ο	Р	0	0		
	Ъ	Б	1	20	9	2	8	1	U	0	
3	С	1	Total	С	N	Ο	Р	0	0		
		1	20	9	2	8	1	0	U		
3	D	1	Total	С	N	Ο	Р	0	0		
	D	1	20	9	2	8	1	U	U		
3	E	1	Total	С	N	Ο	Р	0	0		
	Ľ	1	20	9	2	8	1	U	U		
3	F	1	Total	С	N	О	Р	0	0		
3	I'	1	20	9	2	8	1				

#### • Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	185	Total O 185 185	0	0
4	В	200	Total O 200 200	0	0
4	С	101	Total O 101 101	0	0
4	D	74	Total O 74 74	0	0
4	E	162	Total O 162 162	0	0
4	F	115	Total O 115 115	0	0




## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glucose-1-phosphate thymidylyltransferase







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	109.60Å 254.72Å 82.05Å	Donositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.20 - 2.25	Depositor
rtesolution (A)	46.20 - 2.25	Depositor Depositor EDS Depositor EDS Depositor Depositor Atriage Depositor Depositor Depositor Depositor Atriage EDS Atriage EDS Atriage Atriage EDS Atriage EDS Matriage EDS Matriage EDS Matriage EDS Matriage EDS Matriage EDS Matriage EDS
% Data completeness	99.5 (46.20-2.25)	Depositor
(in resolution range)	99.5 (46.20-2.25)	EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.01 (at 2.24Å)	Xtriage
Refinement program	PHENIX dev_2328	Depositor
D D.	0.157 , 0.197	Depositor
$R, R_{free}$	0.157 , 0.198	DCC
$R_{free}$ test set	1985 reflections (1.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.9	Xtriage
Anisotropy	0.559	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.33 , 51.7	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.48, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	14509	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.59% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: UMP, TRH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.55	0/2329	0.65	$2/3170 \ (0.1\%)$	
1	В	0.54	0/2318	0.66	0/3155	
1	С	0.43	0/2244	0.59	0/3059	
1	D	0.38	0/2201	0.56	0/3012	
1	Е	0.47	0/2304	0.60	0/3137	
1	F	0.43	0/2266	0.58	0/3089	
All	All	0.47	0/13662	0.61	$2/18622 \ (0.0\%)$	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
1	A	154	PHE	CB-CG-CD1	5.35	124.54	120.80
1	A	154	PHE	CB-CG-CD2	-5.30	117.09	120.80

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	$\mathbf{H}(\mathbf{model})$	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	A	2271	0	2217	15	0
1	В	2260	0	2206	11	0
1	С	2195	0	2112	17	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2151	0	2005	11	0
1	Ε	2251	0	2189	8	0
1	F	2214	0	2122	9	0
2	A	35	0	24	0	0
2	В	35	0	24	0	0
2	С	35	0	24	0	0
2	D	35	0	24	1	0
2	Е	35	0	24	0	0
2	F	35	0	24	0	0
3	A	20	0	11	0	0
3	В	20	0	11	0	0
3	С	20	0	11	1	0
3	D	20	0	11	2	0
3	Ε	20	0	11	0	0
3	F	20	0	11	1	0
4	A	185	0	0	0	0
4	В	200	0	0	3	0
4	С	101	0	0	3	0
4	D	74	0	0	2	0
4	Ε	162	0	0	0	1
4	F	115	0	0	1	0
All	All	14509	0	13061	64	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 64 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:121:GLN:HG2	1:C:124:ARG:NH2	1.93	0.84
1:B:128:LYS:NZ	4:B:601:HOH:O	2.12	0.80
1:B:259:ARG:HD2	4:B:746:HOH:O	1.82	0.80
1:D:82:GLN:HE21	1:D:90:GLN:HE21	1.30	0.78
1:A:141:ASP:OD1	1:A:144:ARG:NH1	2.17	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1			$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
4:E:680:HOH:O	4:E:680:HOH:O[2_555]	2.09	0.11



#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	$293/305\ (96\%)$	290 (99%)	2 (1%)	1 (0%)	41	46
1	В	$294/305\ (96\%)$	290 (99%)	3 (1%)	1 (0%)	41	46
1	С	287/305~(94%)	282 (98%)	4 (1%)	1 (0%)	41	46
1	D	291/305~(95%)	284 (98%)	5 (2%)	2 (1%)	22	21
1	E	$292/305\ (96\%)$	289 (99%)	2 (1%)	1 (0%)	41	46
1	F	291/305~(95%)	286 (98%)	4 (1%)	1 (0%)	41	46
All	All	1748/1830 (96%)	1721 (98%)	20 (1%)	7 (0%)	34	37

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	195	GLY
1	A	31	TYR
1	В	31	TYR
1	С	31	TYR
1	D	31	TYR

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	232/249~(93%)	228 (98%)	4 (2%)	60	71	
1	В	231/249~(93%)	228 (99%)	3 (1%)	69	79	
1	С	$220/249\ (88\%)$	216 (98%)	4 (2%)	59	68	



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percei	ntiles
1	D	204/249 (82%)	203 (100%)	1 (0%)	88	92
1	E	229/249 (92%)	228 (100%)	1 (0%)	91	94
1	F	222/249 (89%)	220 (99%)	2 (1%)	78	86
All	All	1338/1494 (90%)	1323 (99%)	15 (1%)	73	82

5 of 15 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	66	MET
1	F	111	ASN
1	С	117	ASP
1	F	117	ASP
1	D	45	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	Ε	90	GLN
1	Ε	290	HIS
1	F	282	GLN
1	С	21	HIS
1	D	21	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



## 5.6 Ligand geometry (i)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trmo	Chain	Res	Link	Во	ond leng	ths	В	ond ang	les
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	UMP	Е	501	-	21,21,21	0.75	1 (4%)	31,31,31	0.81	1 (3%)
3	UMP	A	501	-	21,21,21	0.46	0	31,31,31	0.40	0
2	TRH	Е	500	-	34,37,37	0.52	1 (2%)	52,57,57	1.27	2 (3%)
2	TRH	D	500	-	34,37,37	0.34	0	52,57,57	1.22	2 (3%)
2	TRH	A	500	-	34,37,37	0.37	0	52,57,57	1.41	2 (3%)
2	TRH	F	500	-	34,37,37	0.27	0	52,57,57	1.65	2 (3%)
3	UMP	В	501	-	21,21,21	0.49	0	31,31,31	0.59	1 (3%)
3	UMP	F	501	-	21,21,21	0.47	0	31,31,31	0.43	0
3	UMP	С	501	-	21,21,21	0.67	1 (4%)	31,31,31	0.41	0
3	UMP	D	501	-	21,21,21	0.58	0	31,31,31	0.48	0
2	TRH	С	500	-	34,37,37	0.36	0	52,57,57	1.33	2 (3%)
2	TRH	В	500	-	34,37,37	0.45	1 (2%)	52,57,57	1.06	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UMP	Е	501	-	-	2/10/22/22	0/2/2/2
3	UMP	A	501	-	-	2/10/22/22	0/2/2/2
2	TRH	Е	500	-	-	2/21/53/53	0/3/3/3
2	TRH	D	500	-	-	3/21/53/53	0/3/3/3
2	TRH	A	500	-	-	3/21/53/53	0/3/3/3
2	TRH	F	500	-	-	4/21/53/53	0/3/3/3
3	UMP	В	501	-	-	2/10/22/22	0/2/2/2
3	UMP	F	501	-	-	2/10/22/22	0/2/2/2



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	UMP	С	501	-	-	2/10/22/22	0/2/2/2
3	UMP	D	501	-	-	2/10/22/22	0/2/2/2
2	TRH	С	500	-	-	4/21/53/53	0/3/3/3
2	TRH	В	500	-	-	4/21/53/53	0/3/3/3

#### All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(A)
3	С	501	UMP	P-OP1	2.70	1.59	1.50
3	E	501	UMP	P-OP1	2.58	1.58	1.50
2	Е	500	TRH	P2-O1	2.43	1.66	1.60
2	В	500	TRH	P2-O1	2.09	1.66	1.60

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	F	500	TRH	O5-C1-O1	-10.29	97.91	111.36
2	A	500	TRH	O5-C1-O1	-9.52	98.92	111.36
2	С	500	TRH	O5-C1-O1	-8.63	100.08	111.36
2	Е	500	TRH	O5-C1-O1	-8.20	100.65	111.36
2	D	500	TRH	O5-C1-O1	-7.28	101.85	111.36

There are no chirality outliers.

5 of 32 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	TRH	O5-C1-O1-P2
2	В	500	TRH	O5-C1-O1-P2
2	С	500	TRH	O5-C1-O1-P2
2	D	500	TRH	C1-O1-P2-O4P
2	D	500	TRH	O5-C1-O1-P2


There are no ring outliers.


4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	500	TRH	1	0
3	F	501	UMP	1	0
3	С	501	UMP	1	0
3	D	501	UMP	2	0





The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.








































# 5.7 Other polymers (i)

There are no such residues in this entry.



# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	293/305~(96%)	-0.22	6 (2%) 65 68	19, 31, 61, 104	0
1	В	293/305 (96%)	-0.32	3 (1%) 82 84	17, 28, 51, 102	0
1	С	289/305 (94%)	-0.01	6 (2%) 63 66	25, 45, 74, 95	0
1	D	293/305 (96%)	0.36	26 (8%) 9 10	24, 55, 92, 119	0
1	E	293/305 (96%)	-0.19	6 (2%) 65 68	22, 36, 65, 99	0
1	F	292/305~(95%)	0.05	10 (3%) 45 47	22, 44, 76, 104	0
All	All	1753/1830 (95%)	-0.05	57 (3%) 46 48	17, 38, 78, 119	0

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	150	PHE	7.1
1	С	290	HIS	6.1
1	В	291	GLN	4.1
1	F	188	ASP	4.0
1	D	293	ALA	3.8

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	TRH	D	500	35/35	0.93	0.15	58,66,80,84	0
3	UMP	С	501	20/20	0.95	0.13	34,47,56,63	0
3	UMP	D	501	20/20	0.96	0.10	24,37,50,52	0
2	TRH	С	500	35/35	0.97	0.09	25,38,44,46	0
2	TRH	F	500	35/35	0.97	0.12	27,44,56,58	0
3	UMP	Е	501	20/20	0.97	0.09	19,30,45,49	0
3	UMP	F	501	20/20	0.97	0.11	27,34,45,47	0
2	TRH	В	500	35/35	0.98	0.08	20,26,29,31	0
2	TRH	A	500	35/35	0.98	0.10	23,32,40,42	0
3	UMP	A	501	20/20	0.98	0.12	15,26,36,37	0
3	UMP	В	501	20/20	0.98	0.11	16,25,36,38	0
2	TRH	Е	500	35/35	0.99	0.10	19,27,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


















# Electron density around TRH C 500: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around TRH F 500: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $mF_o$ -DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)













# Electron density around TRH B 500: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $mF_o$ -DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive) Electron density around TRH A 500: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $mF_o$ -DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)














# 6.5 Other polymers (i)

There are no such residues in this entry.

