



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 26, 2020 – 12:11 pm BST

PDB ID : 5IWW  
Title : Crystal structure of RNA editing factor of designer PLS-type PPR/9R protein in complex with MORF9/RIP9  
Authors : Yan, J.; Zhang, Q.; Guan, Z.; Zou, T.; Yin, P.  
Deposited on : 2016-03-23  
Resolution : 2.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1332 (2.68-2.64)
Clashscore	141614	1374 (2.68-2.64)
Ramachandran outliers	138981	1349 (2.68-2.64)
Sidechain outliers	138945	1349 (2.68-2.64)
RSRZ outliers	127900	1318 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	133	
1	B	133	
1	C	133	
2	D	347	

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4987 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Multiple organellar RNA editing factor 9, chloroplastic.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	110	871	559	132	174	6	0	0	0
1	B	110	871	559	132	174	6	0	0	0
1	C	108	857	551	130	170	6	0	0	0

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	74	MET	-	initiating methionine	UNP Q9LPZ1
A	85	SER	CYS	engineered mutation	UNP Q9LPZ1
A	187	SER	CYS	engineered mutation	UNP Q9LPZ1
A	197	LEU	-	expression tag	UNP Q9LPZ1
A	198	GLU	-	expression tag	UNP Q9LPZ1
A	199	HIS	-	expression tag	UNP Q9LPZ1
A	200	HIS	-	expression tag	UNP Q9LPZ1
A	201	HIS	-	expression tag	UNP Q9LPZ1
A	202	HIS	-	expression tag	UNP Q9LPZ1
A	203	HIS	-	expression tag	UNP Q9LPZ1
A	204	HIS	-	expression tag	UNP Q9LPZ1
A	205	HIS	-	expression tag	UNP Q9LPZ1
A	206	HIS	-	expression tag	UNP Q9LPZ1
B	74	MET	-	initiating methionine	UNP Q9LPZ1
B	85	SER	CYS	engineered mutation	UNP Q9LPZ1
B	187	SER	CYS	engineered mutation	UNP Q9LPZ1
B	197	LEU	-	expression tag	UNP Q9LPZ1
B	198	GLU	-	expression tag	UNP Q9LPZ1
B	199	HIS	-	expression tag	UNP Q9LPZ1
B	200	HIS	-	expression tag	UNP Q9LPZ1
B	201	HIS	-	expression tag	UNP Q9LPZ1
B	202	HIS	-	expression tag	UNP Q9LPZ1
B	203	HIS	-	expression tag	UNP Q9LPZ1

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	204	HIS	-	expression tag	UNP Q9LPZ1
B	205	HIS	-	expression tag	UNP Q9LPZ1
B	206	HIS	-	expression tag	UNP Q9LPZ1
C	74	MET	-	initiating methionine	UNP Q9LPZ1
C	85	SER	CYS	engineered mutation	UNP Q9LPZ1
C	187	SER	CYS	engineered mutation	UNP Q9LPZ1
C	197	LEU	-	expression tag	UNP Q9LPZ1
C	198	GLU	-	expression tag	UNP Q9LPZ1
C	199	HIS	-	expression tag	UNP Q9LPZ1
C	200	HIS	-	expression tag	UNP Q9LPZ1
C	201	HIS	-	expression tag	UNP Q9LPZ1
C	202	HIS	-	expression tag	UNP Q9LPZ1
C	203	HIS	-	expression tag	UNP Q9LPZ1
C	204	HIS	-	expression tag	UNP Q9LPZ1
C	205	HIS	-	expression tag	UNP Q9LPZ1
C	206	HIS	-	expression tag	UNP Q9LPZ1

- Molecule 2 is a protein called PLS9-PPR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	300	2332	1501	368	446	17	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total 18	O 18	0	0
3	B	9	Total 9	O 9	0	0
3	C	8	Total 8	O 8	0	0
3	D	21	Total 21	O 21	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Multiple organellar RNA editing factor 9, chloroplastic

Chain A: 




- Molecule 1: Multiple organellar RNA editing factor 9, chloroplastic

Chain B: 





- Molecule 1: Multiple organellar RNA editing factor 9, chloroplastic

Chain C: 



- Molecule 2: PLS9-PPR

Chain D: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	103.19Å 103.19Å 149.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	51.59 – 2.65 76.66 – 2.65	Depositor EDS
% Data completeness (in resolution range)	98.3 (51.59-2.65) 95.4 (76.66-2.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.54 (at 2.65Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.180 , 0.221 0.180 , 0.221	Depositor DCC
$R_{free}$ test set	1250 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.7	Xtrriage
Anisotropy	0.323	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 39.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.056 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	4987	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	61.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.19% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.39	0/893	0.57	0/1214
1	B	0.39	0/893	0.58	0/1214
1	C	0.43	0/879	0.58	0/1194
2	D	0.43	0/2367	0.59	0/3178
All	All	0.42	0/5032	0.58	0/6800

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	871	0	842	7	0
1	B	871	0	842	9	0
1	C	857	0	828	11	0
2	D	2332	0	2379	43	0
3	A	18	0	0	2	0
3	B	9	0	0	2	0
3	C	8	0	0	0	0
3	D	21	0	0	1	0
All	All	4987	0	4891	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 66 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:188:THR:O	3:A:301:HOH:O	1.83	0.97
1:B:85:SER:HA	3:B:301:HOH:O	1.62	0.96
2:D:265:ALA:HB2	2:D:298:MET:HA	1.49	0.92
1:B:164:ASP:O	3:B:301:HOH:O	2.01	0.79
1:C:171:ASN:ND2	1:C:171:ASN:O	2.26	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	108/133 (81%)	106 (98%)	2 (2%)	0	100	100
1	B	108/133 (81%)	104 (96%)	4 (4%)	0	100	100
1	C	106/133 (80%)	100 (94%)	5 (5%)	1 (1%)	17	26
2	D	298/347 (86%)	286 (96%)	9 (3%)	3 (1%)	15	23
All	All	620/746 (83%)	596 (96%)	20 (3%)	4 (1%)	25	37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	181	ASN
2	D	304	LEU
2	D	313	ASP
2	D	292	TYR



### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	98/121 (81%)	91 (93%)	7 (7%)	14	22
1	B	98/121 (81%)	91 (93%)	7 (7%)	14	22
1	C	96/121 (79%)	88 (92%)	8 (8%)	11	16
2	D	254/297 (86%)	234 (92%)	20 (8%)	12	19
All	All	546/660 (83%)	504 (92%)	42 (8%)	13	20

5 of 42 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	171	ASN
2	D	32	LYS
2	D	283	SER
1	C	173	ASP
1	C	181	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	110/133 (82%)	-0.10	0 <a href="#">100</a>   <a href="#">100</a>	41, 51, 76, 100	0
1	B	110/133 (82%)	-0.07	1 (0%) <a href="#">84</a>   <a href="#">83</a>	45, 57, 94, 115	0
1	C	108/133 (81%)	-0.16	0 <a href="#">100</a>   <a href="#">100</a>	46, 62, 102, 122	0
2	D	300/347 (86%)	-0.08	0 <a href="#">100</a>   <a href="#">100</a>	41, 57, 87, 110	0
All	All	628/746 (84%)	-0.10	1 (0%) <a href="#">95</a>   <a href="#">96</a>	41, 57, 94, 122	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	167	ILE	2.3

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.