



# wwPDB EM Validation Summary Report ⓘ

Feb 13, 2024 – 04:29 PM EST

PDB ID : 3J16  
EMDB ID : EMD-2010  
Title : Models of ribosome-bound Dom34p and Rli1p and their ribosomal binding partners  
Authors : Becker, T.; Franckenberg, S.; Wickles, S.; Shoemaker, C.J.; Anger, A.M.; Armache, J.-P.; Sieber, H.; Ungewickell, C.; Berninghausen, O.; Daberkow, I.; Karcher, A.; Thomm, M.; Hopfner, K.-P.; Green, R.; Beckmann, R.  
Deposited on : 2011-12-12  
Resolution : 7.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev70  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.9  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 7.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	386	
2	B	608	
3	J	233	
4	K	155	
5	L	75	
6	F	191	
7	E	63	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	G	312	
9	C	236	
10	H	165	
11	I	137	
12	D	135	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	MG	B	701	-	-	X	-
14	ATP	B	702	-	-	X	-
15	SF4	B	703	-	-	X	-
15	SF4	B	704	-	-	X	-

## 2 Entry composition

There are 16 unique types of molecules in this entry. The entry contains 26208 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dom34p.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	386	3097	1996	483	603	15	0	0

- Molecule 2 is a protein called Rli1p.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	608	4804	3065	831	884	24	0	0

- Molecule 3 is a RNA chain called 28S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
3	J	233	4942	2222	899	1598	223	0	0

- Molecule 4 is a RNA chain called 18S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
4	K	155	3286	1476	591	1069	150	0	0

- Molecule 5 is a RNA chain called P-site tRNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
5	L	75	1595	712	280	529	74	0	0

- Molecule 6 is a protein called 60S ribosomal protein L6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	F	191	1519	963	274	278	4	0	0

- Molecule 7 is a protein called 40S ribosomal protein S30E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	E	55	440	277	90	72	1	0	0

- Molecule 8 is a protein called 60S ribosomal protein L10.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
8	G	199	1541	986	268	282	5	0	0

- Molecule 9 is a protein called 40S ribosomal protein S6E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	C	226	1820	1142	350	325	3	0	0

- Molecule 10 is a protein called 60S ribosomal protein L11.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	H	138	1037	651	190	194	2	0	0

- Molecule 11 is a protein called 40S ribosomal protein S24E.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	I	136	1004	628	189	180	7	0	0

- Molecule 12 is a protein called 40S ribosomal protein S24-A.


Mol	Chain	Residues	Atoms			AltConf	Trace	
			Total	C	N			O
12	D	134	1074	676	208	190	0	0

- Molecule 13 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
			Total	Mg	
13	B	1	1	1	0


- Molecule 14 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:

$C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms				AltConf	
			Total	C	N	O		P
14	B	1	31	10	5	13	3	0

- Molecule 15 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula:  $Fe_4S_4$ ).



Mol	Chain	Residues	Atoms			AltConf
			Total	Fe	S	
15	B	1	8	4	4	0
15	B	1	8	4	4	0


- Molecule 16 is water.

Mol	Chain	Residues	Atoms		AltConf
16	B	1	Total	O	0
			1	1	


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Dom34p




#### • Molecule 2: Rli1p








• Molecule 3: 28S ribosomal RNA




• Molecule 4: 18S ribosomal RNA






• Molecule 5: P-site tRNA





• Molecule 6: 60S ribosomal protein L6




• Molecule 7: 40S ribosomal protein S30E




• Molecule 8: 60S ribosomal protein L10







• Molecule 9: 40S ribosomal protein S6E



• Molecule 10: 60S ribosomal protein L11





• Molecule 11: 40S ribosomal protein S24E





- Molecule 12: 40S ribosomal protein S24-A



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	45700	Depositor
Resolution determination method	Not provided	
CTF correction method	Not provided	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	25	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	4500	Depositor
Magnification	75000	Depositor
Image detector	FEI EAGLE (4k x 4k)	Depositor
Maximum map value	1.731	Depositor
Minimum map value	-0.781	Depositor
Average map value	0.022	Depositor
Map value standard deviation	0.147	Depositor
Recommended contour level	0.32	Depositor
Map size ( $\text{\AA}$ )	455.4, 455.4, 455.4	wwPDB
Map dimensions	368, 368, 368	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	1.2375, 1.2375, 1.2375	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ATP, MG, SF4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.68	20/3149 (0.6%)	2.15	114/4230 (2.7%)
2	B	0.99	6/4893 (0.1%)	1.18	19/6603 (0.3%)
3	J	1.80	184/5523 (3.3%)	2.28	379/8591 (4.4%)
4	K	1.79	103/3671 (2.8%)	2.46	282/5709 (4.9%)
5	L	2.52	94/1781 (5.3%)	2.58	165/2775 (5.9%)
6	F	1.03	3/1540 (0.2%)	1.12	9/2073 (0.4%)
7	E	0.95	1/447 (0.2%)	1.29	7/595 (1.2%)
8	G	1.73	17/1568 (1.1%)	2.33	55/2119 (2.6%)
9	C	0.70	2/1844 (0.1%)	0.84	2/2464 (0.1%)
10	H	2.17	9/1048 (0.9%)	2.61	95/1408 (6.7%)
11	I	1.01	2/1019 (0.2%)	1.04	4/1369 (0.3%)
12	D	0.68	1/1088 (0.1%)	0.87	4/1449 (0.3%)
All	All	1.58	442/27571 (1.6%)	1.99	1135/39385 (2.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	18
2	B	0	4
3	J	0	25
4	K	0	20
5	L	0	5
6	F	0	2
7	E	0	2
8	G	0	13
10	H	0	13
12	D	0	1
All	All	0	103

The worst 5 of 442 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
10	H	88	PRO	CG-CD	50.97	3.18	1.50
5	L	41	G	C2'-C1'	-21.76	1.29	1.53
8	G	5	ARG	C-O	-20.62	0.84	1.23
5	L	22	G	C2'-C1'	-15.84	1.35	1.53
8	G	72	ASP	C-O	-15.71	0.93	1.23

The worst 5 of 1135 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	K	2845	A	N1-C6-N6	40.71	143.02	118.60
4	K	2845	A	C6-N1-C2	39.69	142.41	118.60
4	K	2845	A	C5-C6-N1	-35.84	99.78	117.70
8	G	5	ARG	O-C-N	-35.31	66.20	122.70
10	H	88	PRO	N-CD-CG	-34.39	51.62	103.20

There are no chirality outliers.

5 of 103 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	104	TYR	Sidechain
1	A	113	TYR	Sidechain
1	A	158	THR	Peptide
1	A	170	TYR	Sidechain
1	A	48	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3097	0	3156	158	0
2	B	4804	0	4938	988	0
3	J	4942	0	2531	522	0
4	K	3286	0	1680	563	0
5	L	1595	0	808	24	0
6	F	1519	0	1587	73	0
7	E	440	0	488	18	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	G	1541	0	1584	83	0
9	C	1820	0	1916	115	0
10	H	1037	0	1107	283	0
11	I	1004	0	1048	93	0
12	D	1074	0	1132	22	0
13	B	1	0	0	2	0
14	B	31	0	12	40	0
15	B	16	0	0	24	0
16	B	1	0	0	0	0
All	All	26208	0	21987	2612	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 54.

The worst 5 of 2612 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:H:85:LEU:CD2	10:H:87:GLU:H	1.02	1.62
2:B:56:ILE:HG12	15:B:703:SF4:S3	1.41	1.60
8:G:43:LYS:HG2	10:H:121:PHE:CD1	1.31	1.58
8:G:107:ALA:HB3	8:G:183:PHE:CE2	1.36	1.57
10:H:85:LEU:CD2	10:H:86:LYS:H	1.10	1.56

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	384/386 (100%)	289 (75%)	50 (13%)	45 (12%)	<b>0</b>   <b>6</b>
2	B	606/608 (100%)	561 (93%)	29 (5%)	16 (3%)	<b>5</b>   <b>31</b>

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
6	F	189/191 (99%)	167 (88%)	16 (8%)	6 (3%)	4	26
7	E	53/63 (84%)	42 (79%)	7 (13%)	4 (8%)	1	13
8	G	197/312 (63%)	170 (86%)	15 (8%)	12 (6%)	1	17
9	C	224/236 (95%)	190 (85%)	22 (10%)	12 (5%)	2	19
10	H	136/165 (82%)	47 (35%)	35 (26%)	54 (40%)	0	0
11	I	134/137 (98%)	124 (92%)	9 (7%)	1 (1%)	22	63
12	D	132/135 (98%)	106 (80%)	13 (10%)	13 (10%)	0	9
All	All	2055/2233 (92%)	1696 (82%)	196 (10%)	163 (8%)	2	12

5 of 163 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	ASP
1	A	15	GLY
1	A	46	LYS
1	A	48	THR
1	A	56	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/348 (100%)	318 (91%)	30 (9%)	10	32
2	B	537/537 (100%)	517 (96%)	20 (4%)	34	58
6	F	171/171 (100%)	133 (78%)	38 (22%)	1	6
7	E	48/54 (89%)	40 (83%)	8 (17%)	2	12
8	G	167/254 (66%)	161 (96%)	6 (4%)	35	59
9	C	193/201 (96%)	154 (80%)	39 (20%)	1	7
10	H	112/136 (82%)	96 (86%)	16 (14%)	3	16
11	I	104/105 (99%)	88 (85%)	16 (15%)	2	14
12	D	112/113 (99%)	86 (77%)	26 (23%)	1	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1792/1919 (93%)	1593 (89%)	199 (11%)	9 22

5 of 199 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
9	C	105	ASP
10	H	31	LYS
9	C	124	LEU
9	C	155	ASP
10	H	68	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
10	H	66	ASN
11	I	98	ASN
10	H	68	GLN
10	H	115	GLN
2	B	262	GLN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
3	J	224/233 (96%)	113 (50%)	26 (11%)
4	K	149/155 (96%)	86 (57%)	13 (8%)
5	L	74/75 (98%)	26 (35%)	10 (13%)
All	All	447/463 (96%)	225 (50%)	49 (10%)

5 of 225 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
3	J	40	A
3	J	41	A
3	J	42	G
3	J	45	U
3	J	46	A

5 of 49 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
4	K	1272	C
4	K	2851	A
4	K	2254	U
4	K	2297	U
4	K	3030	G

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
15	SF4	B	704	2	0,12,12	-	-	-	-	-
14	ATP	B	702	13	26,33,33	2.17	8 (30%)	31,52,52	3.29	11 (35%)
15	SF4	B	703	2	0,12,12	-	-	-	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
15	SF4	B	704	2	-	-	0/6/5/5
14	ATP	B	702	13	-	2/18/38/38	0/3/3/3
15	SF4	B	703	2	-	-	0/6/5/5

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	B	702	ATP	C4-N3	6.03	1.44	1.35
14	B	702	ATP	O5'-C5'	-4.55	1.27	1.44
14	B	702	ATP	O4'-C1'	3.92	1.46	1.41
14	B	702	ATP	PA-O5'	-2.97	1.47	1.59
14	B	702	ATP	PB-O2B	-2.54	1.43	1.55

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	B	702	ATP	O5'-C5'-C4'	12.29	151.30	108.99
14	B	702	ATP	O5'-PA-O1A	-6.50	83.69	109.07
14	B	702	ATP	PA-O5'-C5'	5.65	154.82	121.68
14	B	702	ATP	C5'-C4'-C3'	-5.23	95.59	115.18
14	B	702	ATP	O4'-C4'-C3'	4.70	114.42	105.11

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
14	B	702	ATP	C4'-C5'-O5'-PA
14	B	702	ATP	PG-O3B-PB-O1B


There are no ring outliers.

3 monomers are involved in 64 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
15	B	704	SF4	6	0
14	B	702	ATP	40	0
15	B	703	SF4	18	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-2010. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 213



Y Index: 187




Z Index: 173


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.32. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

## 6.6 Mask visualisation [i](#)


This section was not generated. No masks/segmentation were deposited.



## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is 3997  $\text{nm}^3$ ; this corresponds to an approximate mass of 3610 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of  $0.139 \text{\AA}^{-1}$


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-2010 and PDB model 3J16. Per-residue inclusion information can be found in section 3 on page 8.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.32 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.32).



























## 9.4 Atom inclusion [i](#)



At the recommended contour level, 77% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.32) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7360	 0.1180
A	 0.5330	 0.0930
B	 0.5870	 0.0900
C	 0.8300	 0.1020
D	 0.7000	 0.0830
E	 0.8660	 0.1320
F	 0.7730	 0.1020
G	 0.6770	 0.0570
H	 0.7150	 0.1050
I	 0.7300	 0.1050
J	 0.9860	 0.1790
K	 0.9490	 0.1680
L	 0.3030	 0.0930

