



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 01:19 PM JST

PDB ID : 6J4J  
Title : soybean seed H-2 ferritin  
Authors : Zhang, X.; Zang, J.; Chen, H.; Zhao, G.  
Deposited on : 2019-01-09  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtrriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	209	74% 8% • 16%
1	B	209	75% 9% 16%
1	C	209	74% 8% • 16%
1	D	209	74% 8% • 16%
1	E	209	78% 5% • 16%
1	H	209	75% 8% 16%

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 8976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ferritin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	175	1427	906	245	275	1	0	0	0
1	A	175	1427	906	245	275	1	0	0	0
1	B	175	1427	906	245	275	1	0	0	0
1	C	175	1427	906	245	275	1	0	0	0
1	D	175	1427	906	245	275	1	0	0	0
1	E	175	1427	906	245	275	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	155	ASP	GLU	engineered mutation	UNP I1J7H3
A	155	ASP	GLU	engineered mutation	UNP I1J7H3
B	155	ASP	GLU	engineered mutation	UNP I1J7H3
C	155	ASP	GLU	engineered mutation	UNP I1J7H3
D	155	ASP	GLU	engineered mutation	UNP I1J7H3
E	155	ASP	GLU	engineered mutation	UNP I1J7H3

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	H	2	Total 2	Mg 2	0	0
2	A	2	Total 2	Mg 2	0	0
2	B	2	Total 2	Mg 2	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	2	Total Mg 2 2	0	0
2	D	1	Total Mg 1 1	0	0
2	E	1	Total Mg 1 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	H	66	Total O 66 66	0	0
3	A	70	Total O 70 70	0	0
3	B	67	Total O 67 67	0	0
3	C	69	Total O 69 69	0	0
3	D	64	Total O 64 64	0	0
3	E	68	Total O 68 68	0	0






- Molecule 1: Ferritin

Chain D: 74% 8% 16%



- Molecule 1: Ferritin

Chain E: 78% 5% 16%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	130.83Å 130.83Å 171.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.70 – 2.10 46.25 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.5 (40.70-2.10) 99.5 (46.25-2.10)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.10 (at 2.10Å)	Xtrriage
Refinement program	PHENIX (1.10.1_2155: ???)	Depositor
R, $R_{free}$	0.160 , 0.191 0.160 , 0.191	Depositor DCC
$R_{free}$ test set	4203 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.6	Xtrriage
Anisotropy	0.204	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 47.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.022 for -h,k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8976	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	2/1458 (0.1%)	0.48	0/1969
1	B	0.38	0/1458	0.47	0/1969
1	C	0.37	0/1458	0.47	0/1969
1	D	0.37	0/1458	0.48	0/1969
1	E	0.39	0/1458	0.48	0/1969
1	H	0.38	0/1458	0.50	1/1969 (0.1%)
All	All	0.40	2/8748 (0.0%)	0.48	1/11814 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	125	GLU	CD-OE1	-8.32	1.16	1.25
1	A	125	GLU	CD-OE2	-7.13	1.17	1.25

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	133	MET	CG-SD-CE	-5.16	91.94	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1427	0	1378	17	0
1	B	1427	0	1378	16	0
1	C	1427	0	1378	15	0
1	D	1427	0	1378	17	0
1	E	1427	0	1378	11	0
1	H	1427	0	1378	16	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	H	2	0	0	0	0
3	A	70	0	0	2	0
3	B	67	0	0	2	0
3	C	69	0	0	4	0
3	D	64	0	0	5	0
3	E	68	0	0	4	0
3	H	66	0	0	3	0
All	All	8976	0	8268	76	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (76) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:33:ASN:N	3:E:401:HOH:O	2.12	0.82
1:H:64:HIS:CE1	1:A:96:GLU:HG2	2.16	0.81
1:H:33:ASN:N	3:H:401:HOH:O	2.15	0.79
1:H:64:HIS:HE1	1:A:96:GLU:HG2	1.50	0.77
1:A:33:ASN:N	3:A:401:HOH:O	2.19	0.76
1:E:80:LEU:HG	3:E:468:HOH:O	1.88	0.73
1:B:89:GLU:HG3	1:B:92:ARG:HH21	1.56	0.70
1:H:140:GLU:HG3	1:H:181:ILE:HD12	1.74	0.70
1:H:92:ARG:HE	1:A:92:ARG:HE	1.39	0.68
1:D:33:ASN:N	3:D:402:HOH:O	2.27	0.66
1:E:140:GLU:HG3	1:E:181:ILE:HD12	1.77	0.66
1:D:133:MET:HE3	3:D:464:HOH:O	1.96	0.65
1:C:108:ARG:NH1	3:C:401:HOH:O	2.21	0.64
1:C:140:GLU:HG3	1:C:181:ILE:HD12	1.78	0.64
1:B:140:GLU:HG3	1:B:181:ILE:HD12	1.79	0.63
1:A:125:GLU:H	1:A:125:GLU:CD	2.03	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:133:MET:HE2	3:E:468:HOH:O	2.00	0.61
1:B:89:GLU:HG3	1:B:92:ARG:NH2	2.17	0.60
1:A:89:GLU:HG3	1:A:92:ARG:HH22	1.67	0.59
1:A:89:GLU:HG3	1:A:92:ARG:NH2	2.17	0.59
1:D:140:GLU:HG3	1:D:181:ILE:HD12	1.83	0.59
1:D:96:GLU:HG2	1:E:64:HIS:CE1	2.39	0.58
1:C:89:GLU:HG3	1:C:92:ARG:NH2	2.18	0.57
1:B:80:LEU:HD12	1:B:204:LEU:HD23	1.86	0.56
1:H:92:ARG:HE	1:A:92:ARG:NE	2.01	0.56
1:C:33:ASN:N	3:C:402:HOH:O	2.40	0.54
1:C:203:LYS:HE2	1:C:207:ASP:OD1	2.10	0.52
1:H:64:HIS:ND1	1:A:57:TYR:OH	2.43	0.51
1:H:113:PRO:HD3	1:A:116:SER:HB2	1.92	0.51
1:B:64:HIS:ND1	1:C:57:TYR:OH	2.43	0.51
1:B:114:ILE:HB	1:C:114:ILE:HB	1.93	0.51
1:D:64:HIS:NE2	1:E:96:GLU:HG2	2.27	0.49
1:A:99:ILE:HG23	1:A:109:VAL:HG21	1.95	0.49
1:B:57:TYR:OH	1:C:64:HIS:ND1	2.45	0.48
1:C:80:LEU:HD12	1:C:201:ASP:OD1	2.13	0.48
1:H:33:ASN:N	1:H:33:ASN:OD1	2.45	0.48
1:H:62:VAL:HG22	1:H:117:PRO:HB3	1.97	0.47
1:C:89:GLU:HG3	1:C:92:ARG:HH21	1.79	0.47
1:B:99:ILE:HG23	1:B:109:VAL:HG21	1.97	0.47
1:E:61:TYR:CD2	1:E:117:PRO:HD3	2.50	0.47
1:B:92:ARG:O	1:B:96:GLU:HG3	2.15	0.47
1:D:207:ASP:OD2	3:D:401:HOH:O	2.21	0.46
1:B:161:GLN:OE1	3:B:401:HOH:O	2.20	0.46
1:C:92:ARG:NH1	3:C:403:HOH:O	2.40	0.45
1:D:200:PHE:HE2	3:D:464:HOH:O	1.97	0.45
1:A:33:ASN:N	3:A:405:HOH:O	2.49	0.44
1:D:111:LEU:HB3	1:E:61:TYR:CZ	2.52	0.44
1:D:203:LYS:NZ	1:D:207:ASP:OD2	2.49	0.44
1:D:204:LEU:HD12	1:D:204:LEU:HA	1.88	0.44
1:B:101:TYR:HE1	3:B:401:HOH:O	2.01	0.43
1:A:203:LYS:HG3	1:A:204:LEU:N	2.33	0.43
1:D:62:VAL:HG22	1:D:117:PRO:HB3	1.99	0.43
1:H:146:LYS:HA	1:H:146:LYS:HD2	1.87	0.43
1:A:112:HIS:HB3	1:A:113:PRO:HD2	2.00	0.43
1:E:108:ARG:HG2	3:E:423:HOH:O	2.18	0.43
1:C:172:TYR:CZ	1:C:176:LYS:HE3	2.53	0.43
1:H:96:GLU:HG2	1:A:64:HIS:NE2	2.34	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140:GLU:HG3	1:A:181:ILE:HD12	1.99	0.42
1:D:80:LEU:HD21	3:D:464:HOH:O	2.18	0.42
1:D:61:TYR:CD2	1:D:117:PRO:HD3	2.54	0.42
1:H:108:ARG:NH1	3:H:403:HOH:O	2.43	0.42
1:H:185:VAL:HG23	3:H:402:HOH:O	2.18	0.42
1:B:113:PRO:HD3	1:C:116:SER:HB2	2.01	0.42
1:D:61:TYR:CZ	1:E:111:LEU:HB3	2.55	0.42
1:E:33:ASN:O	1:E:108:ARG:NH2	2.53	0.42
1:H:77:LEU:HD23	1:H:77:LEU:HA	1.93	0.41
1:B:125:GLU:HG3	1:B:126:LYS:N	2.36	0.41
1:C:78:LYS:HD3	3:C:409:HOH:O	2.21	0.41
1:H:204:LEU:HD12	1:H:204:LEU:HA	1.94	0.41
1:B:92:ARG:HD3	1:C:92:ARG:HD3	2.02	0.41
1:D:33:ASN:O	1:D:108:ARG:NH1	2.53	0.41
1:B:204:LEU:HD12	1:B:204:LEU:HA	1.88	0.41
1:D:60:SER:HB2	1:D:91:GLU:HB3	2.02	0.41
1:A:82:LYS:HE2	1:A:205:LEU:O	2.22	0.40
1:B:85:LYS:HE3	1:B:85:LYS:HB2	1.95	0.40
1:D:146:LYS:HD3	1:D:146:LYS:HA	1.85	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	173/209 (83%)	171 (99%)	2 (1%)	0	100	100
1	B	173/209 (83%)	172 (99%)	1 (1%)	0	100	100
1	C	173/209 (83%)	172 (99%)	1 (1%)	0	100	100
1	D	173/209 (83%)	172 (99%)	1 (1%)	0	100	100
1	E	173/209 (83%)	172 (99%)	1 (1%)	0	100	100

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	173/209 (83%)	172 (99%)	1 (1%)	0	100	100
All	All	1038/1254 (83%)	1031 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	153/180 (85%)	148 (97%)	5 (3%)	38	40
1	B	153/180 (85%)	151 (99%)	2 (1%)	69	75
1	C	153/180 (85%)	147 (96%)	6 (4%)	32	33
1	D	153/180 (85%)	149 (97%)	4 (3%)	46	50
1	E	153/180 (85%)	150 (98%)	3 (2%)	55	60
1	H	153/180 (85%)	149 (97%)	4 (3%)	46	50
All	All	918/1080 (85%)	894 (97%)	24 (3%)	46	50

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	61	TYR
1	H	64	HIS
1	H	139	LEU
1	H	176	LYS
1	A	61	TYR
1	A	92	ARG
1	A	113	PRO
1	A	152	SER
1	A	203	LYS
1	B	61	TYR
1	B	64	HIS
1	C	33	ASN
1	C	61	TYR

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	64	HIS
1	C	90	GLU
1	C	108	ARG
1	C	199	HIS
1	D	61	TYR
1	D	64	HIS
1	D	133	MET
1	D	139	LEU
1	E	61	TYR
1	E	64	HIS
1	E	176	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	64	HIS
1	E	33	ASN
1	E	64	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	175/209 (83%)	-0.36	0 100 100	26, 32, 44, 70	0
1	B	175/209 (83%)	-0.47	0 100 100	27, 33, 45, 58	0
1	C	175/209 (83%)	-0.44	0 100 100	28, 33, 46, 63	0
1	D	175/209 (83%)	-0.46	0 100 100	27, 32, 45, 64	0
1	E	175/209 (83%)	-0.38	0 100 100	27, 31, 43, 61	0
1	H	175/209 (83%)	-0.49	0 100 100	28, 32, 45, 63	0
All	All	1050/1254 (83%)	-0.43	0 100 100	26, 32, 45, 70	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	A	302	1/1	0.88	0.07	57,57,57,57	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	MG	B	302	1/1	0.95	0.07	57,57,57,57	0
2	MG	E	301	1/1	0.96	0.11	39,39,39,39	0
2	MG	D	301	1/1	0.97	0.13	47,47,47,47	0
2	MG	B	301	1/1	0.98	0.05	39,39,39,39	0
2	MG	H	302	1/1	0.98	0.23	40,40,40,40	0
2	MG	C	301	1/1	0.98	0.27	45,45,45,45	0
2	MG	C	302	1/1	0.98	0.07	37,37,37,37	0
2	MG	A	301	1/1	0.98	0.09	37,37,37,37	0
2	MG	H	301	1/1	0.98	0.07	39,39,39,39	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.