

wwPDB X-ray Structure Validation Summary Report (i)

Oct 22, 2023 – 12:58 PM JST

structure of Flavihumibacter petaseus GH31 alpha-galactosidase mu-
304A in complex with alpha-1,4-galactobiose
, M.; Miyazaki, T.
I-21
(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity Mogul Xtriage (Phenix) EDS	:	4.02b-467 1.8.5 (274361), CSD as541be (2020) 1.13 2.36
buster-report Percentile statistics Refmac	: : :	1.1.7 (2018) 20191225.v01 (using entries in the PDB archive December 25th 2019) 5.8.0158 7.0.044 (Gargrove)
Ideal geometry (DNA, RNA) Validation Pipeline (wwPDB-VP)		Parkinson et al. (1996) 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{l} \textbf{Whole archive} \\ (\#\textbf{Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	541	.%	7%	• 6%
1	В	541	88%	6%	6%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 17074 atoms, of which 8080 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GH31 alpha-galactosidase.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
1	А	510	Total 8073	C 2612		N 704	0 732	S 23	91	3	0
1	В	508	Total 8003	C 2592		N 698	0 724	S 23	89	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	6	MET	-	initiating methionine	UNP A0A0E9MUN5
А	7	GLY	-	expression tag	UNP A0A0E9MUN5
А	8	SER	-	expression tag	UNP A0A0E9MUN5
А	9	SER	-	expression tag	UNP A0A0E9MUN5
А	10	HIS	-	expression tag	UNP A0A0E9MUN5
A	11	HIS	-	expression tag	UNP A0A0E9MUN5
А	12	HIS	-	expression tag	UNP A0A0E9MUN5
A	13	HIS	-	expression tag	UNP A0A0E9MUN5
А	14	HIS	-	expression tag	UNP A0A0E9MUN5
А	15	HIS	-	expression tag	UNP A0A0E9MUN5
А	16	SER	-	expression tag	UNP A0A0E9MUN5
А	17	SER	-	expression tag	UNP A0A0E9MUN5
А	18	GLY	-	expression tag	UNP A0A0E9MUN5
А	19	LEU	-	expression tag	UNP A0A0E9MUN5
А	20	VAL	-	expression tag	UNP A0A0E9MUN5
А	21	PRO	-	expression tag	UNP A0A0E9MUN5
А	22	ARG	-	expression tag	UNP A0A0E9MUN5
А	23	GLY	-	expression tag	UNP A0A0E9MUN5
А	24	SER	-	expression tag	UNP A0A0E9MUN5
А	25	HIS	-	expression tag	UNP A0A0E9MUN5
А	26	MET	-	expression tag	UNP A0A0E9MUN5
А	27	ALA	-	expression tag	UNP A0A0E9MUN5
А	28	SER	-	expression tag	UNP A0A0E9MUN5
А	304	ALA	ASP	engineered mutation	UNP A0A0E9MUN5
В	6	MET	-	initiating methionine	UNP A0A0E9MUN5

There are 48 discrepancies between the modelled and reference sequences:


Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
В	7	GLY	-	expression tag	UNP A0A0E9MUN5
В	8	SER	-	expression tag	UNP A0A0E9MUN5
В	9	SER	-	expression tag	UNP A0A0E9MUN5
В	10	HIS	-	expression tag	UNP A0A0E9MUN5
В	11	HIS	-	expression tag	UNP A0A0E9MUN5
В	12	HIS	-	expression tag	UNP A0A0E9MUN5
В	13	HIS	-	expression tag	UNP A0A0E9MUN5
В	14	HIS	-	expression tag	UNP A0A0E9MUN5
В	15	HIS	-	expression tag	UNP A0A0E9MUN5
В	16	SER	-	expression tag	UNP A0A0E9MUN5
В	17	SER	-	expression tag	UNP A0A0E9MUN5
В	18	GLY	-	expression tag	UNP A0A0E9MUN5
В	19	LEU	-	expression tag	UNP A0A0E9MUN5
В	20	VAL	-	expression tag	UNP A0A0E9MUN5
В	21	PRO	-	expression tag	UNP A0A0E9MUN5
В	22	ARG	-	expression tag	UNP A0A0E9MUN5
В	23	GLY	-	expression tag	UNP A0A0E9MUN5
В	24	SER	-	expression tag	UNP A0A0E9MUN5
В	25	HIS	-	expression tag	UNP A0A0E9MUN5
В	26	MET	-	expression tag	UNP A0A0E9MUN5
В	27	ALA	-	expression tag	UNP A0A0E9MUN5
В	28	SER	-	expression tag	UNP A0A0E9MUN5
В	304	ALA	ASP	engineered mutation	UNP A0A0E9MUN5

Continued from previous page...


• Molecule 2 is alpha-D-galactopyranose (three-letter code: GLA) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).






M	ol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	2	А	1	Total 22				4	0
2	2	В	1	Total 22		H 11	O 5	4	0

• Molecule 3 is beta-D-galactopyranose (three-letter code: GAL) (formula: $C_6H_{12}O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	А	1	Total				4	0
		_	24	-		•	_	
3	В	1	Total	\mathbf{C}	Η	Ο	4	0
5	D	T	24	6	12	6	4	0





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total C H O 10 2 6 2	1	0
4	А	1	Total C H O 10 2 6 2	1	0
4	А	1	Total C H O 10 2 6 2	1	0
4	А	1	Total C H O 10 2 6 2	1	0
4	А	1	Total C H O 10 2 6 2	1	0
4	В	1	Total C H O 10 2 6 2	1	0
4	В	1	Total C H O 10 2 6 2	1	0
4	В	1	Total C H O 10 2 6 2	1	0
4	В	1	Total C H O 10 2 6 2	1	0
4	В	1	Total C H O 10 2 6 2	1	0
4	В	1	Total C H O 10 2 6 2	1	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	А	1	Total 1	Cl 1	0	0
				Со		next page

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Cl 1 1	0	0


• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	409	Total O 409 409	0	0
6	В	385	Total O 385 385	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: GH31 alpha-galactosidase



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants	71.50Å 72.54Å 81.20Å	Depositor
a, b, c, α , β , γ	101.92° 104.40° 103.71°	-
Resolution (Å)	46.04 - 1.90	Depositor
	46.00 - 1.90	EDS
% Data completeness	$97.4 \ (46.04 - 1.90)$	Depositor
(in resolution range)	$97.4 \ (46.00 - 1.90)$	EDS
R _{merge}	0.07	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.11 (at 1.89 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0405	Depositor
D D	0.136 , 0.192	Depositor
R, R_{free}	0.147 , 0.199	DCC
R_{free} test set	5585 reflections (4.94%)	wwPDB-VP
Wilson B-factor $(Å^2)$	13.8	Xtriage
Anisotropy	1.256	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.45 , 48.0	EDS
L-test for twinning ²	$< L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	0.003 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	17074	wwPDB-VP
Average B, all atoms $(Å^2)$	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.03% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GAL, CL, GLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	Bond lengths		ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.70	5/4191~(0.1%)	0.89	7/5679~(0.1%)
1	В	0.65	3/4148~(0.1%)	0.87	9/5622~(0.2%)
All	All	0.67	8/8339~(0.1%)	0.88	16/11301~(0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	5
1	В	0	3
All	All	0	8

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
1	А	517	GLU	CD-OE2	8.60	1.35	1.25
1	В	287	GLU	CD-OE2	7.62	1.34	1.25
1	В	325	GLU	CD-OE1	7.17	1.33	1.25
1	А	287	GLU	CD-OE2	7.07	1.33	1.25
1	А	236	GLU	CD-OE2	-6.41	1.18	1.25

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	198	ARG	NE-CZ-NH2	-9.70	115.45	120.30
1	В	358	ARG	NE-CZ-NH1	8.32	124.46	120.30
1	В	459	ARG	NE-CZ-NH1	7.78	124.19	120.30
1	В	296	TYR	CB-CG-CD1	7.20	125.32	121.00

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$Ideal(^{o})$
1	А	368	ARG	NE-CZ-NH1	6.82	123.71	120.30

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	448	ARG	Sidechain
1	А	483	ARG	Sidechain
1	А	506	ARG	Sidechain
1	А	53[A]	ARG	Sidechain
1	А	53[B]	ARG	Sidechain

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	4071	4002	3994	16	0
1	В	4037	3966	3955	12	0
2	А	11	11	10	0	0
2	В	11	11	10	2	0
3	А	12	12	11	0	0
3	В	12	12	11	2	0
4	А	20	30	30	3	0
4	В	24	36	36	0	0
5	А	1	0	0	0	0
5	В	1	0	0	0	0
6	А	409	0	0	0	0
6	В	385	0	0	5	0
All	All	8994	8080	8057	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 29 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:601:GLA:C1	3:B:602:GAL:O4	1.63	1.47
2:B:601:GLA:C1	3:B:602:GAL:C4	2.60	0.78
1:A:331:GLY:HA2	1:A:342:TYR:CE2	2.37	0.59
1:B:99:GLU:HG3	6:B:950:HOH:O	2.04	0.58
1:B:288:LYS:HE2	6:B:939:HOH:O	2.04	0.57

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	509/541~(94%)	494 (97%)	13 (3%)	2~(0%)	34 24
1	В	504/541~(93%)	489 (97%)	13 (3%)	2(0%)	34 24
All	All	1013/1082~(94%)	983~(97%)	26 (3%)	4 (0%)	34 24

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	187	ASN
1	В	187	ASN
1	А	157	ILE
1	В	157	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	427/450~(95%)	424 (99%)	3~(1%)	84 84		
1	В	422/450 (94%)	418 (99%)	4 (1%)	78 79		
All	All	849/900~(94%)	842~(99%)	7 (1%)	81 82		

5 of 7 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	В	82	SER
1	В	185	ASP
1	В	359	ASP
1	В	307	ASP
1	А	359	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	А	257	HIS
1	В	74	ASN
1	В	257	HIS
1	А	60	GLN
1	А	33	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 17 ligands modelled in this entry, 2 are monoatomic - leaving 15 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Turne	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	les
1VIOI	Type	Unam	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	GLA	В	601	-	$11,\!11,\!12$	1.73	2 (18%)	$15,\!15,\!17$	1.19	1 (6%)
4	EDO	А	603	-	$3,\!3,\!3$	1.54	1 (33%)	$2,\!2,\!2$	1.18	0
4	EDO	В	605	-	3,3,3	1.31	0	2,2,2	0.78	0
3	GAL	А	602	-	12,12,12	0.91	1 (8%)	17,17,17	0.68	0
4	EDO	А	606	-	3,3,3	1.46	0	2,2,2	0.95	0
4	EDO	А	605	-	3,3,3	0.09	0	2,2,2	0.01	0
3	GAL	В	602	-	$12,\!12,\!12$	0.95	0	$17,\!17,\!17$	0.80	0
4	EDO	В	607	-	3, 3, 3	0.32	0	$2,\!2,\!2$	0.47	0
4	EDO	В	604	-	3, 3, 3	0.40	0	$2,\!2,\!2$	0.69	0
4	EDO	В	606	-	3, 3, 3	0.72	0	$2,\!2,\!2$	0.57	0
4	EDO	А	604	-	$3,\!3,\!3$	0.75	0	$2,\!2,\!2$	0.24	0
2	GLA	А	601	-	11,11,12	1.16	1 (9%)	$15,\!15,\!17$	0.97	0
4	EDO	В	603	-	3,3,3	0.10	0	2,2,2	0.32	0
4	EDO	В	608	-	3,3,3	0.61	0	2,2,2	0.40	0
4	EDO	А	607	-	$3,\!3,\!3$	0.78	0	2,2,2	0.97	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLA	В	601	-	-	0/2/19/22	0/1/1/1
4	EDO	А	603	-	-	1/1/1/1	-
4	EDO	В	605	-	-	0/1/1/1	-
3	GAL	А	602	-	-	0/2/22/22	0/1/1/1
4	EDO	А	606	-	-	0/1/1/1	-
4	EDO	А	605	-	-	0/1/1/1	-
3	GAL	В	602	-	-	0/2/22/22	0/1/1/1
4	EDO	В	607	-	-	0/1/1/1	-
4	EDO	В	604	-	-	0/1/1/1	-
4	EDO	В	606	-	-	0/1/1/1	-
4	EDO	А	604	-	-	0/1/1/1	-
2	GLA	А	601	-	-	0/2/19/22	0/1/1/1

Continued on next page...



001000	Continuacia from pretious page									
Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings			
4	EDO	В	603	-	-	0/1/1/1	-			
4	EDO	В	608	-	-	0/1/1/1	-			
4	EDO	А	607	-	-	1/1/1/1	-			

Continued from previous page...

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	В	601	GLA	C4-C5	3.46	1.60	1.53
2	В	601	GLA	O5-C5	2.96	1.49	1.43
2	А	601	GLA	C4-C5	2.38	1.58	1.53
3	А	602	GAL	O4-C4	2.26	1.48	1.43
4	А	603	EDO	O1-C1	2.10	1.52	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	В	601	GLA	O4-C4-C3	-3.15	103.06	110.35

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	А	603	EDO	O1-C1-C2-O2
4	А	607	EDO	O1-C1-C2-O2

There are no ring outliers.


3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	601	GLA	2	0
3	В	602	GAL	2	0
4	А	607	EDO	3	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the




average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	510/541~(94%)	-0.53	3 (0%) 89 90	15, 22, 42, 76	0
1	В	508/541~(93%)	-0.47	16 (3%) 49 51	16, 23, 47, 73	0
All	All	1018/1082 (94%)	-0.50	19 (1%) 66 69	15, 23, 45, 76	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	546	THR	5.6
1	А	251	ASP	3.6
1	А	403	LYS	3.6
1	В	525	ARG	3.5
1	В	403	LYS	3.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	А	607	4/4	0.90	0.13	38,40,41,41	1
4	EDO	В	608	4/4	0.90	0.19	48,53,59,61	1
4	EDO	В	605	4/4	0.93	0.07	29,32,35,35	1
4	EDO	А	606	4/4	0.93	0.08	30,35,39,40	1
4	EDO	В	606	4/4	0.95	0.11	52,67,69,73	1
4	EDO	А	603	4/4	0.95	0.18	34,41,47,49	1
4	EDO	В	604	4/4	0.97	0.07	$25,\!31,\!38,\!41$	1
3	GAL	В	602	12/12	0.98	0.07	21,28,31,33	4
2	GLA	А	601	11/12	0.98	0.07	21,23,24,24	4
2	GLA	В	601	11/12	0.98	0.06	21,22,23,24	4
4	EDO	В	607	4/4	0.98	0.06	30,31,36,36	1
3	GAL	А	602	12/12	0.98	0.06	21,26,29,32	4
4	EDO	В	603	4/4	0.99	0.07	20,21,22,23	1
4	EDO	А	604	4/4	0.99	0.08	22,23,24,25	1
4	EDO	А	605	4/4	0.99	0.07	24,26,29,31	1
5	CL	А	608	1/1	1.00	0.04	28,28,28,28	0
5	CL	В	609	1/1	1.00	0.06	26,26,26,26	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



















6.5 Other polymers (i)

There are no such residues in this entry.

