

wwPDB X-ray Structure Validation Summary Report (i)

Jun 14, 2020 - 11:35 am BST

PDB ID	:	2JBF
Title	:	Structure of PBP-A, L158E mutant. Acyl-enzyme complex with penicillin- G.
Authors	:	Evrard, C.; Declercq, JP.
Deposited on		
Resolution	:	1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity		4.02b-467 1.8.5 (274361), CSD as541be (2020)
9		
Xtriage (Phenix)		1.13
EDS	:	2.11
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	4695(1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	298	% • 77%	11% •	11%
1	В	298	83%	5% •	11%
1	С	298	% • 78%	10%	11%
1	D	298	% • 80%	11%	• 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PNM	А	300	-	-	-	Х
2	PNM	С	300	-	-	Х	-
2	PNM	D	300	-	-	Х	Х



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 9215 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	264	Total	С	Ν	Ο	S	0	0	0
	A	204	2035	1277	360	387	11	0	0	0
1	В	265	Total	С	Ν	Ο	S	0	0	0
	D	205	2042	1281	361	389	11	0		0
1	С	264	Total	С	Ν	Ο	S	0	0	0
		204	2035	1277	360	387	11	0	0	0
1	П	274	Total	С	Ν	Ο	S	0	0	0
		$\begin{array}{c c} \hline \end{array}$ 274	2101	1317	370	403	11			U

• Molecule 1 is a protein called TLL2115 PROTEIN.

There are 92 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q8DH45
A	278	GLU	-	expression tag	UNP Q8DH45
A	279	GLN	-	expression tag	UNP Q8DH45
A	280	LYS	-	expression tag	UNP Q8DH45
A	281	LEU	-	expression tag	UNP Q8DH45
A	282	ILE	-	expression tag	UNP Q8DH45
A	283	SER	-	expression tag	UNP Q8DH45
A	284	GLU	-	expression tag	UNP Q8DH45
A	285	GLU	-	expression tag	UNP Q8DH45
A	286	ASP	-	expression tag	UNP Q8DH45
A	287	LEU	-	expression tag	UNP Q8DH45
A	288	ASN	-	expression tag	UNP Q8DH45
A	289	SER	-	expression tag	UNP Q8DH45
A	290	ALA	-	expression tag	UNP Q8DH45
A	291	VAL	-	expression tag	UNP Q8DH45
A	292	ASP	-	expression tag	UNP Q8DH45
A	293	HIS	-	expression tag	UNP Q8DH45
А	294	HIS	-	expression tag	UNP Q8DH45
А	295	HIS	-	expression tag	UNP Q8DH45
А	296	HIS	-	expression tag	UNP Q8DH45
А	297	HIS	-	expression tag	UNP Q8DH45



Chain	Residue	Modelled	Actual	Comment	Reference
А	298	HIS	-	expression tag	UNP Q8DH45
А	158	GLU	LEU	engineered mutation	UNP Q8DH45
В	1	MET	-	expression tag	UNP Q8DH45
В	278	GLU	-	expression tag	UNP Q8DH45
В	279	GLN	-	expression tag	UNP Q8DH45
В	280	LYS	-	expression tag	UNP Q8DH45
В	281	LEU	-	expression tag	UNP Q8DH45
В	282	ILE	-	expression tag	UNP Q8DH45
В	283	SER	-	expression tag	UNP Q8DH45
В	284	GLU	-	expression tag	UNP Q8DH45
В	285	GLU	-	expression tag	UNP Q8DH45
В	286	ASP	-	expression tag	UNP Q8DH45
В	287	LEU	-	expression tag	UNP Q8DH45
В	288	ASN	-	expression tag	UNP Q8DH45
В	289	SER	-	expression tag	UNP Q8DH45
В	290	ALA	-	expression tag	UNP Q8DH45
В	291	VAL	-	expression tag	UNP Q8DH45
В	292	ASP	-	expression tag	UNP Q8DH45
В	293	HIS	-	expression tag	UNP Q8DH45
В	294	HIS	-	expression tag	UNP Q8DH45
В	295	HIS	-	expression tag	UNP Q8DH45
В	296	HIS	-	expression tag	UNP Q8DH45
В	297	HIS	-	expression tag	UNP Q8DH45
В	298	HIS	-	expression tag	UNP Q8DH45
В	158	GLU	LEU	engineered mutation	UNP Q8DH45
С	1	MET	-	expression tag	UNP Q8DH45
С	278	GLU	-	expression tag	UNP Q8DH45
С	279	GLN	-	expression tag	UNP Q8DH45
С	280	LYS	-	expression tag	UNP Q8DH45
С	281	LEU	-	expression tag	UNP Q8DH45
С	282	ILE	-	expression tag	UNP Q8DH45
С	283	SER	-	expression tag	UNP Q8DH45
С	284	GLU	-	expression tag	UNP Q8DH45
С	285	GLU	-	expression tag	UNP Q8DH45
С	286	ASP	-	expression tag	UNP Q8DH45
С	287	LEU	-	expression tag	UNP Q8DH45
С	288	ASN	-	expression tag	UNP Q8DH45
С	289	SER	-	expression tag	UNP Q8DH45
	22.0	ALA	-	expression tag	UNP Q8DH45
С	290				•
C C	290 291	VAL	-	expression tag	UNP Q8DH45
			-	expression tag expression tag	-

Continued from previous page...




Chain	Residue	Modelled	Actual	Comment	Reference
С	294	HIS	-	expression tag	UNP Q8DH45
С	295	HIS	-	expression tag	UNP Q8DH45
С	296	HIS	-	expression tag	UNP Q8DH45
С	297	HIS	-	expression tag	UNP Q8DH45
С	298	HIS	-	expression tag	UNP Q8DH45
С	158	GLU	LEU	engineered mutation	UNP Q8DH45
D	1	MET	-	expression tag	UNP Q8DH45
D	278	GLU	-	expression tag	UNP Q8DH45
D	279	GLN	-	expression tag	UNP Q8DH45
D	280	LYS	-	expression tag	UNP Q8DH45
D	281	LEU	-	expression tag	UNP Q8DH45
D	282	ILE	-	expression tag	UNP Q8DH45
D	283	SER	-	expression tag	UNP Q8DH45
D	284	GLU	-	expression tag	UNP Q8DH45
D	285	GLU	-	expression tag	UNP Q8DH45
D	286	ASP	-	expression tag	UNP Q8DH45
D	287	LEU	-	expression tag	UNP Q8DH45
D	288	ASN	-	expression tag	UNP Q8DH45
D	289	SER	-	expression tag	UNP Q8DH45
D	290	ALA	-	expression tag	UNP Q8DH45
D	291	VAL	-	expression tag	UNP Q8DH45
D	292	ASP	-	expression tag	UNP Q8DH45
D	293	HIS	-	expression tag	UNP Q8DH45
D	294	HIS	-	expression tag	UNP Q8DH45
D	295	HIS	-	expression tag	UNP Q8DH45
D	296	HIS	-	expression tag	UNP Q8DH45
D	297	HIS	-	expression tag	UNP Q8DH45
D	298	HIS	-	expression tag	UNP Q8DH45
D	158	GLU	LEU	engineered mutation	UNP Q8DH45


Continued from previous page...

• Molecule 2 is OPEN FORM - PENICILLIN G (three-letter code: PNM) (formula: $C_{16}H_{20}N_2O_4S$).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	Δ	1	Total	С	Ν	Ο	S	0	0
	Л	T	23	16	2	4	1	0	0
2	В	1	Total	С	Ν	Ο	\mathbf{S}	0	0
	D	I	23	16	2	4	1	0	0
2	С	1	Total	С	Ν	Ο	\mathbf{S}	0	0
		I	23	16	2	4	1	0	0
2	п	1	Total	С	Ν	Ο	S	0	0
	D	L	23	16	2	4	1	U	0





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	А	1	Total 6	${ m C} { m 3}$	O 3	0	0


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	189	Total O 189 189	0	0
4	В	226	Total O 226 226	0	0
4	С	275	Total O 275 275	0	0
4	D	214	Total O 214 214	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: TLL2115 PROTEIN







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	86.58Å 91.53Å 145.71Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	77.61 - 1.70	Depositor
Resolution (A)	74.43 - 1.70	EDS
% Data completeness	99.3 (77.61-1.70)	Depositor
(in resolution range)	99.3(74.43-1.70)	EDS
R _{merge}	0.04	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.36 (at 1.70 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
D D.	0.178 , 0.218	Depositor
R, R_{free}	0.186 , (Not available)	DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor $(Å^2)$	22.1	Xtriage
Anisotropy	0.079	Xtriage
Bulk solvent $k_{sol}(e/A^3)$, $B_{sol}(A^2)$	0.35 , 47.6	EDS
L-test for twinning ²	$ \langle L \rangle = 0.46, \langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9215	wwPDB-VP
Average B, all atoms $(Å^2)$	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.73% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, $\rm PNM$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	Bond lengths Bon		ond angles
	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.65	0/2068	0.74	2/2805~(0.1%)
1	В	0.72	0/2075	0.81	3/2815~(0.1%)
1	С	0.76	0/2068	0.82	2/2805~(0.1%)
1	D	0.66	0/2137	0.77	3/2902~(0.1%)
All	All	0.70	0/8348	0.79	10/11327~(0.1%)

There are no bond length outliers.

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	D	79	ARG	NE-CZ-NH1	8.49	124.54	120.30
1	В	199	ARG	NE-CZ-NH1	7.88	124.24	120.30
1	В	199	ARG	NE-CZ-NH2	-7.38	116.61	120.30
1	D	79	ARG	NE-CZ-NH2	-7.25	116.67	120.30
1	А	79	ARG	NE-CZ-NH2	-6.36	117.12	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2035	0	2065	25	0
1	В	2042	0	2072	15	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	С	2035	0	2065	37	0
1	D	2101	0	2127	32	0
2	А	23	0	18	5	0
2	В	23	0	18	2	0
2	С	23	0	18	13	0
2	D	23	0	18	9	0
3	А	6	0	8	0	0
4	А	189	0	0	4	0
4	В	226	0	0	6	0
4	С	275	0	0	8	0
4	D	214	0	0	7	0
All	All	9215	0	8409	113	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 113 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:161:MET:SD	2:D:300:PNM:H171	2.14	0.87
1:D:161:MET:SD	2:D:300:PNM:C17	2.64	0.85
1:C:225:ILE:HG13	1:C:226:VAL:HG23	1.58	0.85
1:D:35:THR:HG22	1:D:248:LYS:HB2	1.60	0.83
1:C:161:MET:SD	2:C:300:PNM:H171	2.19	0.81

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	262/298~(88%)	254 (97%)	7 (3%)	1 (0%)	34 18



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	263/298~(88%)	257~(98%)	6 (2%)	0	100	100
1	С	262/298~(88%)	254 (97%)	8 (3%)	0	100	100
1	D	272/298~(91%)	266~(98%)	6 (2%)	0	100	100
All	All	1059/1192~(89%)	$1031 \ (97\%)$	27 (2%)	1 (0%)	51	33

Continued from previous page...

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	32	PRO

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	221/252~(88%)	215~(97%)	6 (3%)	44	26
1	В	222/252~(88%)	217~(98%)	5(2%)	50	33
1	С	221/252~(88%)	217~(98%)	4 (2%)	59	43
1	D	229/252~(91%)	221~(96%)	8 (4%)	36	17
All	All	893/1008 (89%)	870 (97%)	23 (3%)	46	28

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	200	THR
1	С	85	ARG
1	D	170	ARG
1	С	15	ARG
1	С	122	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:



Mol	Chain	Res	Type
1	В	269	GLN
1	D	262	GLN
1	С	218	HIS
1	В	181	GLN
1	С	25	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tune	Chain	Res	Link	Bo	ond leng	ths	B	ond ang	jles
	Type	Cham	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	А	2647	-	5, 5, 5	0.45	0	$5,\!5,\!5$	0.48	0
2	PNM	А	300	1	17,24,24	1.14	1(5%)	22,34,34	1.44	2 (9%)
2	PNM	В	300	1	17,24,24	1.22	1 (5%)	22,34,34	1.15	2 (9%)
2	PNM	С	300	1	17,24,24	1.44	2 (11%)	22,34,34	0.95	0
2	PNM	D	300	1	17,24,24	1.29	2 (11%)	22,34,34	1.78	6 (27%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	А	2647	-	-	3/4/4/4	-
2	PNM	А	300	1	-	4/8/33/33	0/2/2/2
2	PNM	В	300	1	-	2/8/33/33	0/2/2/2
2	PNM	С	300	1	-	4/8/33/33	0/2/2/2
2	PNM	D	300	1	-	4/8/33/33	0/2/2/2

'-' means no outliers of that kind were identified.

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	С	300	PNM	C17-C18	4.19	1.58	1.51
2	D	300	PNM	C17-C18	2.99	1.56	1.51
2	В	300	PNM	C17-C18	2.98	1.56	1.51
2	D	300	PNM	C17-C15	2.93	1.58	1.51
2	А	300	PNM	C17-C18	2.80	1.56	1.51

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	D	300	PNM	C18-C17-C15	4.35	125.47	112.57
2	А	300	PNM	C6-N14-C15	-4.16	118.67	123.13
2	D	300	PNM	C9-C2-S1	3.63	115.29	109.21
2	А	300	PNM	C3-C2-S1	-3.16	97.54	103.81
2	В	300	PNM	C3-C2-S1	-3.01	97.83	103.81

There are no chirality outliers.

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	2647	GOL	C1-C2-C3-O3
3	А	2647	GOL	O2-C2-C3-O3
2	С	300	PNM	C15-C17-C18-C19
2	D	300	PNM	C15-C17-C18-C19
2	А	300	PNM	C15-C17-C18-C19

There are no ring outliers.

4 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	300	PNM	5	0




	Chain	1	1 0		Symm-Clashes
2	В	300	PNM	2	0
2	С	300	PNM	13	0
2	D	300	PNM	9	0


Continued from previous page...

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



















5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	$\mathbf{Q}{<}0.9$
1	А	264/298~(88%)	-0.18	3 (1%) 80 83	16, 34, 59, 69	0
1	В	265/298~(88%)	-0.42	0 100 100	16, 26, 41, 58	0
1	С	264/298~(88%)	-0.33	4 (1%) 73 77	12, 22, 44, 54	0
1	D	274/298~(91%)	-0.28	3 (1%) 80 83	16, 28, 45, 59	0
All	All	1067/1192~(89%)	-0.30	10 (0%) 84 87	12, 27, 50, 69	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	\mathbf{Res}	Type	RSRZ
1	С	225	ILE	4.0
1	D	2	PRO	3.7
1	D	274	LEU	3.5
1	С	274	LEU	3.4
1	С	30	ARG	3.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
2	PNM	С	300	23/23	0.62	0.40	$33,\!58,\!67,\!68$	0
3	GOL	А	2647	6/6	0.65	0.22	56, 57, 58, 60	0
2	PNM	В	300	23/23	0.68	0.32	$32,\!53,\!65,\!66$	0
2	PNM	D	300	23/23	0.69	0.41	$37,\!55,\!68,\!69$	0
2	PNM	А	300	23/23	0.71	0.44	48,62,74,76	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















6.5 Other polymers (i)

There are no such residues in this entry.

