



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 21, 2024 – 11:12 pm BST

PDB ID : 2JE1  
Title : The crystal Structure of the tumor suppressor protein pp32 (Anp32a) :structural insights into the Anp32 family of proteins  
Authors : Huyton, T.; Wolberger, C.  
Deposited on : 2007-01-12  
Resolution : 2.69 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	80% 15% . .
1	B	149	82% 15% . .
1	C	149	79% 15% 5% . .
1	D	149	82% 16% . .

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 4907 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACIDIC LEUCINE-RICH NUCLEAR PHOSPHOPROTEIN 32 FAMILY MEMBER A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	149	Total 1189	C 746	N 203	O 236	S 4	11	0	0
1	B	149	Total 1189	C 746	N 203	O 236	S 4	11	0	0
1	C	149	Total 1189	C 746	N 203	O 236	S 4	15	0	0
1	D	149	Total 1189	C 746	N 203	O 236	S 4	12	0	0

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	Total 6	C 3	O 3	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	34	Total O 34 34	0	0
3	B	34	Total O 34 34	0	0
3	C	34	Total O 34 34	0	0
3	D	43	Total O 43 43	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ACIDIC LEUCINE-RICH NUCLEAR PHOSPHOPROTEIN 32 FAMILY MEMBER A

Chain A:  80% 15% ..





- Molecule 1: ACIDIC LEUCINE-RICH NUCLEAR PHOSPHOPROTEIN 32 FAMILY MEMBER A

Chain B:  82% 15% ..




- Molecule 1: ACIDIC LEUCINE-RICH NUCLEAR PHOSPHOPROTEIN 32 FAMILY MEMBER A

Chain C:  79% 15% 5% ..



- Molecule 1: ACIDIC LEUCINE-RICH NUCLEAR PHOSPHOPROTEIN 32 FAMILY MEMBER A

Chain D:  82% 16% ..



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	106.51Å 106.51Å 133.15Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	25.00 – 2.69 24.08 – 2.69	Depositor EDS
% Data completeness (in resolution range)	97.7 (25.00-2.69) 97.7 (24.08-2.69)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.68Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.238 , 0.303 0.234 , 0.290	Depositor DCC
$R_{free}$ test set	1189 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.7	Xtrriage
Anisotropy	0.078	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 22.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.458 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	4907	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 51.99 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.1877e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	8.17	9/1202 (0.7%)	4.20	12/1619 (0.7%)
1	B	7.04	5/1202 (0.4%)	2.95	10/1619 (0.6%)
1	C	9.03	4/1202 (0.3%)	4.07	10/1619 (0.6%)
1	D	3.88	5/1202 (0.4%)	5.62	7/1619 (0.4%)
All	All	7.30	23/4808 (0.5%)	4.31	39/6476 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	3
1	D	0	2
All	All	0	9

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	28	ARG	CZ-NH2	252.47	4.61	1.33
1	C	28	ARG	CZ-NH2	209.82	4.05	1.33
1	B	5	ARG	CZ-NH2	180.08	3.67	1.33
1	C	75	ARG	CZ-NH2	172.28	3.57	1.33
1	C	75	ARG	CZ-NH1	151.67	3.30	1.33
1	B	5	ARG	CZ-NH1	121.15	2.90	1.33
1	B	28	ARG	CZ-NH1	98.89	2.61	1.33
1	A	28	ARG	CZ-NH1	83.41	2.41	1.33
1	D	5	ARG	CZ-NH1	75.54	2.31	1.33
1	D	28	ARG	CZ-NH2	71.08	2.25	1.33

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	5	ARG	CZ-NH2	68.60	2.22	1.33
1	A	5	ARG	CZ-NH1	63.07	2.15	1.33
1	A	75	ARG	CZ-NH2	54.12	2.03	1.33
1	B	28	ARG	CZ-NH2	-48.22	0.70	1.33
1	D	28	ARG	CZ-NH1	-46.32	0.72	1.33
1	A	5	ARG	CZ-NH2	-36.69	0.85	1.33
1	C	28	ARG	CZ-NH1	29.98	1.72	1.33
1	A	75	ARG	CZ-NH1	-26.39	0.98	1.33
1	D	5	ARG	CB-CG	-14.16	1.14	1.52
1	A	62	LYS	CE-NZ	10.68	1.75	1.49
1	A	75	ARG	CG-CD	-8.96	1.29	1.51
1	A	28	ARG	CB-CG	7.06	1.71	1.52
1	B	28	ARG	CG-CD	-6.35	1.36	1.51

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	5	ARG	NE-CZ-NH1	-199.73	20.43	120.30
1	C	75	ARG	NE-CZ-NH2	-121.21	59.70	120.30
1	C	75	ARG	NE-CZ-NH1	97.66	169.13	120.30
1	A	28	ARG	NE-CZ-NH1	-96.73	71.93	120.30
1	A	5	ARG	NE-CZ-NH2	-91.50	74.55	120.30
1	B	5	ARG	NE-CZ-NH2	-72.93	83.84	120.30
1	A	75	ARG	NE-CZ-NH2	-68.86	85.87	120.30
1	B	28	ARG	NE-CZ-NH1	-64.99	87.80	120.30
1	D	28	ARG	NE-CZ-NH1	64.45	152.52	120.30
1	D	28	ARG	NE-CZ-NH2	-55.80	92.40	120.30
1	A	28	ARG	NH1-CZ-NH2	50.02	174.42	119.40
1	B	28	ARG	NE-CZ-NH2	45.27	142.94	120.30
1	D	5	ARG	NE-CZ-NH2	-43.19	98.71	120.30
1	C	28	ARG	NE-CZ-NH1	-34.03	103.28	120.30
1	B	5	ARG	NH1-CZ-NH2	-33.17	82.91	119.40
1	D	28	ARG	NH1-CZ-NH2	-28.37	88.19	119.40
1	A	5	ARG	NH1-CZ-NH2	26.51	148.56	119.40
1	A	5	ARG	NE-CZ-NH1	25.33	132.97	120.30
1	A	75	ARG	NH1-CZ-NH2	23.76	145.53	119.40
1	A	75	ARG	NE-CZ-NH1	-20.88	109.86	120.30
1	D	5	ARG	NH1-CZ-NH2	-20.16	97.22	119.40
1	B	28	ARG	CG-CD-NE	15.21	143.74	111.80
1	A	28	ARG	NE-CZ-NH2	-13.50	113.55	120.30
1	B	28	ARG	CB-CG-CD	12.13	143.15	111.60
1	A	75	ARG	CB-CG-CD	11.19	140.68	111.60

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	28	ARG	NE-CZ-NH2	-10.94	114.83	120.30
1	B	5	ARG	NE-CZ-NH1	10.44	125.52	120.30
1	C	28	ARG	NH1-CZ-NH2	10.18	130.60	119.40
1	C	33	LYS	CB-CG-CD	9.73	136.90	111.60
1	C	75	ARG	NH1-CZ-NH2	-8.79	109.74	119.40
1	A	28	ARG	CA-CB-CG	-6.67	98.72	113.40
1	B	9	LEU	CA-CB-CG	6.56	130.39	115.30
1	C	142	LEU	CA-CB-CG	6.42	130.07	115.30
1	D	142	LEU	CA-CB-CG	6.28	129.75	115.30
1	B	28	ARG	NH1-CZ-NH2	-6.24	112.53	119.40
1	B	142	LEU	CA-CB-CG	5.82	128.69	115.30
1	A	142	LEU	CA-CB-CG	5.80	128.63	115.30
1	C	9	LEU	CA-CB-CG	5.12	127.08	115.30
1	C	60	LEU	CA-CB-CG	5.11	127.05	115.30

There are no chirality outliers.

All (9) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	28	ARG	Sidechain
1	A	75	ARG	Sidechain
1	B	28	ARG	Sidechain
1	B	5	ARG	Sidechain
1	C	148	TYR	Peptide
1	C	28	ARG	Sidechain
1	C	75	ARG	Sidechain
1	D	28	ARG	Sidechain
1	D	5	ARG	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1189	0	1220	15	0
1	B	1189	0	1220	9	5
1	C	1189	0	1220	12	0
1	D	1189	0	1220	11	5

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	6	0	8	3	0
3	A	34	0	0	0	0
3	B	34	0	0	0	0
3	C	34	0	0	2	0
3	D	43	0	0	2	0
All	All	4907	0	4888	43	5

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:LYS:CE	1:A:62:LYS:NZ	1.75	1.49
1:A:102:ASP:CB	2:A:1150:GOL:H2	1.89	1.02
1:A:102:ASP:HB2	2:A:1150:GOL:H2	1.00	1.00
1:A:102:ASP:HB2	2:A:1150:GOL:C2	1.94	0.97
1:D:28:ARG:NH1	1:D:28:ARG:NH2	2.34	0.75
1:A:123:CYS:O	1:A:126:THR:HB	1.91	0.70
1:D:12:ARG:HG2	3:D:2004:HOH:O	1.92	0.70
1:C:102:ASP:OD2	1:C:105:THR:HG22	1.92	0.69
1:D:3:MET:HE2	1:D:35:GLU:O	1.92	0.68
1:B:102:ASP:OD2	1:B:105:THR:HG23	1.94	0.67
1:C:123:CYS:O	1:C:126:THR:HB	1.95	0.66
1:A:148:TYR:O	1:A:149:ASP:HB2	1.97	0.64
1:D:46:PHE:CD1	1:D:68:LYS:HG2	2.33	0.63
1:D:50:ILE:HG12	1:D:72:SER:HB2	1.83	0.58
1:A:62:LYS:NZ	1:A:62:LYS:CD	2.65	0.57
1:B:2:GLU:H	1:B:5:ARG:HH21	1.55	0.53
1:D:123:CYS:O	1:D:126:THR:HB	2.08	0.53
1:C:148:TYR:O	1:C:149:ASP:HB2	2.08	0.53
1:C:126:THR:HG21	3:C:2022:HOH:O	2.08	0.52
1:D:148:TYR:O	1:D:149:ASP:HB2	2.09	0.52
1:C:110:LYS:HD2	3:C:2030:HOH:O	2.09	0.52
1:D:107:GLU:HG2	3:D:2031:HOH:O	2.10	0.51
1:D:46:PHE:HD1	1:D:68:LYS:HG2	1.75	0.50
1:A:12:ARG:CZ	1:A:12:ARG:HB2	2.41	0.49
1:A:14:ARG:NH2	1:B:21:GLU:OE2	2.46	0.48
1:B:3:MET:HE2	1:B:35:GLU:O	2.14	0.48
1:C:60:LEU:HD22	1:C:61:PRO:HD2	1.95	0.47
1:B:148:TYR:O	1:B:149:ASP:HB2	2.14	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:102:ASP:OD2	1:A:105:THR:HG23	2.15	0.46
1:B:11:LEU:O	1:B:14:ARG:HD3	2.15	0.46
1:C:21:GLU:OE1	1:D:14:ARG:NH2	2.48	0.46
1:C:14:ARG:NH2	1:D:21:GLU:OE1	2.48	0.46
1:C:50:ILE:HG12	1:C:72:SER:HB2	1.98	0.46
1:A:107:GLU:O	1:A:110:LYS:HG2	2.15	0.45
1:A:69:LEU:HB2	1:A:90:LEU:HD11	1.99	0.44
1:C:32:GLY:HA2	1:C:53:GLY:O	2.18	0.44
1:B:21:GLU:HG2	1:B:46:PHE:HB3	1.99	0.43
1:A:21:GLU:OE2	1:B:14:ARG:NH2	2.52	0.43
1:A:32:GLY:HA2	1:A:53:GLY:O	2.19	0.43
1:B:107:GLU:O	1:B:110:LYS:HG2	2.20	0.42
1:C:21:GLU:HG3	1:C:46:PHE:HD2	1.85	0.41
1:A:62:LYS:HE2	1:A:86:LYS:NZ	2.36	0.40
1:C:12:ARG:CZ	1:C:12:ARG:HB2	2.46	0.40

All (5) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:28:ARG:NH1	1:D:28:ARG:NE[6_555]	1.51	0.69
1:B:28:ARG:NH1	1:D:28:ARG:CG[6_555]	1.57	0.63
1:B:28:ARG:NH1	1:D:28:ARG:CD[6_555]	1.62	0.58
1:B:28:ARG:CG	1:D:28:ARG:CD[6_555]	1.82	0.38
1:B:28:ARG:NH1	1:D:28:ARG:CZ[6_555]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	147/149 (99%)	140 (95%)	7 (5%)	0	<b>100</b> <b>100</b>

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	147/149 (99%)	140 (95%)	7 (5%)	0	100	100
1	C	147/149 (99%)	139 (95%)	8 (5%)	0	100	100
1	D	147/149 (99%)	141 (96%)	6 (4%)	0	100	100
All	All	588/596 (99%)	560 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	139/139 (100%)	125 (90%)	14 (10%)	7	17
1	B	139/139 (100%)	125 (90%)	14 (10%)	7	17
1	C	139/139 (100%)	123 (88%)	16 (12%)	5	13
1	D	139/139 (100%)	127 (91%)	12 (9%)	10	24
All	All	556/556 (100%)	500 (90%)	56 (10%)	7	17

All (56) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	5	ARG
1	A	15	THR
1	A	22	LEU
1	A	33	LYS
1	A	60	LEU
1	A	62	LYS
1	A	69	LEU
1	A	104	SER
1	A	110	LYS
1	A	124	GLU
1	A	128	LEU
1	A	130	ASP

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>
1	A	149	ASP
1	B	1	MET
1	B	22	LEU
1	B	28	ARG
1	B	54	LEU
1	B	60	LEU
1	B	64	ASN
1	B	69	LEU
1	B	104	SER
1	B	107	GLU
1	B	110	LYS
1	B	124	GLU
1	B	128	LEU
1	B	143	THR
1	B	149	ASP
1	C	1	MET
1	C	9	LEU
1	C	11	LEU
1	C	12	ARG
1	C	15	THR
1	C	60	LEU
1	C	69	LEU
1	C	104	SER
1	C	105	THR
1	C	107	GLU
1	C	110	LYS
1	C	128	LEU
1	C	130	ASP
1	C	131	TYR
1	C	143	THR
1	C	149	ASP
1	D	1	MET
1	D	5	ARG
1	D	9	LEU
1	D	11	LEU
1	D	15	THR
1	D	54	LEU
1	D	60	LEU
1	D	69	LEU
1	D	110	LYS
1	D	124	GLU
1	D	128	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	149	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	26	ASN
1	C	26	ASN
1	C	51	ASN
1	C	89	ASN
1	D	26	ASN
1	D	51	ASN
1	D	89	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	A	1150	-	5,5,5	0.48	0	5,5,5	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1150	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1150	GOL	O1-C1-C2-C3
2	A	1150	GOL	O1-C1-C2-O2
2	A	1150	GOL	O2-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1150	GOL	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	149/149 (100%)	-0.16	0 100 100	28, 31, 34, 38	3 (2%)
1	B	149/149 (100%)	-0.18	0 100 100	29, 31, 34, 39	3 (2%)
1	C	149/149 (100%)	-0.24	0 100 100	28, 31, 34, 39	4 (2%)
1	D	149/149 (100%)	-0.15	0 100 100	28, 31, 34, 39	3 (2%)
All	All	596/596 (100%)	-0.18	0 100 100	28, 31, 34, 39	13 (2%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	A	1150	6/6	0.90	0.30	60,63,63,63	0



## 6.5 Other polymers [i](#)

There are no such residues in this entry.