



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 29, 2024 – 02:12 PM JST

PDB ID : 8JKK  
Title : Crystal Structure of the dioxygenase CcTet from Coprinopsis cinerea bound to 12bp 5-methylcytosine (5mC) containing duplex DNA  
Authors : Zhang, L.; Zhang, L.  
Deposited on : 2023-06-01  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	415	
1	D	415	
1	G	415	
1	J	415	
2	B	12	
2	E	12	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	H	12	 58% 42%
2	K	12	 33% 50% 33% 17%
3	C	12	 42% 58%
3	F	12	 8% 25% 67% 8%
3	I	12	 17% 67% 17%
3	L	12	 17% 42% 58% 17% 8%

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 14841 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2OGFeDO JBP1/TET oxygenase domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	376	Total 2955	C 1889	N 523	O 529	S 14	0	0	0
1	D	383	Total 3007	C 1921	N 535	O 536	S 15	0	0	0
1	G	376	Total 2947	C 1886	N 520	O 527	S 14	0	0	0
1	J	383	Total 3007	C 1921	N 533	O 538	S 15	0	0	0

- Molecule 2 is a DNA chain called DNA (5'-D(P\*CP\*GP\*AP\*TP\*CP\*(5CM)P\*GP\*CP\*TP\*AP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	B	12	Total 244	C 116	N 44	O 72	P 12	0	0	0
2	E	12	Total 244	C 116	N 44	O 72	P 12	0	0	0
2	H	12	Total 244	C 116	N 44	O 72	P 12	0	0	0
2	K	12	Total 244	C 116	N 44	O 72	P 12	0	0	0

- Molecule 3 is a DNA chain called DNA (5'-D(P\*CP\*GP\*TP\*AP\*GP\*CP\*TP\*GP\*AP\*TP\*CP\*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	12	Total 247	C 117	N 45	O 73	P 12	0	0	0
3	F	12	Total 247	C 117	N 45	O 73	P 12	0	0	0
3	I	12	Total 247	C 117	N 45	O 73	P 12	0	0	0

*Continued on next page...*


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	L	11	Total 228	C 108	N 42	O 67	P 11	0	0	0

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Mn 1	0	0
4	D	1	Total 1	Mn 1	0	0
4	G	1	Total 1	Mn 1	0	0
4	J	1	Total 1	Mn 1	0	0

- Molecule 5 is N-OXALYLGLYCINE (three-letter code: OGA) (formula: C<sub>4</sub>H<sub>5</sub>NO<sub>5</sub>).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	Total 10	C 4	N 1	O 5	0	0
5	D	1	Total 10	C 4	N 1	O 5	0	0
5	G	1	Total 10	C 4	N 1	O 5	0	0
5	J	1	Total 10	C 4	N 1	O 5	0	0



- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	298	Total O 298 298	0	0
6	B	16	Total O 16 16	0	0
6	C	9	Total O 9 9	0	0
6	D	275	Total O 275 275	0	0
6	E	21	Total O 21 21	0	0
6	F	23	Total O 23 23	0	0
6	G	188	Total O 188 188	0	0
6	H	22	Total O 22 22	0	0
6	I	10	Total O 10 10	0	0
6	J	70	Total O 70 70	0	0
6	K	3	Total O 3 3	0	0
6	L	1	Total O 1 1	0	0






- Molecule 1: 2OGFeDO JBP1/TET oxygenase domain-containing protein




- Molecule 2: DNA (5'-D(P\*CP\*GP\*AP\*TP\*CP\*(5CM)P\*GP\*CP\*TP\*AP\*CP\*G)-3')




- Molecule 2: DNA (5'-D(P\*CP\*GP\*AP\*TP\*CP\*(5CM)P\*GP\*CP\*TP\*AP\*CP\*G)-3')



- Molecule 2: DNA (5'-D(P\*CP\*GP\*AP\*TP\*CP\*(5CM)P\*GP\*CP\*TP\*AP\*CP\*G)-3')



- Molecule 2: DNA (5'-D(P\*CP\*GP\*AP\*TP\*CP\*(5CM)P\*GP\*CP\*TP\*AP\*CP\*G)-3')





- Molecule 3: DNA (5'-D(P\*CP\*GP\*TP\*AP\*GP\*CP\*TP\*GP\*AP\*TP\*CP\*G)-3')

Chain C: 




- Molecule 3: DNA (5'-D(P\*CP\*GP\*TP\*AP\*GP\*CP\*TP\*GP\*AP\*TP\*CP\*G)-3')

Chain F: 





- Molecule 3: DNA (5'-D(P\*CP\*GP\*TP\*AP\*GP\*CP\*TP\*GP\*AP\*TP\*CP\*G)-3')

Chain I: 



- Molecule 3: DNA (5'-D(P\*CP\*GP\*TP\*AP\*GP\*CP\*TP\*GP\*AP\*TP\*CP\*G)-3')

Chain L: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	86.57Å 127.63Å 212.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	21.98 – 2.30 49.93 – 2.30	Depositor EDS
% Data completeness (in resolution range)	90.5 (21.98-2.30) 90.5 (49.93-2.30)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	5.98 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, $R_{free}$	0.187 , 0.216 0.189 , 0.216	Depositor DCC
$R_{free}$ test set	4770 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.5	Xtrriage
Anisotropy	0.200	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	14841	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.95% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: OGA, 5CM, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.63	6/3033 (0.2%)	0.76	7/4128 (0.2%)
1	D	1.13	38/3084 (1.2%)	0.84	3/4194 (0.1%)
1	G	0.63	6/3025 (0.2%)	0.77	6/4118 (0.1%)
1	J	0.69	7/3085 (0.2%)	0.92	19/4197 (0.5%)
2	B	0.90	0/249	1.38	3/379 (0.8%)
2	E	1.75	5/249 (2.0%)	1.51	4/379 (1.1%)
2	H	0.92	0/249	1.19	0/379
2	K	0.87	0/249	1.39	4/379 (1.1%)
3	C	0.98	0/276	1.07	0/424
3	F	1.09	0/276	1.21	1/424 (0.2%)
3	I	1.26	3/276 (1.1%)	1.31	2/424 (0.5%)
3	L	1.02	1/255 (0.4%)	1.19	3/392 (0.8%)
All	All	0.86	66/14306 (0.5%)	0.91	52/19817 (0.3%)

The worst 5 of 66 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	5	DC	P-OP2	-11.96	1.28	1.49
1	J	248	PRO	N-CA	11.47	1.66	1.47
1	D	251	GLU	CD-OE1	-9.46	1.15	1.25
1	G	395	ARG	CZ-NH1	8.79	1.44	1.33
1	D	251	GLU	CD-OE2	-8.60	1.16	1.25

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	106	LYS	CD-CE-NZ	15.62	147.62	111.70
1	J	32	ARG	NE-CZ-NH1	-11.84	114.38	120.30
2	B	1	DC	OP1-P-O3'	-11.02	80.95	105.20
2	K	1	DC	OP1-P-O3'	-10.78	81.49	105.20
1	J	32	ARG	CA-CB-CG	10.49	136.49	113.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2955	0	2964	28	0
1	D	3007	0	3029	27	0
1	G	2947	0	2959	33	0
1	J	3007	0	3020	34	0
2	B	244	0	137	7	0
2	E	244	0	137	5	0
2	H	244	0	137	3	0
2	K	244	0	137	5	0
3	C	247	0	136	6	0
3	F	247	0	136	6	0
3	I	247	0	136	7	0
3	L	228	0	125	5	0
4	A	1	0	0	0	0
4	D	1	0	0	0	0
4	G	1	0	0	0	0
4	J	1	0	0	0	0
5	A	10	0	3	0	0
5	D	10	0	3	0	0
5	G	10	0	3	0	0
5	J	10	0	3	0	0
6	A	298	0	0	6	0
6	B	16	0	0	2	0
6	C	9	0	0	0	0
6	D	275	0	0	2	0
6	E	21	0	0	0	0
6	F	23	0	0	0	0
6	G	188	0	0	7	0
6	H	22	0	0	0	0
6	I	10	0	0	0	0
6	J	70	0	0	1	0
6	K	3	0	0	1	0
6	L	1	0	0	0	0
All	All	14841	0	13065	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:57:LEU:HB2	1:D:261:ILE:HD11	1.31	1.06
1:D:57:LEU:CB	1:D:261:ILE:HD11	2.01	0.90
1:A:178:ARG:HG2	1:A:208:LEU:HD21	1.61	0.80
1:D:57:LEU:HB2	1:D:261:ILE:CD1	2.14	0.75
1:J:123:ILE:HG13	1:J:124:LEU:HD22	1.76	0.68

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	372/415 (90%)	360 (97%)	11 (3%)	1 (0%)	41	50
1	D	379/415 (91%)	368 (97%)	10 (3%)	1 (0%)	41	50
1	G	372/415 (90%)	361 (97%)	10 (3%)	1 (0%)	41	50
1	J	379/415 (91%)	367 (97%)	11 (3%)	1 (0%)	41	50
All	All	1502/1660 (90%)	1456 (97%)	42 (3%)	4 (0%)	41	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	259	ILE
1	G	259	ILE
1	D	259	ILE
1	J	259	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	315/348 (90%)	308 (98%)	7 (2%)	52 69
1	D	320/348 (92%)	311 (97%)	9 (3%)	43 60
1	G	314/348 (90%)	307 (98%)	7 (2%)	52 69
1	J	320/348 (92%)	310 (97%)	10 (3%)	40 55
All	All	1269/1392 (91%)	1236 (97%)	33 (3%)	46 63

5 of 33 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	J	180	ARG
1	J	203	ASN
1	J	411	GLN
1	D	229	TRP
1	D	225	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	115	ASN
1	G	168	GLN
1	J	35	GLN
1	D	237	GLN
1	A	35	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	5CM	B	6	2	17,21,22	2.12	8 (47%)	24,30,33	2.42	11 (45%)
2	5CM	E	6	2	17,21,22	3.03	8 (47%)	24,30,33	2.50	10 (41%)
2	5CM	H	6	2	17,21,22	1.90	6 (35%)	24,30,33	2.36	11 (45%)
2	5CM	K	6	2	17,21,22	2.09	7 (41%)	24,30,33	2.39	10 (41%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	5CM	B	6	2	-	6/7/21/22	0/2/2/2
2	5CM	E	6	2	-	6/7/21/22	0/2/2/2
2	5CM	H	6	2	-	6/7/21/22	0/2/2/2
2	5CM	K	6	2	-	6/7/21/22	0/2/2/2

The worst 5 of 29 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	6	5CM	C6-N1	-6.27	1.27	1.38
2	E	6	5CM	O2-C2	-5.83	1.12	1.23
2	B	6	5CM	C4-N3	-4.05	1.27	1.34
2	E	6	5CM	C2-N3	-3.94	1.28	1.36
2	E	6	5CM	C4-N3	-3.82	1.27	1.34

The worst 5 of 42 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	6	5CM	C1'-N1-C2	5.38	127.18	117.74
2	B	6	5CM	O3'-C3'-C2'	-5.35	91.76	110.90
2	K	6	5CM	O2-C2-N3	-5.10	114.03	122.33
2	B	6	5CM	O2-C2-N3	-5.05	114.12	122.33
2	K	6	5CM	O3'-C3'-C2'	-5.01	92.98	110.90

There are no chirality outliers.

5 of 24 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	6	5CM	C3'-C4'-C5'-O5'
2	H	6	5CM	O4'-C4'-C5'-O5'
2	B	6	5CM	C3'-C4'-C5'-O5'
2	B	6	5CM	O4'-C4'-C5'-O5'
2	E	6	5CM	C3'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	6	5CM	2	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	OGA	G	502	4	9,9,9	1.14	1 (11%)	10,11,11	1.61	2 (20%)
5	OGA	D	502	4	9,9,9	1.11	0	10,11,11	1.51	0
5	OGA	J	502	4	9,9,9	1.22	0	10,11,11	1.50	1 (10%)
5	OGA	A	502	4	9,9,9	1.26	0	10,11,11	1.47	2 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	OGA	G	502	4	-	0/8/9/9	-
5	OGA	D	502	4	-	1/8/9/9	-
5	OGA	J	502	4	-	1/8/9/9	-
5	OGA	A	502	4	-	0/8/9/9	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	G	502	OGA	O3-C5	-2.14	1.23	1.30

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	502	OGA	O2-C1-C2	3.10	122.29	113.15
5	J	502	OGA	O2-C1-C2	2.63	120.89	113.15
5	A	502	OGA	O2-C1-C2	2.28	119.88	113.15
5	A	502	OGA	O2'-C2-C1	-2.21	118.20	121.32
5	G	502	OGA	O1-C1-C2	-2.03	117.66	122.18

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	D	502	OGA	N1-C4-C5-O4
5	J	502	OGA	N1-C4-C5-O4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	376/415 (90%)	0.38	28 (7%) 14 19	18, 28, 51, 89	0
1	D	383/415 (92%)	0.12	7 (1%) 68 74	19, 29, 50, 79	0
1	G	376/415 (90%)	0.11	14 (3%) 41 48	24, 38, 56, 94	0
1	J	383/415 (92%)	0.54	41 (10%) 6 8	36, 51, 76, 105	0
2	B	11/12 (91%)	0.09	1 (9%) 9 12	34, 42, 50, 64	0
2	E	11/12 (91%)	0.33	1 (9%) 9 12	30, 33, 65, 73	0
2	H	11/12 (91%)	-0.22	0 100 100	33, 37, 53, 59	0
2	K	11/12 (91%)	2.01	4 (36%) 0 0	54, 61, 108, 111	0
3	C	12/12 (100%)	0.05	0 100 100	39, 46, 49, 50	0
3	F	12/12 (100%)	0.18	1 (8%) 11 15	28, 45, 70, 81	0
3	I	12/12 (100%)	-0.09	0 100 100	44, 47, 49, 51	0
3	L	11/12 (91%)	1.72	5 (45%) 0 0	57, 62, 119, 123	0
All	All	1609/1756 (91%)	0.30	102 (6%) 20 25	18, 38, 65, 123	0

The worst 5 of 102 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	118	HIS	6.5
1	A	178	ARG	6.4
1	J	181	VAL	5.7
1	J	117	ASP	5.5
1	J	172	ARG	5.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	5CM	B	6	20/21	0.91	0.17	25,32,50,52	0
2	5CM	H	6	20/21	0.91	0.15	29,37,48,54	0
2	5CM	K	6	20/21	0.92	0.12	39,45,57,63	0
2	5CM	E	6	20/21	0.96	0.12	20,28,36,39	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	OGA	J	502	10/10	0.94	0.13	36,41,44,44	0
5	OGA	D	502	10/10	0.98	0.10	17,20,26,38	0
5	OGA	A	502	10/10	0.98	0.13	20,23,25,26	0
5	OGA	G	502	10/10	0.99	0.10	28,30,32,33	0
4	MN	J	501	1/1	0.99	0.11	37,37,37,37	0
4	MN	G	501	1/1	1.00	0.10	33,33,33,33	0
4	MN	A	501	1/1	1.00	0.11	23,23,23,23	0
4	MN	D	501	1/1	1.00	0.14	21,21,21,21	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.