



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 4, 2024 – 09:26 am GMT

PDB ID : 5JMY
Title : NEPRILYSIN COMPLEXED WITH LBQ657
Authors : Schiering, N.; Wiesmann, C.
Deposited on : 2016-04-29
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	698	 2% 89% 10%
1	B	698	 5% 88% 11%
2	C	2	 50% 50%
2	D	2	 50% 50%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 12295 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Neprilysin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	696	5612	3549	961	1076	26	0	2	0
1	B	696	5595	3538	957	1074	26	0	0	0

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	2	28	16	2	10	0	0	0
2	D	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0
3	B	1	14	8	1	5	0	0
3	B	1	14	8	1	5	0	0
3	B	1	14	8	1	5	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
4	A	1	1	1	0	0
4	B	1	1	1	0	0

- Molecule 5 is Sacubitrilat (three-letter code: 6LD) (formula: C₂₂H₂₅NO₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
5	A	1	28	22	1	5	0	0
5	B	1	28	22	1	5	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
6	A	514	514	514	0	0
6	B	376	376	376	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Neprilysin



- Molecule 1: Neprilysin



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



MAG1
MAG2

- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 

MAG1
MAG2

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	59.74Å 109.14Å 248.00Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.28 – 2.00 29.08 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.2 (29.28-2.00) 98.2 (29.08-2.00)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.11 (at 2.00Å)	Xtrriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.186 , 0.228 0.201 , 0.246	Depositor DCC
R_{free} test set	4992 reflections (4.60%)	wwPDB-VP
Wilson B-factor (Å ²)	32.0	Xtrriage
Anisotropy	0.053	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12295	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 6LD, ZN, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/5731	0.65	0/7752
1	B	0.50	0/5713	0.65	0/7727
All	All	0.51	0/11444	0.65	0/15479

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5612	0	5460	33	0
1	B	5595	0	5446	36	0
2	C	28	0	25	0	0
2	D	28	0	25	0	0
3	A	42	0	39	0	0
3	B	42	0	39	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	28	0	0	0	0
5	B	28	0	0	0	0
6	A	514	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	376	0	0	1	0
All	All	12295	0	11034	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:202:LEU:HD23	1:A:218:ILE:HD11	1.70	0.72
1:B:202:LEU:HD23	1:B:218:ILE:HD11	1.72	0.71
1:A:305:ALA:HB2	1:A:339:ASN:HB3	1.77	0.67
1:B:168:VAL:H	1:B:368:ASN:HD21	1.44	0.65
1:B:183:ALA:HB2	1:B:315:ILE:HG21	1.79	0.64
1:A:168:VAL:H	1:A:368:ASN:HD21	1.43	0.64
1:A:333:VAL:HG11	1:A:523:LYS:HB3	1.79	0.62
1:B:333:VAL:HG11	1:B:523:LYS:HB3	1.79	0.62
1:A:67:ARG:HH22	1:A:688:ASN:HD21	1.47	0.62
1:A:407:TRP:CZ2	1:B:95:VAL:HG11	2.34	0.61
1:B:502:LYS:H	1:B:509:ASN:HD21	1.50	0.59
1:A:502:LYS:H	1:A:509:ASN:HD21	1.52	0.58
1:A:67:ARG:HH22	1:A:688:ASN:ND2	2.03	0.56
1:B:589:PHE:HB3	1:B:749:TRP:CZ2	2.41	0.56
1:A:589:PHE:HB3	1:A:749:TRP:CZ2	2.41	0.56
1:A:96:ILE:HD11	1:A:696:THR:HG23	1.90	0.53
1:B:67:ARG:HH22	1:B:688:ASN:ND2	2.07	0.52
1:B:121:LEU:HD13	1:B:414:VAL:HG21	1.92	0.52
1:B:67:ARG:HH22	1:B:688:ASN:HD21	1.57	0.51
1:A:67:ARG:HH12	1:A:688:ASN:ND2	2.09	0.51
1:B:67:ARG:HH12	1:B:688:ASN:ND2	2.09	0.51
1:A:62:ILE:HG13	1:B:62:ILE:HG13	1.94	0.50
1:A:278:LEU:HD11	1:A:366:LEU:HD23	1.94	0.50
1:A:535:ILE:HD13	1:A:546:SER:HB2	1.94	0.50
1:B:67:ARG:HH12	1:B:688:ASN:HD22	1.60	0.50
1:A:597:ASN:ND2	1:A:599:ASP:H	2.10	0.50
1:B:570:ASN:O	1:B:574:TYR:HD1	1.94	0.49
1:B:597:ASN:ND2	1:B:599:ASP:H	2.11	0.49
1:A:372:TRP:CE2	1:A:376:MET:HG3	2.48	0.48
1:B:372:TRP:CE2	1:B:376:MET:HG3	2.48	0.48
1:A:347:ALA:HB1	1:A:350:TYR:HB3	1.95	0.48
1:B:278:LEU:HD11	1:B:366:LEU:HD23	1.93	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96:ILE:HD11	1:B:696:THR:HG23	1.95	0.48
1:A:121:LEU:HD13	1:A:414[A]:VAL:HG21	1.96	0.48
1:A:570:ASN:O	1:A:574:TYR:HD1	1.96	0.48
1:B:464:THR:HG23	1:B:607:THR:HG23	1.96	0.47
1:A:68:LEU:HD22	1:A:687:LEU:HD13	1.96	0.47
1:A:460:MET:HE3	1:A:465:LYS:HA	1.96	0.47
1:A:67:ARG:HH12	1:A:688:ASN:HD22	1.63	0.47
1:B:535:ILE:HD13	1:B:546:SER:HB2	1.97	0.47
1:A:597:ASN:HD22	1:A:597:ASN:C	2.18	0.46
1:B:314:GLU:HB3	1:B:359:THR:HG22	1.97	0.46
1:B:597:ASN:C	1:B:597:ASN:HD22	2.18	0.46
1:A:90:TRP:CZ2	1:A:104:GLY:HA2	2.51	0.45
1:B:315:ILE:HD11	1:B:358:LEU:HD23	1.98	0.45
1:B:715:ASN:O	1:B:719:ILE:HD12	2.16	0.45
1:B:304:LEU:HD13	1:B:323:LEU:HA	1.99	0.45
1:A:459:TRP:HB2	1:A:613:ASN:HB3	1.99	0.45
1:B:68:LEU:HD22	1:B:687:LEU:HD13	2.00	0.44
1:A:303:THR:H	1:A:306:GLN:HE21	1.64	0.44
1:A:270:LEU:HB2	6:A:1264:HOH:O	2.18	0.44
1:B:90:TRP:CZ2	1:B:104:GLY:HA2	2.52	0.43
1:B:717:ARG:O	1:B:721:THR:HG23	2.19	0.43
1:B:464:THR:HG22	1:B:606:TRP:HA	2.01	0.42
1:B:248:MET:HG2	1:B:372:TRP:CE2	2.54	0.42
1:B:304:LEU:HA	1:B:307:ILE:HD12	2.01	0.42
1:A:248:MET:HG2	1:A:372:TRP:CE2	2.55	0.42
1:B:490:ASN:HB2	6:B:1145:HOH:O	2.19	0.42
1:A:109:LEU:HD11	1:A:562:PRO:HD2	2.02	0.41
1:B:184:GLU:HG3	1:B:320:PHE:CE1	2.56	0.41
1:A:63:LYS:HD2	1:B:69:ILE:CG2	2.50	0.41
1:B:347:ALA:O	1:B:351:LEU:HG	2.20	0.41
1:A:312:SER:O	1:A:355:LYS:HG3	2.20	0.41
1:A:643:THR:HA	1:A:710:VAL:O	2.21	0.41
1:B:129:ILE:HD11	1:B:492:LEU:HD23	2.02	0.40
1:A:717:ARG:O	1:A:721:THR:HG23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	696/698 (100%)	677 (97%)	19 (3%)	0	100	100
1	B	694/698 (99%)	672 (97%)	22 (3%)	0	100	100
All	All	1390/1396 (100%)	1349 (97%)	41 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	607/607 (100%)	578 (95%)	29 (5%)	25	22
1	B	605/607 (100%)	574 (95%)	31 (5%)	24	19
All	All	1212/1214 (100%)	1152 (95%)	60 (5%)	24	20

All (60) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	91	LEU
1	A	154	GLU
1	A	156	LEU
1	A	160	LEU
1	A	171	GLU
1	A	175	GLN
1	A	189	GLN
1	A	197	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	260	ARG
1	A	274	LYS
1	A	278	LEU
1	A	298	LEU
1	A	314	GLU
1	A	318	LYS
1	A	333	VAL
1	A	355	LYS
1	A	368	ASN
1	A	434	GLU
1	A	483	ASP
1	A	497	LEU
1	A	498	GLU
1	A	519	SER
1	A	579	MET
1	A	597	ASN
1	A	608	GLN
1	A	667	LYS
1	A	670	GLU
1	A	687	LEU
1	A	748	VAL
1	B	91	LEU
1	B	151	ARG
1	B	154	GLU
1	B	156	LEU
1	B	160	LEU
1	B	171	GLU
1	B	175	GLN
1	B	182	THR
1	B	189	GLN
1	B	197	LYS
1	B	260	ARG
1	B	278	LEU
1	B	298	LEU
1	B	333	VAL
1	B	338	THR
1	B	368	ASN
1	B	394	ARG
1	B	434	GLU
1	B	442	LEU
1	B	456	ASP
1	B	483	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	497	LEU
1	B	498	GLU
1	B	597	ASN
1	B	608	GLN
1	B	629	SER
1	B	667	LYS
1	B	687	LEU
1	B	743	GLU
1	B	747	ARG
1	B	748	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (25) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	105	ASN
1	A	257	GLN
1	A	306	GLN
1	A	368	ASN
1	A	490	ASN
1	A	509	ASN
1	A	550	ASN
1	A	597	ASN
1	A	619	GLN
1	A	656	GLN
1	A	662	GLN
1	A	688	ASN
1	B	105	ASN
1	B	213	ASN
1	B	257	GLN
1	B	368	ASN
1	B	452	GLN
1	B	509	ASN
1	B	550	ASN
1	B	551	GLN
1	B	597	ASN
1	B	619	GLN
1	B	656	GLN
1	B	662	GLN
1	B	688	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	C	1	1,2	14,14,15	0.22	0	17,19,21	0.55	0
2	NAG	C	2	2	14,14,15	0.31	0	17,19,21	0.79	1 (5%)
2	NAG	D	1	1,2	14,14,15	0.30	0	17,19,21	0.55	0
2	NAG	D	2	2	14,14,15	0.29	0	17,19,21	0.86	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	NAG	D	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	2	NAG	C1-O5-C5	2.95	116.18	112.19

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2	NAG	C1-O5-C5	2.71	115.87	112.19

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	802	1	14,14,15	0.29	0	17,19,21	0.77	1 (5%)
3	NAG	B	801	1	14,14,15	0.32	0	17,19,21	0.78	1 (5%)
5	6LD	B	807	4	29,29,29	0.98	0	36,38,38	1.04	3 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	6LD	A	807	4	29,29,29	1.17	0	36,38,38	1.14	2 (5%)
3	NAG	B	805	1	14,14,15	0.34	0	17,19,21	0.50	0
3	NAG	A	805	1	14,14,15	0.31	0	17,19,21	0.60	0
3	NAG	A	801	1	14,14,15	0.31	0	17,19,21	0.63	0
3	NAG	B	802	1	14,14,15	0.30	0	17,19,21	0.73	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	802	1	-	1/6/23/26	0/1/1/1
3	NAG	B	801	1	-	0/6/23/26	0/1/1/1
5	6LD	B	807	4	-	3/25/25/25	0/2/2/2
5	6LD	A	807	4	-	2/25/25/25	0/2/2/2
3	NAG	B	805	1	-	0/6/23/26	0/1/1/1
3	NAG	A	805	1	-	0/6/23/26	0/1/1/1
3	NAG	A	801	1	-	0/6/23/26	0/1/1/1
3	NAG	B	802	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	807	6LD	C11-C13-C14	-3.21	106.49	113.78
5	B	807	6LD	O20-C17-C19	-3.20	116.17	122.02
5	A	807	6LD	C16-C14-N15	-3.11	103.04	110.49
3	A	802	NAG	C1-O5-C5	2.83	116.03	112.19
3	B	801	NAG	C1-O5-C5	2.62	115.74	112.19
3	B	802	NAG	C1-O5-C5	2.59	115.69	112.19
5	B	807	6LD	C11-C13-C14	-2.55	107.99	113.78
5	B	807	6LD	C19-C17-N15	2.40	120.00	115.83

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	B	807	6LD	N15-C17-C19-C23
5	B	807	6LD	O20-C17-C19-C23

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	A	807	6LD	O20-C17-C19-C23
5	A	807	6LD	N15-C17-C19-C23
3	A	802	NAG	O5-C5-C6-O6
5	B	807	6LD	C17-C19-C23-C26

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	696/698 (99%)	-0.07	13 (1%) 66 65	19, 34, 60, 83	0
1	B	696/698 (99%)	0.37	34 (4%) 29 28	22, 45, 73, 88	0
All	All	1392/1396 (99%)	0.15	47 (3%) 45 44	19, 40, 70, 88	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	333	VAL	5.8
1	B	323	LEU	5.2
1	B	317	GLY	4.1
1	B	600	GLY	3.8
1	B	179	ALA	3.7
1	B	80	THR	3.4
1	A	434	GLU	3.2
1	B	260	ARG	3.2
1	B	319	PRO	3.1
1	A	309	ASN	3.1
1	B	739	TYR	3.1
1	B	175	GLN	3.0
1	B	297	LEU	3.0
1	B	212	VAL	3.0
1	B	748	VAL	2.9
1	B	318	LYS	2.9
1	B	736	LYS	2.9
1	B	180	SER	2.9
1	B	666	LYS	2.8
1	B	607	THR	2.7
1	B	749	TRP	2.6
1	B	309	ASN	2.6
1	B	623	TYR	2.6
1	A	263	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	620	CYS	2.5
1	B	737	ASN	2.5
1	B	462	ALA	2.5
1	B	735	ARG	2.4
1	B	531	LYS	2.4
1	B	335	ILE	2.4
1	A	654	LEU	2.4
1	A	653	GLY	2.4
1	A	55	ILE	2.3
1	B	467	ARG	2.3
1	A	531	LYS	2.3
1	B	455	ASP	2.3
1	A	200	ILE	2.3
1	B	340	GLU	2.2
1	A	325	PHE	2.2
1	B	498	GLU	2.2
1	A	180	SER	2.2
1	A	652	GLY	2.2
1	B	654	LEU	2.2
1	B	466	LYS	2.1
1	A	175	GLN	2.1
1	B	337	ILE	2.1
1	A	336	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	D	1	14/15	0.76	0.20	73,82,87,92	0
2	NAG	D	2	14/15	0.78	0.25	95,97,100,103	0
2	NAG	C	2	14/15	0.90	0.13	43,48,53,54	0
2	NAG	C	1	14/15	0.97	0.08	27,34,38,41	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NAG	B	801	14/15	0.80	0.29	64,74,80,83	0
3	NAG	B	805	14/15	0.82	0.24	79,80,81,82	0
3	NAG	B	802	14/15	0.86	0.28	62,65,79,79	0
3	NAG	A	805	14/15	0.86	0.24	67,73,78,79	0
3	NAG	A	801	14/15	0.92	0.17	49,55,57,62	0
3	NAG	A	802	14/15	0.94	0.15	51,53,60,61	0
5	6LD	A	807	28/28	0.95	0.12	17,22,34,38	0
5	6LD	B	807	28/28	0.95	0.12	18,27,35,39	0
4	ZN	B	806	1/1	0.99	0.07	31,31,31,31	0
4	ZN	A	806	1/1	1.00	0.07	24,24,24,24	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 6LD A 807:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 6LD B 807:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.