



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 10, 2023 – 03:41 PM EDT

PDB ID : 7JMP  
Title : Crystal structure of SARS-CoV-2 receptor binding domain in complex with neutralizing antibody COVA2-39  
Authors : Wu, N.C.; Yuan, M.; Liu, H.; Zhu, X.; Wilson, I.A.  
Deposited on : 2020-08-02  
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5722 (1.74-1.70)
Clashscore	141614	6152 (1.74-1.70)
Ramachandran outliers	138981	6051 (1.74-1.70)
Sidechain outliers	138945	6051 (1.74-1.70)
RSRZ outliers	127900	5629 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	 16% (Poor fit), 63% (0 outliers), 8% (1 outlier), 29% (2+ outliers)
2	H	232	 % (Poor fit), 88% (0 outliers), 6% (1 outlier), 6% (2+ outliers)
3	L	216	 94% (0 outliers), 4% (1 outlier), 2% (2+ outliers)

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4953 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	165	1274	822	210	238	4	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	542	SER	-	expression tag	UNP P0DTC2
A	543	GLY	-	expression tag	UNP P0DTC2
A	544	HIS	-	expression tag	UNP P0DTC2
A	545	HIS	-	expression tag	UNP P0DTC2
A	546	HIS	-	expression tag	UNP P0DTC2
A	547	HIS	-	expression tag	UNP P0DTC2
A	548	HIS	-	expression tag	UNP P0DTC2
A	549	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called COVA2-39 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	218	1620	1017	276	320	7	0	1	0

- Molecule 3 is a protein called COVA2-39 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	211	1563	980	258	319	6	0	4	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	87	Total	O	0	0
			87	87		
5	H	186	Total	O	0	0
			186	186		
5	L	209	Total	O	0	0
			209	209		



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	68.91Å 80.44Å 72.01Å 90.00° 104.88° 90.00°	Depositor
Resolution (Å)	34.82 – 1.71 34.82 – 1.71	Depositor EDS
% Data completeness (in resolution range)	96.6 (34.82-1.71) 96.6 (34.82-1.71)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 1.71Å)	Xtrriage
Refinement program	PHENIX 1.16_3549	Depositor
R, $R_{free}$	0.177 , 0.209 0.177 , 0.209	Depositor DCC
$R_{free}$ test set	3841 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.0	Xtrriage
Anisotropy	0.183	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 47.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.017 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	4953	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.60	1/1309 (0.1%)	0.62	0/1778
2	H	0.71	2/1656 (0.1%)	0.76	0/2258
3	L	0.62	0/1611	0.70	0/2203
All	All	0.65	3/4576 (0.1%)	0.70	0/6239

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	H	108	ARG	CD-NE	-7.19	1.34	1.46
1	A	484	GLU	CD-OE1	-6.13	1.19	1.25
2	H	92	CYS	CB-SG	-5.74	1.72	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1274	0	1140	12	0
2	H	1620	0	1574	12	0
3	L	1563	0	1503	7	0
4	A	14	0	13	0	0
5	A	87	0	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	H	186	0	0	2	0
5	L	209	0	0	2	0
All	All	4953	0	4230	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:129:LYS:NZ	5:L:301:HOH:O	1.92	1.01
1:A:355:ARG:HD3	1:A:398:ASP:OD1	1.72	0.89
2:H:199:ASN:HD21	2:H:201:LYS:HE2	1.48	0.78
1:A:381:GLY:HA3	1:A:430:THR:HG22	1.73	0.70
2:H:199:ASN:ND2	2:H:201:LYS:HE2	2.18	0.58
1:A:417:LYS:HD2	5:H:469:HOH:O	2.05	0.54
3:L:194:GLN:HG2	3:L:203:GLU:HG3	1.90	0.54
2:H:87:THR:CG2	2:H:108:ARG:NH2	2.72	0.53
1:A:424:LYS:HD3	5:A:1104:HOH:O	2.08	0.52
2:H:4:LEU:HD22	2:H:22:CYS:SG	2.50	0.52
3:L:53:LYS:NZ	5:L:304:HOH:O	2.41	0.52
1:A:353:TRP:O	1:A:466:ARG:NH1	2.44	0.51
2:H:167:PRO:HG2	3:L:165:SER:OG	2.13	0.48
2:H:87:THR:HG22	2:H:108:ARG:NH2	2.28	0.48
2:H:42:GLY:HA3	3:L:163:THR:HG21	1.97	0.46
1:A:396:TYR:HB2	1:A:514:SER:OG	2.15	0.45
3:L:120:PRO:HD3	3:L:132:LEU:HD12	1.99	0.45
3:L:129:LYS:HE3	3:L:129:LYS:HB2	1.68	0.45
1:A:376:THR:HB	1:A:435:ALA:HB3	1.98	0.44
1:A:431:GLY:HA2	1:A:515:PHE:CE2	2.52	0.44
2:H:171:GLN:HG3	2:H:175:LEU:O	2.18	0.44
1:A:354:ASN:O	1:A:398:ASP:HA	2.18	0.43
1:A:498:GLN:H	1:A:501:ASN:ND2	2.17	0.42
2:H:199:ASN:HD21	2:H:201:LYS:CE	2.27	0.42
1:A:431:GLY:HA2	1:A:515:PHE:CD2	2.55	0.42
2:H:43:LYS:HA	2:H:43:LYS:HE2	2.03	0.41
1:A:353:TRP:HZ3	1:A:355:ARG:HE	1.70	0.40
2:H:100(C):GLU:HB3	5:H:374:HOH:O	2.21	0.40
2:H:42:GLY:C	2:H:43:LYS:HD2	2.42	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	157/231 (68%)	149 (95%)	8 (5%)	0	100	100
2	H	215/232 (93%)	211 (98%)	4 (2%)	0	100	100
3	L	213/216 (99%)	208 (98%)	5 (2%)	0	100	100
All	All	585/679 (86%)	568 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	126/203 (62%)	124 (98%)	2 (2%)	62	47
2	H	179/194 (92%)	177 (99%)	2 (1%)	73	62
3	L	174/181 (96%)	172 (99%)	2 (1%)	73	62
All	All	479/578 (83%)	473 (99%)	6 (1%)	73	55

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	355	ARG
1	A	392	PHE
2	H	108	ARG
2	H	149	PRO
3	L	137[A]	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
3	L	137[B]	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	501	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	1001	1	14,14,15	1.66	1 (7%)	17,19,21	2.15	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	1001	1	-	4/6/23/26	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1001	NAG	O5-C1	6.09	1.53	1.43

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1001	NAG	C1-O5-C5	8.65	123.92	112.19

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1001	NAG	O5-C5-C6-O6
4	A	1001	NAG	C8-C7-N2-C2
4	A	1001	NAG	O7-C7-N2-C2
4	A	1001	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	165/231 (71%)	0.88	36 (21%) 0 0	21, 45, 81, 94	0
2	H	218/232 (93%)	0.06	2 (0%) 84 87	19, 28, 42, 54	0
3	L	211/216 (97%)	-0.19	0 100 100	18, 27, 40, 51	0
All	All	594/679 (87%)	0.20	38 (6%) 19 21	18, 30, 61, 94	0

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	395	VAL	7.0
1	A	381	GLY	6.5
1	A	392	PHE	6.1
1	A	339	GLY	5.7
1	A	515	PHE	5.6
1	A	382	VAL	5.6
1	A	338	PHE	5.0
1	A	368	LEU	4.9
1	A	358	ILE	4.9
1	A	513	LEU	4.5
1	A	394	ASN	4.3
1	A	380	TYR	4.1
1	A	369	TYR	4.0
1	A	383	SER	4.0
1	A	379	CYS	3.8
1	A	393	THR	3.6
1	A	516	GLU	3.6
1	A	359	SER	3.6
1	A	340	GLU	3.4
1	A	384	PRO	3.3
1	A	432	CYS	3.3
1	A	345	THR	3.1
1	A	415	THR	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	430	THR	2.9
1	A	357	ARG	2.9
1	A	367	VAL	2.8
1	A	445	VAL	2.8
1	A	341	VAL	2.8
1	A	416	GLY	2.7
1	A	514	SER	2.7
1	A	396	TYR	2.6
2	H	37	VAL	2.5
2	H	50	VAL	2.5
1	A	372	ALA	2.5
1	A	431	GLY	2.5
1	A	366	SER	2.3
1	A	446	GLY	2.2
1	A	355	ARG	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	NAG	A	1001	14/15	0.82	0.22	80,83,88,89	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.