



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 13, 2024 – 08:04 AM EDT

PDB ID : 4JSN  
Title : structure of mTORdeltaN-mLST8 complex  
Authors : Pavletich, N.P.  
Deposited on : 2013-03-22  
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1174	 6% 68% 19% 10%
1	B	1174	 3% 69% 18% 10%
2	C	326	 3% 67% 26% 10%
2	D	326	 2% 68% 26% 10%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 22097 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Serine/threonine-protein kinase mTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	1058	8608	5472	1521	1552	63	0	0	0
1	A	1054	8577	5451	1517	1546	63	0	0	0


- Molecule 2 is a protein called Target of rapamycin complex subunit LST8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	317	2456	1526	436	476	18	0	0	0
2	C	317	2456	1526	436	476	18	0	0	0







● Molecule 1: Serine/threonine-protein kinase mTOR



● Molecule 2: Target of rapamycin complex subunit LST8



• Molecule 2: Target of rapamycin complex subunit LST8



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	139.40Å 163.20Å 207.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.86 – 3.20 29.84 – 3.20	Depositor EDS
% Data completeness (in resolution range)	86.6 (29.86-3.20) 86.5 (29.84-3.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 3.18Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.215 , 0.256 0.219 , 0.253	Depositor DCC
$R_{free}$ test set	1857 reflections (2.46%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	56.0	Xtrriage
Anisotropy	0.333	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 42.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	22097	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	87.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 43.98 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6294e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/8772	0.85	10/11872 (0.1%)
1	B	0.55	0/8805	0.85	9/11920 (0.1%)
2	C	0.55	0/2514	0.88	2/3426 (0.1%)
2	D	0.58	0/2514	0.90	0/3426
All	All	0.55	0/22605	0.86	21/30644 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
2	C	0	1
2	D	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2378	ARG	NE-CZ-NH1	7.37	123.99	120.30
1	B	2378	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	B	2378	ARG	NE-CZ-NH1	6.88	123.74	120.30
1	A	2378	ARG	NE-CZ-NH2	-6.23	117.18	120.30
1	A	2344	LEU	CA-CB-CG	6.14	129.41	115.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	1444	GLU	Peptide
1	A	1680	ASP	Peptide
1	B	1444	GLU	Peptide
1	B	1680	ASP	Peptide
2	D	267	GLU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8577	0	8559	114	1
1	B	8608	0	8593	111	0
2	C	2456	0	2341	43	0
2	D	2456	0	2341	43	0
All	All	22097	0	21834	299	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 299 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2380:THR:HG22	1:A:2383:LEU:HG	1.53	0.91
1:B:2380:THR:HG22	1:B:2383:LEU:HG	1.52	0.89
1:B:1930:ILE:HD11	1:B:1934:THR:HG21	1.57	0.87
1:A:1930:ILE:HD11	1:A:1934:THR:HG21	1.58	0.84
2:C:219:HIS:NE2	2:C:237:THR:HG22	1.97	0.78

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1612:ARG:NH2	1:A:1612:ARG:NH2[2_554]	1.96	0.24

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1046/1174 (89%)	967 (92%)	72 (7%)	7 (1%)	22	61
1	B	1052/1174 (90%)	967 (92%)	77 (7%)	8 (1%)	19	58
2	C	315/326 (97%)	285 (90%)	24 (8%)	6 (2%)	8	39
2	D	315/326 (97%)	284 (90%)	24 (8%)	7 (2%)	6	35
All	All	2728/3000 (91%)	2503 (92%)	197 (7%)	28 (1%)	15	54

5 of 28 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	1630	VAL
1	B	1692	VAL
1	B	1937	GLN
2	D	74	ASN
1	A	1692	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	927/1024 (90%)	831 (90%)	96 (10%)	7	28
1	B	931/1024 (91%)	834 (90%)	97 (10%)	7	28
2	C	269/276 (98%)	233 (87%)	36 (13%)	4	18
2	D	269/276 (98%)	234 (87%)	35 (13%)	4	19
All	All	2396/2600 (92%)	2132 (89%)	264 (11%)	6	26

5 of 264 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	C	82	ASP
2	C	122	GLN
2	C	270	ARG
2	D	13	VAL
1	B	2519	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 54 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1492	GLN
1	A	1807	GLN
2	C	137	HIS
1	A	1496	GLN
1	A	1693	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	1054/1174 (89%)	0.12	71 (6%) 17 10	38, 86, 178, 243	0
1	B	1058/1174 (90%)	-0.14	36 (3%) 45 29	28, 72, 150, 238	0
2	C	317/326 (97%)	-0.08	11 (3%) 44 28	40, 77, 147, 222	0
2	D	317/326 (97%)	-0.36	6 (1%) 66 53	27, 52, 116, 192	0
All	All	2746/3000 (91%)	-0.06	124 (4%) 33 21	27, 74, 170, 243	0

The worst 5 of 124 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1579	ALA	8.6
1	A	1580	GLY	7.1
1	B	1580	GLY	6.6
1	A	1559	LEU	6.2
1	B	2436	THR	6.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.