



Full wwPDB NMR Structure Validation Report ⓘ

Aug 16, 2020 – 07:32 PM BST

PDB ID : 2JUE
Title : Solution structure of the all-D kalata B1
Authors : Daly, N.L.; Sando, L.; Craik, D.
Deposited on : 2007-08-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.13.1
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	29	90% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA and RNA chains that are outliers for geometric criteria:

Mol	Chain	Compound	Res	Total models with violations	
				Chirality	Geometry
1	A	DTH	12	20	-
1	A	DTH	16	20	-
1	A	DTH	23	20	-
1	A	DTH	4	20	-
1	A	DTH	9	20	-

2 Ensemble composition and analysis

This entry contains 20 models.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms (5) was below the domain threshold value (8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 376 atoms, of which 179 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Kalata-B1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	29	376	117	179	35	39	6	0

There is a discrepancy between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	29	DVA	SER	SEE REMARK 999	UNP P56254

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Kalata-B1




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Kalata-B1



4.2.2 Score per residue for model 2


- Molecule 1: Kalata-B1



4.2.3 Score per residue for model 3


- Molecule 1: Kalata-B1


Chain A:  93% 7%



4.2.4 Score per residue for model 4


- Molecule 1: Kalata-B1

Chain A:  86% 14%



4.2.5 Score per residue for model 5


- Molecule 1: Kalata-B1


Chain A:  79% 21%



4.2.6 Score per residue for model 6


- Molecule 1: Kalata-B1


Chain A:  86% 14%



4.2.7 Score per residue for model 7


- Molecule 1: Kalata-B1

Chain A:  86% 14%




4.2.8 Score per residue for model 8

- Molecule 1: Kalata-B1




4.2.9 Score per residue for model 9

- Molecule 1: Kalata-B1




4.2.10 Score per residue for model 10

- Molecule 1: Kalata-B1




4.2.11 Score per residue for model 11

- Molecule 1: Kalata-B1




4.2.12 Score per residue for model 12

- Molecule 1: Kalata-B1




4.2.13 Score per residue for model 13

- Molecule 1: Kalata-B1




4.2.14 Score per residue for model 14

- Molecule 1: Kalata-B1




4.2.15 Score per residue for model 15

- Molecule 1: Kalata-B1




4.2.16 Score per residue for model 16

- Molecule 1: Kalata-B1




4.2.17 Score per residue for model 17

- Molecule 1: Kalata-B1



4.2.18 Score per residue for model 18


- Molecule 1: Kalata-B1


Chain A:  79% 21%



4.2.19 Score per residue for model 19


- Molecule 1: Kalata-B1


Chain A:  83% 17%



4.2.20 Score per residue for model 20

- Molecule 1: Kalata-B1

Chain A:  79% 21%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	structure solution	
CNS	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DTH, DPR, DLE, DVA, DGL, DSN, DAR, DCY, DTR, DSG

There are no covalent bond-length or bond-angle outliers.

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	5.0±0.0	0.0±0.0
All	All	100	0

There are no bond-length outliers.

There are no bond-angle outliers.

All unique chiral outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	4	DTH	CB	20
1	A	9	DTH	CB	20
1	A	12	DTH	CB	20
1	A	16	DTH	CB	20
1	A	23	DTH	CB	20

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	197	179	159	3±1
All	All	3940	3580	3192	59

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 8.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:27:DLE:HG	1:A:29:DVA:HG13	0.56	1.77	14	4
1:A:1:DCY:SG	1:A:24:DAR:HB2	0.54	2.43	11	7
1:A:12:DTH:HB	1:A:15:DCY:SG	0.53	2.44	17	2
1:A:1:DCY:HA	1:A:24:DAR:NH1	0.51	2.21	8	1
1:A:15:DCY:SG	1:A:24:DAR:HD3	0.51	2.46	20	2
1:A:27:DLE:HD22	1:A:28:DPR:HD3	0.49	1.84	17	8
1:A:24:DAR:C	1:A:26:GLY:H	0.49	2.20	11	1
1:A:12:DTH:HG23	1:A:15:DCY:SG	0.46	2.51	15	1
1:A:3:DGL:CG	1:A:22:DCY:HB2	0.45	2.41	16	2
1:A:3:DGL:HG3	1:A:22:DCY:HB2	0.45	1.87	16	4
1:A:3:DGL:HB3	1:A:22:DCY:HB2	0.45	1.89	19	1
1:A:3:DGL:HB2	1:A:22:DCY:HB2	0.44	1.90	11	1
1:A:1:DCY:SG	1:A:24:DAR:HB3	0.43	2.53	8	1
1:A:18:DSN:O	1:A:19:DTR:C	0.42	2.68	13	19
1:A:19:DTR:HA	1:A:20:DPR:HA	0.42	1.69	19	3
1:A:3:DGL:HG2	1:A:11:DSG:HB2	0.41	1.92	17	1
1:A:24:DAR:C	1:A:26:GLY:N	0.40	2.84	11	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	5/29 (17%)	5±0 (99±4%)	0±0 (1±4%)	0±0 (0±0%)	100	100
All	All	100/580 (17%)	99 (99%)	1 (1%)	0 (0%)	100	100

There are no Ramachandran outliers.

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	0	-	-	-
All	All	0	-	-	-

There are no protein residues with a non-rotameric sidechain to report.

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DTH	A	12	1	1±0,1,2,3	-	-
1	DTH	A	23	1	1±0,1,2,3	-	-
1	DTH	A	4	1	1±0,1,2,3	-	-
1	DTH	A	16	1	1±0,1,2,3	-	-
1	DTH	A	9	1	1±0,1,2,3	-	-

There are no bond-length outliers.

There are no bond-angle outliers.

All unique chiral outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	23	DTH	CB	20
1	A	9	DTH	CB	20
1	A	16	DTH	CB	20
1	A	4	DTH	CB	20
1	A	12	DTH	CB	20

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided