



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2021 – 06:54 AM EDT

PDB ID : 3JWS  
Title : Structure of neuronal nitric oxide synthase R349A mutant heme domain complexed with N1-[(3' S,4'S)-4'-((6"-amino-4"-methylpyridin-2"-yl)methyl)pyrrolidin-3'-yl]-N2-(3'-fluorophenethyl)ethane-1,2-diamine tetrahydrochloride  
Authors : Delker, S.L.; Li, H.; Poulos, T.L.  
Deposited on : 2009-09-18  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2580 (1.96-1.96)
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.


Mol	Chain	Length	Quality of chain
1	A	422	
1	B	422	



Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula:  $C_9H_{15}N_5O_3$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is N-((3S,4S)-4-[(6-amino-4-methylpyridin-2-yl)methyl]pyrrolidin-3-yl)-N'-[2-(3-fluorophenyl)ethyl]ethane-1,2-diamine (three-letter code: JI4) (formula:  $C_{21}H_{30}FN_5$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	F			N
4	A	1	27	21	1	5	0	0
4	B	1	27	21	1	5	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	B	1	4	2	2	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Zn 1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	134	Total 134	O 134	0	0
7	B	197	Total 197	O 197	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, brain



- Molecule 1: Nitric oxide synthase, brain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.94Å 112.19Å 164.47Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.92 – 1.95 43.92 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.8 (43.92-1.95) 97.8 (43.92-1.95)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	0.04	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.93 (at 1.95Å)	Xtrriage
Refinement program	REFMAC 5.5.0089, CNS	Depositor
R, $R_{free}$	0.187 , 0.221 0.225 , 0.254	Depositor DCC
$R_{free}$ test set	3458 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.7	Xtrriage
Anisotropy	0.203	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7166	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, H4B, HEM, ACT, JI4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.55	0/3406	0.60	1/4621 (0.0%)
1	B	0.60	0/3432	0.62	0/4654
All	All	0.57	0/6838	0.61	1/9275 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	523	LEU	CA-CB-CG	5.07	126.97	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3313	0	3221	25	0
1	B	3339	0	3251	15	0
2	A	43	0	30	5	0
2	B	43	0	30	3	0
3	A	17	0	15	0	0
3	B	17	0	15	0	0
4	A	27	0	30	2	0
4	B	27	0	30	2	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	1	0	0	0	0
7	A	134	0	0	4	0
7	B	197	0	0	1	0
All	All	7166	0	6628	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:706:TYR:OH	2:B:750:HEM:O1D	1.90	0.89
2:A:750:HEM:HMC2	2:A:750:HEM:HBC2	1.60	0.82
1:A:706:TYR:OH	2:A:750:HEM:O2D	2.03	0.76
1:A:636:ALA:C	7:A:1053:HOH:O	2.27	0.73
1:A:636:ALA:O	7:A:1053:HOH:O	2.14	0.65
1:B:351:LYS:HE2	1:B:392:SER:OG	1.99	0.63
1:A:639:TYR:N	7:A:1053:HOH:O	2.32	0.61
1:A:355:PHE:HB2	7:A:1024:HOH:O	2.02	0.60
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.85	0.58
1:A:706:TYR:OH	2:A:750:HEM:CGD	2.55	0.54
1:A:337:LEU:HG	4:A:800:JI4:F13	1.98	0.53
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.91	0.53
2:A:750:HEM:HBB2	2:A:750:HEM:HHC	1.93	0.50
1:B:478:GLN:HB2	1:B:481:ARG:HG3	1.92	0.50
1:B:701:THR:HA	1:B:702:PRO:C	2.33	0.48
1:B:391:THR:O	1:B:392:SER:HB2	2.14	0.47
1:B:403:TYR:CE1	1:B:407:HIS:CE1	3.03	0.47
1:A:537:PRO:HB2	1:A:540:LEU:HG	1.97	0.47
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.98	0.46
1:A:316:LEU:HD12	1:A:319:LYS:HD2	1.98	0.45
2:A:750:HEM:HBC2	2:A:750:HEM:CMC	2.37	0.45
1:A:308:THR:O	1:A:309:ASP:HB2	2.16	0.45
1:A:455:LEU:HD12	1:A:587:TRP:HB3	2.00	0.44
1:A:510:TRP:CE2	1:A:521:PRO:HD3	2.53	0.44
1:B:337:LEU:HG	4:B:800:JI4:F13	2.08	0.43
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	2.01	0.43
1:A:332:MET:CE	1:B:301:LEU:HD22	2.49	0.43
1:B:338:PRO:HG3	7:B:1172:HOH:O	2.18	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:451:ASN:HB3	1:A:454:ASN:O	2.19	0.43
1:A:676:TRP:CE2	1:A:680:VAL:HG21	2.54	0.42
1:B:678:TRP:HH2	4:B:800:JI4:H4	1.84	0.42
1:A:701:THR:HA	1:A:702:PRO:C	2.40	0.42
1:A:403:TYR:CE1	1:A:407:HIS:CE1	3.08	0.42
1:A:439:PHE:CZ	1:A:443:CYS:SG	3.13	0.41
1:B:584:PHE:CD1	2:B:750:HEM:CAC	3.04	0.41
1:A:696:LEU:HB3	1:B:330:ILE:HD11	2.01	0.41
1:A:352:ASP:OD2	1:A:352:ASP:N	2.54	0.41
1:B:681:PRO:HG2	1:B:688:THR:HG21	2.03	0.41
1:A:592:GLU:OE1	4:A:800:JI4:N11	2.54	0.40
1:A:299:ARG:O	1:A:317:HIS:CE1	2.74	0.40
1:B:589:MET:HA	1:B:649:VAL:O	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/422 (96%)	396 (98%)	7 (2%)	0	100	100
1	B	407/422 (96%)	402 (99%)	5 (1%)	0	100	100
All	All	810/844 (96%)	798 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/376 (96%)	354 (98%)	9 (2%)	47	38
1	B	365/376 (97%)	359 (98%)	6 (2%)	62	58
All	All	728/752 (97%)	713 (98%)	15 (2%)	53	46

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	350	THR
1	A	454	ASN
1	A	523	LEU
1	A	547	ARG
1	A	555	LYS
1	A	569	ASN
1	A	620	LYS
1	A	645	LYS
1	A	713	THR
1	B	299	ARG
1	B	321	THR
1	B	336	MET
1	B	389	GLU
1	B	454	ASN
1	B	547	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	364	GLN
1	A	425	GLN
1	A	440	ASN
1	A	454	ASN
1	A	527	ASN
1	A	569	ASN
1	A	605	ASN
1	A	628	GLN
1	A	642	GLN
1	A	697	ASN
1	B	364	GLN
1	B	385	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	425	GLN
1	B	454	ASN
1	B	507	GLN
1	B	508	GLN
1	B	527	ASN
1	B	535	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	H4B	B	761	-	16,18,18	0.93	1 (6%)	11,26,26	2.73	6 (54%)
3	H4B	A	760	-	16,18,18	1.04	2 (12%)	11,26,26	3.12	6 (54%)
4	J14	A	800	-	27,29,29	0.63	0	30,38,38	1.48	4 (13%)
4	J14	B	800	-	27,29,29	0.66	0	30,38,38	1.91	6 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	ACT	B	861	-	1,3,3	1.53	0	0,3,3	-	-
5	ACT	A	860	-	1,3,3	1.44	0	0,3,3	-	-
2	HEM	A	750	1	27,50,50	2.18	9 (33%)	17,82,82	2.32	6 (35%)
2	HEM	B	750	1	27,50,50	2.25	8 (29%)	17,82,82	2.50	4 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	B	761	-	-	0/8/17/17	0/2/2/2
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	J14	A	800	-	-	1/13/23/23	0/3/3/3
4	J14	B	800	-	-	1/13/23/23	0/3/3/3
2	HEM	A	750	1	-	1/6/54/54	-
2	HEM	B	750	1	-	0/6/54/54	-

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	750	HEM	C3D-C2D	5.14	1.52	1.37
2	A	750	HEM	C3D-C2D	4.88	1.52	1.37
2	B	750	HEM	C3C-C2C	-4.80	1.33	1.40
2	B	750	HEM	C3B-C2B	-4.67	1.33	1.40
2	A	750	HEM	C3C-C2C	-4.52	1.34	1.40
2	A	750	HEM	C3B-C2B	-4.29	1.34	1.40
2	B	750	HEM	C3C-CAC	3.65	1.55	1.47
2	A	750	HEM	C3B-CAB	3.59	1.55	1.47
2	A	750	HEM	C3C-CAC	3.35	1.54	1.47
2	B	750	HEM	C3B-CAB	3.31	1.54	1.47
3	A	760	H4B	C2-N2	2.74	1.39	1.33
2	B	750	HEM	CAD-C3D	2.42	1.56	1.52
2	A	750	HEM	CMB-C2B	2.40	1.57	1.51
2	A	750	HEM	CMD-C2D	2.35	1.56	1.51
3	B	761	H4B	C7-C6	2.17	1.54	1.52
2	B	750	HEM	CMD-C2D	2.15	1.56	1.51
2	B	750	HEM	CMB-C2B	2.08	1.56	1.51
2	A	750	HEM	CMA-C3A	2.07	1.55	1.51
3	A	760	H4B	C4-N3	2.05	1.36	1.33
2	A	750	HEM	CMC-C2C	2.00	1.56	1.51

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	CBA-CAA-C2A	-7.39	98.86	112.49
3	B	761	H4B	C4-C4A-C8A	6.74	120.56	114.57
4	B	800	JI4	C61-N11-C21	5.67	122.40	118.10
2	A	750	HEM	CBD-CAD-C3D	-5.51	102.33	112.48
3	A	760	H4B	C4-C4A-N5	5.36	123.62	119.12
4	B	800	JI4	C1-N1-C3'	5.34	121.76	114.20
2	A	750	HEM	CBA-CAA-C2A	-5.23	102.84	112.49
2	B	750	HEM	CBD-CAD-C3D	-4.69	103.83	112.48
3	A	760	H4B	C4-C4A-C8A	4.69	118.73	114.57
3	A	760	H4B	N3-C2-N1	-4.03	119.10	125.42
3	A	760	H4B	C4-N3-C2	3.82	122.01	115.93
4	A	800	JI4	C1-N1-C3'	3.76	119.51	114.20
4	A	800	JI4	C61-N11-C21	3.73	120.93	118.10
4	B	800	JI4	C31-C21-N11	-3.52	119.17	122.90
4	A	800	JI4	C5'-N1'-C2'	3.36	113.35	105.42
2	A	750	HEM	C1D-C2D-C3D	-3.36	104.66	107.00
3	A	760	H4B	C2-N1-C8A	3.20	121.71	114.54
4	A	800	JI4	C31-C21-N11	-3.14	119.57	122.90
4	B	800	JI4	C5'-N1'-C2'	3.13	112.80	105.42
3	B	761	H4B	C4-C4A-N5	3.10	121.72	119.12
3	A	760	H4B	N2-C2-N3	3.05	122.00	117.25
3	B	761	H4B	C4-N3-C2	2.82	120.41	115.93
2	B	750	HEM	C1D-C2D-C3D	-2.77	105.07	107.00
2	B	750	HEM	CMA-C3A-C4A	-2.51	124.61	128.46
2	A	750	HEM	CMA-C3A-C4A	-2.42	124.75	128.46
2	A	750	HEM	C4C-C3C-C2C	2.23	108.45	106.90
3	B	761	H4B	C2-N1-C8A	2.11	119.26	114.54
3	B	761	H4B	N3-C2-N1	-2.09	122.14	125.42
2	A	750	HEM	CMC-C2C-C3C	2.08	128.57	124.68
4	B	800	JI4	C2'-C3'-N1	-2.08	110.03	113.73
3	B	761	H4B	C4A-C4-N3	-2.06	118.16	124.01
4	B	800	JI4	C51-C41-C31	2.00	120.48	118.09

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	800	JI4	C2-C1-N1-C3'
4	B	800	JI4	C2-C1-N1-C3'
2	A	750	HEM	C2A-CAA-CBA-CGA


There are no ring outliers.


4 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	800	JI4	2	0
4	B	800	JI4	2	0
2	A	750	HEM	5	0
2	B	750	HEM	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	407/422 (96%)	1.32	91 (22%) <b>0</b> <b>0</b>	26, 51, 96, 119	0
1	B	411/422 (97%)	0.87	53 (12%) <b>3</b> <b>5</b>	26, 40, 64, 88	0
All	All	818/844 (96%)	1.09	144 (17%) <b>1</b> <b>1</b>	26, 45, 88, 119	0

All (144) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	715	VAL	8.0
1	A	716	TRP	7.2
1	A	388	ILE	6.2
1	A	355	PHE	6.0
1	A	619	ARG	6.0
1	B	350	THR	5.8
1	A	352	ASP	5.8
1	A	470	HIS	5.8
1	B	348	VAL	5.8
1	A	506	ILE	5.6
1	B	300	PHE	5.5
1	A	389	GLU	5.2
1	B	619	ARG	4.7
1	A	488	PRO	4.4
1	A	386	LYS	4.4
1	A	390	SER	4.4
1	A	486	LYS	4.4
1	B	351	LYS	4.3
1	A	391	THR	4.3
1	A	469	LYS	4.2
1	A	503	GLU	4.2
1	B	479	LEU	4.1
1	B	564	LEU	4.1
1	B	677	VAL	4.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	511	LYS	3.9
1	A	553	TRP	3.9
1	B	567	VAL	3.9
1	A	382	GLU	3.9
1	A	612	LYS	3.8
1	A	351	LYS	3.8
1	B	321	THR	3.8
1	B	667	ARG	3.7
1	A	385	ASN	3.7
1	A	384	VAL	3.7
1	B	566	ALA	3.6
1	A	479	LEU	3.6
1	A	551	PHE	3.6
1	A	490	GLY	3.6
1	A	321	THR	3.5
1	B	541	VAL	3.4
1	A	676	TRP	3.4
1	B	446	VAL	3.4
1	A	508	GLN	3.3
1	B	442	ILE	3.3
1	B	318	LEU	3.2
1	A	356	PRO	3.2
1	B	676	TRP	3.1
1	A	714	HIS	3.1
1	A	507	GLN	3.1
1	B	352	ASP	3.1
1	A	567	VAL	3.1
1	A	677	VAL	3.1
1	A	712	ASN	3.0
1	A	541	VAL	3.0
1	A	556	ASP	3.0
1	A	376	ALA	3.0
1	A	679	ILE	2.9
1	A	485	TYR	2.9
1	B	329	HIS	2.9
1	A	350	THR	2.9
1	A	354	LEU	2.9
1	A	517	PHE	2.9
1	B	680	VAL	2.9
1	B	480	ILE	2.9
1	A	499	VAL	2.8
1	B	299	ARG	2.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	349	ALA	2.8
1	B	588	TYR	2.8
1	B	718	GLY	2.8
1	B	389	GLU	2.8
1	B	591	THR	2.8
1	B	620	LYS	2.7
1	B	679	ILE	2.7
1	A	394	TYR	2.7
1	A	667	ARG	2.7
1	B	443	CYS	2.7
1	A	487	GLN	2.7
1	B	678	TRP	2.7
1	A	550	LYS	2.7
1	B	565	PRO	2.7
1	A	536	ILE	2.6
1	A	493	LEU	2.6
1	A	682	PRO	2.6
1	A	373	GLY	2.6
1	A	381	LEU	2.6
1	A	680	VAL	2.6
1	A	557	LEU	2.6
1	B	611	ALA	2.6
1	A	608	GLU	2.6
1	A	380	ARG	2.5
1	A	537	PRO	2.5
1	B	474	VAL	2.5
1	A	706	TYR	2.5
1	B	542	LEU	2.5
1	B	589	MET	2.5
1	A	492	THR	2.5
1	B	568	SER	2.5
1	A	611	ALA	2.4
1	A	299	ARG	2.4
1	B	617	ASP	2.4
1	B	593	ILE	2.4
1	B	392	SER	2.4
1	A	530	ASP	2.4
1	A	519	VAL	2.4
1	B	416	VAL	2.4
1	A	520	LEU	2.4
1	A	315	THR	2.4
1	A	442	ILE	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	446	VAL	2.4
1	A	378	MET	2.3
1	A	678	TRP	2.3
1	A	582	CYS	2.3
1	B	550	LYS	2.3
1	B	353	GLN	2.3
1	A	552	ASP	2.3
1	A	393	THR	2.3
1	A	620	LYS	2.3
1	A	379	ASP	2.3
1	A	533	LEU	2.3
1	B	590	GLY	2.2
1	A	685	GLY	2.2
1	A	300	PHE	2.2
1	B	583	PRO	2.2
1	A	467	ASP	2.2
1	B	682	PRO	2.2
1	B	562	TYR	2.2
1	A	554	PHE	2.2
1	B	715	VAL	2.2
1	A	491	SER	2.1
1	B	681	PRO	2.1
1	A	535	GLN	2.1
1	A	372	PHE	2.1
1	A	711	TRP	2.1
1	A	423	LYS	2.1
1	B	439	PHE	2.1
1	B	310	VAL	2.1
1	A	558	GLY	2.1
1	B	335	ILE	2.1
1	A	318	LEU	2.1
1	A	686	SER	2.0
1	A	566	ALA	2.0
1	B	386	LYS	2.0
1	A	482	TYR	2.0
1	A	484	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	J14	A	800	27/27	0.93	0.19	34,39,65,67	0
4	J14	B	800	27/27	0.94	0.22	33,36,57,59	0
3	H4B	A	760	17/17	0.96	0.16	29,32,37,39	0
3	H4B	B	761	17/17	0.96	0.21	26,30,34,36	0
5	ACT	A	860	4/4	0.96	0.23	54,55,56,56	0
5	ACT	B	861	4/4	0.96	0.13	51,52,52,52	0
2	HEM	A	750	43/43	0.97	0.17	29,31,43,45	0
2	HEM	B	750	43/43	0.97	0.17	26,30,44,48	0
6	ZN	A	900	1/1	0.99	0.11	36,36,36,36	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


**Electron density around JI4 A 800:**


$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around JI4 B 800:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)







## 6.5 Other polymers [i](#)

There are no such residues in this entry.