



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 13, 2024 – 04:17 AM EST

PDB ID : 3JWY
Title : Structure of endothelial nitric oxide synthase heme domain complexed with N 1-{(3'R,4'S)-4'-[(6"-amino-4"-methylpyridin-2"-yl)methyl]pyrrolidin-3'-yl}-N 2-(3'-fluorophenethyl)ethane-1,2-diamine tetrahydrochloride
Authors : Delker, S.L.; Li, H.; Poulos, T.L.
Deposited on : 2009-09-18
Resolution : 2.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2391 (2.26-2.22)
Clashscore	141614	2539 (2.26-2.22)
Ramachandran outliers	138981	2489 (2.26-2.22)
Sidechain outliers	138945	2490 (2.26-2.22)
RSRZ outliers	127900	2353 (2.26-2.22)


The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	444	 3% 82% 7% • 9%
1	B	444	 2% 82% 8% • 9%

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	Fe	N			O
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	17	9	5	3	0	0
3	B	1	17	9	5	3	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is N-{(3R,4S)-4-[(6-amino-4-methylpyridin-2-yl)methyl]pyrrolidin-3-yl}-N'-[2-(3-fluorophenyl)ethyl]ethane-1,2-diamine (three-letter code: J11) (formula: C₂₁H₃₀FN₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	F	N		
6	A	1	27	21	1	5	0	0
6	B	1	27	21	1	5	0	0

- Molecule 7 is CACODYLIC ACID (three-letter code: CAD) (formula: C₂H₇AsO₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	As	C	0	0
			3	1	2		
7	B	1	Total	As	C	0	0
			3	1	2		

- Molecule 8 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Zn	0	0
			1	1		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	211	Total	O	0	0
			211	211		
9	B	206	Total	O	0	0
			206	206		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nitric oxide synthase, endothelial



- Molecule 1: Nitric oxide synthase, endothelial



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.11Å 106.93Å 157.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.20 – 2.24 44.20 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.9 (44.20-2.24) 99.9 (44.20-2.24)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 2.24Å)	Xtrriage
Refinement program	REFMAC 5.5.0089, CNS	Depositor
R, R_{free}	0.174 , 0.217 0.182 , 0.223	Depositor DCC
R_{free} test set	2368 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	34.6	Xtrriage
Anisotropy	0.099	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7028	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ACT, H4B, CAD, ZN, J11, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/3299	0.59	0/4494
1	B	0.52	1/3289 (0.0%)	0.59	0/4480
All	All	0.52	1/6588 (0.0%)	0.59	0/8974

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	203	CYS	CB-SG	-5.06	1.73	1.81

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3210	0	3109	26	0
1	B	3200	0	3104	20	0
2	A	43	0	30	1	0
2	B	43	0	30	4	0
3	A	17	0	15	1	0
3	B	17	0	15	1	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	6	0	8	0	0
5	B	6	0	8	1	0
6	A	27	0	30	0	0
6	B	27	0	30	2	0
7	A	3	0	0	2	0
7	B	3	0	0	1	0
8	A	1	0	0	0	0
9	A	211	0	0	2	0
9	B	206	0	0	4	0
All	All	7028	0	6385	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:CYS:SG	7:A:950:CAD:AS	2.48	1.32
1:A:257:ARG:HG3	1:A:257:ARG:HH11	1.36	0.87
1:A:260[B]:ASP:O	1:A:260[B]:ASP:OD2	1.95	0.84
1:A:257:ARG:HG3	1:A:257:ARG:NH1	2.05	0.70
1:A:260[B]:ASP:O	1:A:260[B]:ASP:CG	2.30	0.68
1:B:324:TRP:HB2	7:B:950:CAD:C1	2.24	0.67
1:B:236:ARG:HD2	1:B:242:ASP:OD1	1.94	0.67
1:A:285:ASN:HB3	9:A:1190:HOH:O	1.96	0.65
1:B:126:LEU:HD11	1:B:156:GLU:HG2	1.82	0.60
1:A:384:CYS:CB	7:A:950:CAD:AS	3.11	0.59
1:B:178:GLN:HE22	1:B:181:ARG:HH11	1.51	0.58
1:B:215:ASN:HB2	9:B:1173:HOH:O	2.04	0.57
1:A:91:GLN:HE21	1:A:91:GLN:HA	1.70	0.57
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.45	0.57
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	1.86	0.56
1:B:389:THR:HG22	1:B:389:THR:O	2.06	0.56
1:A:378:LEU:HB2	9:A:1001:HOH:O	2.06	0.56
1:B:378:LEU:HB2	9:B:1006:HOH:O	2.06	0.55
1:A:249:GLN:HB2	1:A:252:ARG:HG2	1.88	0.55
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.90	0.52
1:B:249:GLN:HB2	1:B:252:ARG:HG3	1.90	0.52
1:A:260[A]:ASP:OD1	1:A:260[A]:ASP:N	2.36	0.52
2:B:500:HEM:HBA1	6:B:800:J11:H71A	1.92	0.52
1:B:215:ASN:HB2	9:B:1179:HOH:O	2.10	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:449:TRP:HA	3:B:600:H4B:N1	2.26	0.50
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.41	0.50
1:A:465:GLU:HB3	1:B:105:LEU:HD22	1.93	0.50
1:A:126:LEU:HD23	1:A:130:ARG:HH21	1.78	0.49
1:A:152:LEU:O	1:A:156:GLU:HG2	2.14	0.47
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.97	0.46
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.55	0.46
2:A:500:HEM:HBB2	2:A:500:HEM:HHC	1.99	0.45
1:A:126:LEU:HD11	1:A:156:GLU:HA	1.98	0.45
1:B:72:ARG:HH11	1:B:72:ARG:HG2	1.82	0.45
1:B:366:THR:O	1:B:370:CYS:HB2	2.17	0.44
1:A:449:TRP:HA	3:A:600:H4B:N1	2.31	0.44
1:A:455:SER:O	1:A:458:LEU:HB2	2.17	0.44
1:A:472:SER:HA	1:A:473:PRO:C	2.38	0.44
1:A:151:ARG:HD3	1:A:168:ARG:NH2	2.32	0.44
1:A:196:GLN:HG2	1:A:219:TYR:CZ	2.54	0.43
1:B:231:THR:O	1:B:354:PRO:HD2	2.17	0.43
1:B:480:ASP:HA	1:B:481:PRO:HD3	1.94	0.43
1:A:257:ARG:HH11	1:A:257:ARG:CG	2.18	0.42
1:B:233:PHE:HB3	1:B:234:PRO:HD2	2.01	0.42
1:B:477:TYR:HH	2:B:500:HEM:CGD	2.33	0.41
6:B:800:J11:N11	6:B:800:J11:H5'A	2.35	0.41
5:B:880:GOL:H2	9:B:1148:HOH:O	2.20	0.41
1:A:252:ARG:HA	1:A:252:ARG:HD2	1.90	0.40
2:B:500:HEM:CMC	2:B:500:HEM:HBC2	2.51	0.40
1:B:150:GLU:O	1:B:154:GLU:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/444 (90%)	390 (98%)	8 (2%)	2 (0%)	29	28
1	B	398/444 (90%)	391 (98%)	6 (2%)	1 (0%)	41	44
All	All	798/888 (90%)	781 (98%)	14 (2%)	3 (0%)	41	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	260	ASP
1	A	260[A]	ASP
1	A	260[B]	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/377 (91%)	324 (94%)	19 (6%)	21	20
1	B	342/377 (91%)	326 (95%)	16 (5%)	26	27
All	All	685/754 (91%)	650 (95%)	35 (5%)	24	23

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	72	ARG
1	A	91	GLN
1	A	100	ARG
1	A	102	LEU
1	A	146	GLN
1	A	156	GLU
1	A	194	LYS
1	A	257	ARG
1	A	259	GLN
1	A	260[A]	ASP
1	A	260[B]	ASP
1	A	293	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	294	LEU
1	A	328	LEU
1	A	340	ASN
1	A	378	LEU
1	A	390	ARG
1	A	416	LYS
1	B	72	ARG
1	B	92	GLN
1	B	100	ARG
1	B	102	LEU
1	B	105	LEU
1	B	143	SER
1	B	207	GLN
1	B	218	LYS
1	B	225	ASN
1	B	260	ASP
1	B	278	GLN
1	B	330	LEU
1	B	340	ASN
1	B	378	LEU
1	B	414	LEU
1	B	458	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	191	GLN
1	A	259	GLN
1	A	278	GLN
1	A	340	ASN
1	A	376	ASN
1	A	468	ASN
1	B	124	GLN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 1 is monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ACT	A	860	-	3,3,3	0.82	0	3,3,3	0.61	0
7	CAD	A	950	-	0,2,4	-	-	0,1,6	-	-
4	ACT	B	860	-	3,3,3	0.79	0	3,3,3	0.72	0
2	HEM	B	500	1	41,50,50	1.93	7 (17%)	45,82,82	1.67	9 (20%)
3	H4B	B	600	-	16,18,18	0.91	0	11,26,26	2.61	6 (54%)
2	HEM	A	500	1	41,50,50	1.81	6 (14%)	45,82,82	1.51	9 (20%)
5	GOL	B	880	-	5,5,5	0.32	0	5,5,5	0.87	0
3	H4B	A	600	-	16,18,18	0.90	0	11,26,26	2.70	6 (54%)
7	CAD	B	950	-	0,2,4	-	-	0,1,6	-	-
5	GOL	A	880	-	5,5,5	0.35	0	5,5,5	0.57	0
6	J11	B	800	-	27,29,29	0.68	0	30,38,38	1.85	6 (20%)
6	J11	A	800	-	27,29,29	0.72	0	30,38,38	1.61	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	500	1	-	3/12/54/54	-
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
2	HEM	A	500	1	-	4/12/54/54	-
5	GOL	B	880	-	-	2/4/4/4	-
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
5	GOL	A	880	-	-	1/4/4/4	-
6	J11	B	800	-	-	7/13/23/23	0/3/3/3
6	J11	A	800	-	-	5/13/23/23	0/3/3/3

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C3D-C2D	7.44	1.52	1.36
2	A	500	HEM	C3D-C2D	6.65	1.50	1.36
2	B	500	HEM	C3C-C2C	-4.35	1.34	1.40
2	A	500	HEM	C3C-C2C	-3.82	1.35	1.40
2	A	500	HEM	C3C-CAC	3.67	1.55	1.47
2	B	500	HEM	C3C-CAC	3.61	1.55	1.47
2	B	500	HEM	CAB-C3B	3.08	1.55	1.47
2	B	500	HEM	FE-ND	3.04	2.11	1.96
2	A	500	HEM	CAB-C3B	2.90	1.55	1.47
2	B	500	HEM	CMD-C2D	2.15	1.55	1.50
2	B	500	HEM	CMB-C2B	2.09	1.55	1.50
2	A	500	HEM	C3B-C2B	-2.04	1.33	1.37
2	A	500	HEM	CAA-C2A	2.00	1.55	1.52

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	800	J11	C1-N1-C3'	5.47	121.94	114.20
2	B	500	HEM	C4D-ND-C1D	5.02	110.25	105.07
6	B	800	J11	C61-N11-C21	4.93	121.83	118.10
3	B	600	H4B	C4-C4A-N5	4.72	123.08	119.12
2	A	500	HEM	C4D-ND-C1D	4.38	109.60	105.07
3	A	600	H4B	C8A-C4A-C4	4.34	118.43	114.57
3	A	600	H4B	C4-C4A-N5	4.29	122.72	119.12
3	B	600	H4B	C8A-C4A-C4	4.13	118.23	114.57
6	A	800	J11	C61-N11-C21	3.61	120.84	118.10
3	A	600	H4B	C2-N3-C4	3.43	121.38	115.93
2	B	500	HEM	C4B-CHC-C1C	3.43	127.08	122.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	800	J11	C5'-N1'-C2'	3.40	113.45	105.42
3	A	600	H4B	N1-C2-N3	-3.29	120.25	125.42
6	A	800	J11	C1-N1-C3'	3.19	118.72	114.20
3	B	600	H4B	N1-C2-N3	-3.11	120.54	125.42
6	B	800	J11	C5'-N1'-C2'	3.04	112.61	105.42
2	B	500	HEM	CBA-CAA-C2A	-2.98	107.54	112.62
2	B	500	HEM	C1B-NB-C4B	2.95	108.12	105.07
3	B	600	H4B	C2-N3-C4	2.90	120.53	115.93
2	B	500	HEM	C4C-CHD-C1D	2.86	126.34	122.56
2	A	500	HEM	CMA-C3A-C4A	-2.85	124.08	128.46
2	A	500	HEM	CMD-C2D-C1D	2.72	129.18	125.04
6	A	800	J11	C14-C13-C12	-2.70	119.78	123.29
3	B	600	H4B	C2-N1-C8A	2.70	120.59	114.54
6	B	800	J11	C3-C4-C11	-2.70	106.62	112.87
2	B	500	HEM	C3B-C2B-C1B	2.64	108.45	106.49
3	A	600	H4B	C2-N1-C8A	2.61	120.39	114.54
2	A	500	HEM	C4C-CHD-C1D	2.50	125.86	122.56
2	B	500	HEM	CAD-C3D-C4D	2.48	129.00	124.66
2	A	500	HEM	CHB-C1B-NB	2.47	127.44	124.38
6	B	800	J11	C14-C13-C12	-2.47	120.08	123.29
2	A	500	HEM	CAD-C3D-C4D	2.45	128.93	124.66
6	A	800	J11	C2-C1-N1	2.42	115.01	111.06
2	A	500	HEM	C4B-CHC-C1C	2.42	125.75	122.56
3	A	600	H4B	N2-C2-N3	2.41	120.99	117.25
6	B	800	J11	C41-C31-C21	-2.35	118.78	120.32
6	A	800	J11	C4-C11-C12	-2.32	116.73	120.54
2	B	500	HEM	CAD-CBD-CGD	-2.24	108.78	113.60
6	A	800	J11	C3-N2-C2	2.20	123.81	113.45
2	B	500	HEM	CMD-C2D-C1D	2.20	128.38	125.04
3	B	600	H4B	N2-C2-N3	2.18	120.64	117.25
2	A	500	HEM	CAD-CBD-CGD	-2.11	109.05	113.60
2	A	500	HEM	CBA-CAA-C2A	-2.10	109.03	112.62
6	A	800	J11	F13-C13-C14	2.06	122.03	118.54

There are no chirality outliers.

All (22) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	HEM	C2A-CAA-CBA-CGA
2	B	500	HEM	C2A-CAA-CBA-CGA
5	B	880	GOL	C1-C2-C3-O3
6	A	800	J11	C4-C3-N2-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	800	J11	C2'-C3'-N1-C1
6	B	800	J11	C4'-C3'-N1-C1
6	B	800	J11	N2-C3-C4-C11
6	B	800	J11	N1-C1-C2-N2
5	B	880	GOL	O2-C2-C3-O3
6	B	800	J11	C2-C1-N1-C3'
6	B	800	J11	C1-C2-N2-C3
2	A	500	HEM	C3D-CAD-CBD-CGD
6	A	800	J11	C16-C11-C4-C3
6	A	800	J11	C12-C11-C4-C3
6	A	800	J11	C2'-C3'-N1-C1
2	A	500	HEM	CAD-CBD-CGD-O2D
6	A	800	J11	N1-C1-C2-N2
2	A	500	HEM	CAD-CBD-CGD-O1D
6	B	800	J11	C4-C3-N2-C2
2	B	500	HEM	CAD-CBD-CGD-O2D
5	A	880	GOL	C1-C2-C3-O3
2	B	500	HEM	CAD-CBD-CGD-O1D


There are no ring outliers.


8 monomers are involved in 12 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	A	950	CAD	2	0
2	B	500	HEM	4	0
3	B	600	H4B	1	0
2	A	500	HEM	1	0
5	B	880	GOL	1	0
3	A	600	H4B	1	0
7	B	950	CAD	1	0
6	B	800	J11	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	403/444 (90%)	-0.02	13 (3%) 47 47	22, 33, 57, 73	0
1	B	402/444 (90%)	-0.25	8 (1%) 65 66	21, 36, 57, 75	0
All	All	805/888 (90%)	-0.14	21 (2%) 56 57	21, 34, 57, 75	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	GLY	4.9
1	A	259	GLN	4.5
1	A	69	LYS	4.4
1	B	259	GLN	4.0
1	A	160	ALA	3.9
1	A	123	GLU	3.1
1	B	130	ARG	2.6
1	B	121	PRO	2.5
1	A	91	GLN	2.5
1	B	123	GLU	2.5
1	B	146	GLN	2.4
1	A	126	LEU	2.4
1	A	67	GLY	2.3
1	A	127	SER	2.3
1	B	257	ARG	2.2
1	A	156	GLU	2.2
1	B	124	GLN	2.1
1	B	144	GLY	2.1
1	A	157	ALA	2.1
1	A	451	VAL	2.1
1	A	285	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	J11	A	800	27/27	0.90	0.23	22,36,63,65	0
6	J11	B	800	27/27	0.91	0.20	26,38,61,64	0
5	GOL	A	880	6/6	0.92	0.17	37,39,40,42	0
4	ACT	A	860	4/4	0.95	0.11	39,41,41,42	0
3	H4B	B	600	17/17	0.96	0.17	23,26,27,28	0
5	GOL	B	880	6/6	0.97	0.22	30,33,34,35	0
4	ACT	B	860	4/4	0.97	0.15	40,41,41,42	0
2	HEM	A	500	43/43	0.97	0.18	21,24,40,46	0
3	H4B	A	600	17/17	0.98	0.16	24,27,29,29	0
2	HEM	B	500	43/43	0.98	0.14	22,26,43,48	0
7	CAD	A	950	3/5	0.98	0.09	47,47,48,50	0
7	CAD	B	950	3/5	0.99	0.08	49,49,51,51	0
8	ZN	A	900	1/1	1.00	0.11	26,26,26,26	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around J11 A 800:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around J11 B 800:**


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM A 500:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.