



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 10:13 PM JST

PDB ID : 6JXG  
Title : Crystall Structure of Beta-glucosidase D2-BGL from Chaetomella Raphigera  
Authors : Wang, A.H.-J.; Lee, C.C.; Kao, M.R.; Ho, T.H.D.  
Deposited on : 2019-04-23  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	722	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5856 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-glucosidase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	713	5277	3332	890	1040	15	0	0	0

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0
2	A	1	14	8	1	5	0	0

- Molecule 3 is alpha-D-mannopyranose (three-letter code: MAN) (formula:  $C_6H_{12}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			11	6	5		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	526	Total	O	0	0
			526	526		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Beta-glucosidase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.77Å 77.66Å 123.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.19 – 1.90 24.18 – 1.90	Depositor EDS
% Data completeness (in resolution range)	93.6 (24.19-1.90) 93.7 (24.18-1.90)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.73 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.145 , 0.195 0.157 , 0.200	Depositor DCC
$R_{free}$ test set	2504 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.1	Xtrriage
Anisotropy	0.569	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 45.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5856	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	17.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.98	0/5395	1.01	21/7360 (0.3%)

There are no bond length outliers.

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	169	MET	CG-SD-CE	-16.57	73.69	100.20
1	A	672	ARG	NE-CZ-NH1	11.14	125.87	120.30
1	A	672	ARG	NE-CZ-NH2	-10.55	115.03	120.30
1	A	29	VAL	CG1-CB-CG2	8.67	124.78	110.90
1	A	94	LEU	CB-CG-CD2	7.34	123.48	111.00
1	A	545	VAL	CG1-CB-CG2	7.14	122.32	110.90
1	A	702	ARG	NE-CZ-NH2	-6.90	116.85	120.30
1	A	580	THR	CB-CA-C	-6.69	93.54	111.60
1	A	545	VAL	CA-CB-CG1	5.56	119.24	110.90
1	A	558	ASP	CB-CG-OD1	5.55	123.30	118.30
1	A	214	PHE	CB-CG-CD1	-5.49	116.96	120.80
1	A	64	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	167	ASP	CB-CG-OD2	5.43	123.19	118.30
1	A	219	LEU	CB-CG-CD1	5.36	120.12	111.00
1	A	347	LYS	CD-CE-NZ	5.32	123.93	111.70
1	A	167	ASP	CB-CG-OD1	-5.29	113.54	118.30
1	A	448	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	29	VAL	N-CA-CB	-5.18	100.09	111.50
1	A	702	ARG	NE-CZ-NH1	5.12	122.86	120.30
1	A	587	LEU	CA-CB-CG	5.04	126.89	115.30
1	A	558	ASP	CB-CG-OD2	-5.03	113.77	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5277	0	5141	29	0
2	A	42	0	39	2	0
3	A	11	0	10	0	0
4	A	526	0	0	7	2
All	All	5856	0	5190	30	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:566:ASN:O	4:A:901:HOH:O	1.84	0.93
1:A:580:THR:HG21	1:A:622:ASN:OD1	1.82	0.80
1:A:580:THR:HG22	1:A:624:GLY:HA3	1.66	0.77
1:A:334:ILE:HD12	1:A:518:ILE:HD13	1.73	0.70
1:A:595:SER:OG	4:A:902:HOH:O	1.92	0.70
1:A:55:CYS:H	1:A:104:ASN:HD22	1.39	0.68
1:A:39:CYS:H	1:A:42:ASN:HD21	1.38	0.68
1:A:598:SER:OG	1:A:600:THR:HG23	1.95	0.67
1:A:409:THR:HG21	4:A:1073:HOH:O	1.94	0.66
1:A:175:ASP:OD2	2:A:803:NAG:O3	2.06	0.61
1:A:711:LYS:CD	1:A:711:LYS:H	2.16	0.58
1:A:455:ASN:C	1:A:455:ASN:HD22	2.08	0.57
1:A:349:PRO:O	1:A:409:THR:HB	2.05	0.57
1:A:295:VAL:HG23	1:A:297:GLN:HG3	1.87	0.57
1:A:711:LYS:H	1:A:711:LYS:HD3	1.70	0.56
1:A:29:VAL:HG13	1:A:263:TRP:CE3	2.42	0.55
1:A:462:ASN:ND2	4:A:907:HOH:O	2.39	0.54
1:A:116:LYS:HE3	1:A:117:ILE:HG13	1.90	0.53
2:A:801:NAG:C6	4:A:973:HOH:O	2.58	0.52
1:A:409:THR:CG2	4:A:1073:HOH:O	2.56	0.50
1:A:224:ASP:HB2	1:A:601:LEU:HD12	1.97	0.47
1:A:383:TRP:CD1	1:A:441:GLY:HA3	2.51	0.45
1:A:564:GLN:HE21	1:A:647:PRO:HD3	1.81	0.44

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:475:LYS:O	1:A:499:LYS:HE2	2.17	0.44
1:A:580:THR:HG22	1:A:624:GLY:CA	2.43	0.43
1:A:659:LYS:HB2	1:A:662:LYS:HG3	2.00	0.43
1:A:656:ILE:HD11	1:A:664:GLY:HA3	2.01	0.43
1:A:23:THR:HG23	4:A:1287:HOH:O	2.19	0.42
1:A:198:MET:HA	1:A:230:VAL:O	2.20	0.41
1:A:292:TRP:CZ2	1:A:298:ASP:HA	2.56	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1329:HOH:O	4:A:1363:HOH:O[3_644]	1.91	0.29
4:A:1326:HOH:O	4:A:1339:HOH:O[3_554]	2.12	0.08

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	711/722 (98%)	684 (96%)	25 (4%)	2 (0%)	41 31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	341	ASN
1	A	128	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	550/555 (99%)	526 (96%)	24 (4%)	28 19

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	VAL
1	A	36	LYS
1	A	94	LEU
1	A	116	LYS
1	A	214	PHE
1	A	219	LEU
1	A	233	TRP
1	A	299	GLN
1	A	341	ASN
1	A	347	LYS
1	A	409	THR
1	A	451	LEU
1	A	455	ASN
1	A	545	VAL
1	A	580	THR
1	A	587	LEU
1	A	591	TYR
1	A	600	THR
1	A	601	LEU
1	A	634	LEU
1	A	638	LEU
1	A	672	ARG
1	A	689	THR
1	A	711	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	42	ASN
1	A	104	ASN
1	A	151	GLN
1	A	236	GLN
1	A	455	ASN

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type
1	A	462	ASN
1	A	536	GLN
1	A	564	GLN
1	A	568	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	802	1	14,14,15	0.91	1 (7%)	17,19,21	1.19	2 (11%)
2	NAG	A	803	1	14,14,15	0.87	1 (7%)	17,19,21	3.27	10 (58%)
3	MAN	A	804	1	11,11,12	1.09	2 (18%)	15,15,17	2.06	5 (33%)
2	NAG	A	801	1	14,14,15	0.55	0	17,19,21	1.47	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	802	1	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1	-	2/6/23/26	0/1/1/1
3	MAN	A	804	1	-	0/2/19/22	0/1/1/1
2	NAG	A	801	1	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	802	NAG	C1-C2	2.44	1.56	1.52
2	A	803	NAG	O4-C4	-2.12	1.38	1.43
3	A	804	MAN	O2-C2	-2.11	1.38	1.43
3	A	804	MAN	O5-C1	-2.07	1.40	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	803	NAG	C1-C2-N2	5.99	120.72	110.49
2	A	803	NAG	C3-C4-C5	-5.32	100.75	110.24
3	A	804	MAN	C1-C2-C3	4.87	115.66	109.67
2	A	803	NAG	O5-C5-C4	-4.75	99.27	110.83
2	A	803	NAG	C8-C7-N2	4.58	123.86	116.10
2	A	803	NAG	O7-C7-C8	-4.22	114.22	122.06
2	A	803	NAG	O5-C5-C6	3.75	113.08	107.20
2	A	803	NAG	O4-C4-C5	3.63	118.30	109.30
3	A	804	MAN	C1-O5-C5	3.35	116.74	112.19
3	A	804	MAN	C3-C4-C5	-2.96	104.96	110.24
2	A	803	NAG	O6-C6-C5	-2.95	101.16	111.29
2	A	801	NAG	C3-C4-C5	-2.93	105.02	110.24
2	A	803	NAG	C4-C3-C2	-2.86	106.83	111.02
2	A	802	NAG	O4-C4-C3	-2.81	103.86	110.35
2	A	803	NAG	C2-N2-C7	2.64	126.66	122.90
3	A	804	MAN	O2-C2-C1	-2.46	104.12	109.15
2	A	802	NAG	C1-C2-N2	2.35	114.51	110.49
2	A	801	NAG	O6-C6-C5	-2.33	103.31	111.29
3	A	804	MAN	O3-C3-C2	-2.31	105.58	109.99
2	A	801	NAG	O5-C1-C2	-2.23	107.77	111.29
2	A	801	NAG	C4-C3-C2	-2.23	107.76	111.02
2	A	801	NAG	O5-C5-C4	-2.04	105.85	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	803	NAG	C8-C7-N2-C2
2	A	803	NAG	O7-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	803	NAG	1	0
2	A	801	NAG	1	0

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	A	802	1	14,14,15	0.91	1 (7%)	17,19,21	1.19	2 (11%)
2	NAG	A	803	1	14,14,15	0.87	1 (7%)	17,19,21	3.27	10 (58%)
3	MAN	A	804	1	11,11,12	1.09	2 (18%)	15,15,17	2.06	5 (33%)
2	NAG	A	801	1	14,14,15	0.55	0	17,19,21	1.47	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	802	1	-	0/6/23/26	0/1/1/1
2	NAG	A	803	1	-	2/6/23/26	0/1/1/1
3	MAN	A	804	1	-	0/2/19/22	0/1/1/1
2	NAG	A	801	1	-	0/6/23/26	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	802	NAG	C1-C2	2.44	1.56	1.52

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	803	NAG	O4-C4	-2.12	1.38	1.43
3	A	804	MAN	O2-C2	-2.11	1.38	1.43
3	A	804	MAN	O5-C1	-2.07	1.40	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	803	NAG	C1-C2-N2	5.99	120.72	110.49
2	A	803	NAG	C3-C4-C5	-5.32	100.75	110.24
3	A	804	MAN	C1-C2-C3	4.87	115.66	109.67
2	A	803	NAG	O5-C5-C4	-4.75	99.27	110.83
2	A	803	NAG	C8-C7-N2	4.58	123.86	116.10
2	A	803	NAG	O7-C7-C8	-4.22	114.22	122.06
2	A	803	NAG	O5-C5-C6	3.75	113.08	107.20
2	A	803	NAG	O4-C4-C5	3.63	118.30	109.30
3	A	804	MAN	C1-O5-C5	3.35	116.74	112.19
3	A	804	MAN	C3-C4-C5	-2.96	104.96	110.24
2	A	803	NAG	O6-C6-C5	-2.95	101.16	111.29
2	A	801	NAG	C3-C4-C5	-2.93	105.02	110.24
2	A	803	NAG	C4-C3-C2	-2.86	106.83	111.02
2	A	802	NAG	O4-C4-C3	-2.81	103.86	110.35
2	A	803	NAG	C2-N2-C7	2.64	126.66	122.90
3	A	804	MAN	O2-C2-C1	-2.46	104.12	109.15
2	A	802	NAG	C1-C2-N2	2.35	114.51	110.49
2	A	801	NAG	O6-C6-C5	-2.33	103.31	111.29
3	A	804	MAN	O3-C3-C2	-2.31	105.58	109.99
2	A	801	NAG	O5-C1-C2	-2.23	107.77	111.29
2	A	801	NAG	C4-C3-C2	-2.23	107.76	111.02
2	A	801	NAG	O5-C5-C4	-2.04	105.85	110.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	803	NAG	C8-C7-N2-C2
2	A	803	NAG	O7-C7-N2-C2

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	803	NAG	1	0
2	A	801	NAG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	713/722 (98%)	-0.28	14 (1%) 65 68	9, 15, 30, 62	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	713	SER	4.2
1	A	596	ALA	3.8
1	A	691	GLY	3.3
1	A	680	THR	3.0
1	A	1	PRO	3.0
1	A	690	SER	3.0
1	A	406	ALA	2.9
1	A	4	GLY	2.7
1	A	408	GLY	2.7
1	A	348	LYS	2.5
1	A	299	GLN	2.4
1	A	594	THR	2.3
1	A	595	SER	2.2
1	A	409	THR	2.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	A	803	14/15	0.89	0.23	25,33,40,44	0
2	NAG	A	801	14/15	0.92	0.20	27,35,40,43	0
3	MAN	A	804	11/12	0.95	0.16	23,28,34,34	0

*Continued on next page...*



*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	802	14/15	0.96	0.13	19,22,26,26	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	NAG	A	803	14/15	0.89	0.23	25,33,40,44	0
2	NAG	A	801	14/15	0.92	0.20	27,35,40,43	0
3	MAN	A	804	11/12	0.95	0.16	23,28,34,34	0
2	NAG	A	802	14/15	0.96	0.13	19,22,26,26	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.