



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 10, 2023 – 09:26 PM EDT

PDB ID : 4K3V
Title : Structure of Staphylococcus aureus MntC
Authors : Parris, K.D.; Mosyak, L.
Deposited on : 2013-04-11
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	 3% 75% 7% 15%
1	B	309	 2% 81% 8% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4404 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC superfamily ATP binding cassette transporter, binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	253	2027	1288	340	393	6	0	0	0
1	B	278	2221	1406	371	437	7	0	0	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	A	1	1	1	0	0
2	B	1	1	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	62	62	62	0	0
3	B	92	92	92	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	67.48Å 68.36Å 107.91Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.00 – 2.20 22.30 – 2.20	Depositor EDS
% Data completeness (in resolution range)	79.0 (23.00-2.20) 79.0 (22.30-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.19 (at 2.19Å)	Xtrriage
Refinement program	BUSTER 2.11.1	Depositor
R, R_{free}	0.198 , 0.219 0.204 , 0.234	Depositor DCC
R_{free} test set	1039 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtrriage
Anisotropy	0.828	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.096 for k,h,-l	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4404	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2064	0.71	2/2772 (0.1%)
1	B	0.46	0/2261	0.67	0/3035
All	All	0.47	0/4325	0.69	2/5807 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	178	ILE	N-CA-C	-7.39	91.04	111.00
1	A	177	ASP	C-N-CA	5.04	134.31	121.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2027	0	2031	10	0
1	B	2221	0	2224	17	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	62	0	0	0	0
3	B	92	0	0	3	0
All	All	4404	0	4255	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:263:THR:HG23	1:B:264:ASP:H	1.39	0.87
1:A:186:ILE:HG21	1:A:235:LEU:HD23	1.65	0.79
1:A:186:ILE:CG2	1:A:235:LEU:HD23	2.20	0.71
1:B:263:THR:HG21	3:B:419:HOH:O	1.96	0.64
1:A:25:LEU:HD22	1:A:101:VAL:HG21	1.78	0.64
1:A:187:THR:HG23	1:A:189:GLU:O	1.97	0.64
1:B:25:LEU:CD2	1:B:101:VAL:HG21	2.28	0.63
1:A:180:LYS:HA	1:A:183:ARG:HD3	1.84	0.59
1:B:25:LEU:HD22	1:B:101:VAL:HG21	1.86	0.57
1:B:263:THR:HG23	1:B:264:ASP:N	2.16	0.54
1:A:25:LEU:CD2	1:A:101:VAL:HG21	2.37	0.54
1:B:54:VAL:HG13	1:B:58:ASP:HB2	1.88	0.53
1:B:166:LYS:HG3	3:B:488:HOH:O	2.09	0.52
1:B:25:LEU:HD21	1:B:101:VAL:HG21	1.92	0.52
1:B:238:GLU:HG2	1:B:241:VAL:HG22	1.92	0.51
1:A:31:ASN:O	1:A:156:GLN:HG3	2.15	0.47
1:B:209:ILE:O	1:B:212:GLU:HG2	2.15	0.47
1:B:143:ILE:HG23	1:B:150:LYS:HB2	1.97	0.47
1:B:222:GLN:NE2	3:B:417:HOH:O	2.48	0.46
1:A:83:GLU:HG3	1:A:93:LEU:HD11	1.99	0.45
1:A:143:ILE:HG23	1:A:150:LYS:HB2	1.99	0.44
1:A:210:ASN:HA	1:A:214:GLN:HE21	1.82	0.44
1:B:255:LYS:HD3	1:B:255:LYS:HA	1.89	0.43
1:B:110:LEU:HD22	1:B:208:GLU:HG2	2.01	0.42
1:B:48:ASP:OD1	1:B:50:HIS:HD2	2.03	0.41
1:B:271:THR:HG23	1:B:273:GLY:H	1.85	0.41
1:B:237:VAL:O	1:B:260:GLU:HA	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/309 (80%)	236 (96%)	9 (4%)	2 (1%)	19	19
1	B	276/309 (89%)	267 (97%)	7 (2%)	2 (1%)	22	22
All	All	523/618 (85%)	503 (96%)	16 (3%)	4 (1%)	19	19

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	HIS
1	B	123	HIS
1	A	78	GLY
1	B	78	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/270 (83%)	220 (99%)	3 (1%)	69	81
1	B	245/270 (91%)	241 (98%)	4 (2%)	62	76
All	All	468/540 (87%)	461 (98%)	7 (2%)	65	78

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150	LYS
1	A	154	GLU
1	A	172	LYS
1	B	150	LYS
1	B	174	LYS
1	B	197	LYS
1	B	271	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	198	GLN
1	A	214	GLN
1	A	234	HIS
1	B	79	ASN
1	B	111	ASN
1	B	222	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	253/309 (81%)	0.29	8 (3%) 47 45	18, 40, 71, 89	0
1	B	278/309 (89%)	0.02	6 (2%) 62 59	16, 35, 60, 73	0
All	All	531/618 (85%)	0.15	14 (2%) 56 53	16, 37, 66, 89	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	233	LYS	5.0
1	A	235	LEU	3.7
1	B	114	GLU	3.3
1	A	118	ASP	2.9
1	B	225	GLU	2.6
1	A	226	PHE	2.6
1	A	217	PRO	2.5
1	B	112	GLY	2.4
1	B	117	LYS	2.4
1	B	173	ASP	2.3
1	A	221	ARG	2.2
1	A	178	ILE	2.1
1	A	270	GLY	2.1
1	B	115	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MN	A	301	1/1	0.99	0.05	30,30,30,30	0
2	MN	B	301	1/1	1.00	0.09	20,20,20,20	0

6.5 Other polymers [i](#)

There are no such residues in this entry.