

wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 01:19 pm BST

PDB ID : 5K57 BMRB ID : 30097

Title : HDD domain from human Ddi2

Authors : Veverka, V. Deposited on : 2016-05-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &: & v1.2 \\ BMRB \ Restraints \ Analysis &: & v1.2 \\ \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 93%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	NMR archive
Metric	$(\# \mathrm{Entries})$	$(\# \mathrm{Entries})$
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain				
1	A	98	43%		56%		



2 Ensemble composition and analysis (i)

This entry contains 30 models. Model 13 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues					
Well-defined core Residue range (total) Backbone RMSD (Å) Medoid mode					
1	A:21-A:63 (43)	0.19	13		

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 4, 7, 10, 12, 13, 15, 17, 21, 22, 25, 27, 29
2	3, 6, 9, 14, 16, 18, 24, 28, 30
3	2, 5, 8, 11, 20, 26
4	19, 23



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1557 atoms, of which 775 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called Protein DDI1 homolog 2.

Mol	Chain	Residues	Atoms					Trace	
1	Λ	08	Total	С	Н	N	О	S	0
1 A	98	1557	479	775	144	158	1	0	

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP Q5TDH0




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: Protein DDI1 homolog 2



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 13. Colouring as in section 4.1 above.

• Molecule 1: Protein DDI1 homolog 2





Refinement protocol and experimental data overview (i) 5



The models were refined using the following method: simulated annealing.

Of the 100 calculated structures, 30 were deposited, based on the following criterion: structures with the lowest energy.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
YASARA	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1190
Number of shifts mapped to atoms	1190
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	93%



6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		В	Sond lengths	Bond angles		
		RMSZ	#Z>5	RMSZ	#Z>5	
1	A	0.60 ± 0.03	$0\pm0/343~(~0.0\pm~0.0\%)$	0.89 ± 0.04	$1\pm1/465~(~0.3\pm~0.2\%)$	
All	All	0.60	0/10290 (0.0%)	0.89	36/13950 (0.3%)	

There are no bond-length outliers.

5 of 6 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	hain Res Type Atoms Z Observed(Observed(0)	Ideal(0)	Models			
MIOI	Chain	nes	Type	Atoms	Z Obser	$\operatorname{Observed}(^{o})$	$\mathbf{Ideal}(^o)$	Worst	Total
1	A	25	ARG	NE-CZ-NH1	7.38	123.99	120.30	10	11
1	A	25	ARG	NE-CZ-NH2	-5.92	117.34	120.30	10	1
1	A	59	ARG	NE-CZ-NH1	5.77	123.18	120.30	11	4
1	A	37	LEU	CA-CB-CG	5.68	128.37	115.30	23	14
1	A	41	ARG	NE-CZ-NH1	5.36	122.98	120.30	3	5

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	337	356	356	0±1
All	All	10110	10680	10680	14

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.



All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom 2	Clash(Å)	$Distance(\mathring{A})$	${f Models}$	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:21:PRO:HB3	1:A:61:LEU:HD22	0.52	1.82	18	13
1:A:48:ALA:HB1	1:A:57:PHE:HA	0.41	1.92	23	1

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	sed Favoured Allowed		Outliers	Percentiles		
1	A	43/98 (44%)	43±0 (99±1%)	0±0 (1±1%)	0±0 (0±0%)	100 100		
All	All	1290/2940 (44%)	1279 (99%)	11 (1%)	0 (0%)	100 100		

There are no Ramachandran outliers.

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric		Percentiles		
1	A	38/87 (44%)	35±1 (93±3%)	3±1 (7±3%)	19	68	
All	All	1140/2610 (44%)	1063 (93%)	77 (7%)	19	68	

5 of 11 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	35	LEU	30
1	A	59	ARG	10
1	A	26	ASP	9
1	A	34	GLU	9
1	A	53	ASP	5



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 93% for the well-defined parts and 87% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: hdd.star

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1190
Number of shifts mapped to atoms	1190
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\mathrm{C}_{\alpha}$	97	-0.20 ± 0.11	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	94	0.12 ± 0.07	None needed ($< 0.5 \text{ ppm}$)
¹³ C'	96	-0.23 ± 0.16	None needed (< 0.5 ppm)
^{15}N	89	-0.34 ± 0.32	None needed ($< 0.5 \text{ ppm}$)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 93%, i.e. 573 atoms were assigned a chemical shift out of a possible 615. 0 out of 14 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	207/208 (100%)	83/83 (100%)	85/86~(99%)	39/39 (100%)
Sidechain	357/390 (92%)	244/256 (95%)	108/121 (89%)	5/13 (38%)



Continued from previous page...

	Total	$^{1}\mathbf{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	9/17 (53%)	6/9 (67%)	3/7~(43%)	0/1 (0%)
Overall	573/615 (93%)	333/348 (96%)	$196/214 \ (92\%)$	44/53 (83%)


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1917
Intra-residue ($ i-j =0$)	495
Sequential (i-j =1)	536
Medium range ($ i-j >1$ and $ i-j <5$)	593
Long range (i-j ≥5)	293
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	108
Number of unmapped restraints	0
Number of restraints per residue	20.7
Number of long range restraints per residue ¹	3.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	8.2	0.2
0.2-0.5 (Medium)	3.6	0.5
>0.5 (Large)	11.6	1.79



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins (°)	Average number of violations per model	$\mathbf{Max} \ (^{\circ})$
1.0-10.0 (Small)	3.2	5.1
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Dordensinda dom o	C	% ¹	Vic	olated ⁵	3	Consis	tentl	${f y}$ Violated 4
Restraints type	Count	70	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	495	25.8	13	2.6	0.7	2	0.4	0.1
Backbone-Backbone	5	0.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	410	21.4	13	3.2	0.7	2	0.5	0.1
Sidechain-Sidechain	80	4.2	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	536	28.0	16	3.0	0.8	1	0.2	0.1
Backbone-Backbone	103	5.4	2	1.9	0.1	0	0.0	0.0
Backbone-Sidechain	387	20.2	12	3.1	0.6	0	0.0	0.0
Sidechain-Sidechain	46	2.4	2	4.3	0.1	1	2.2	0.1
Medium range ($ i-j >1 \& i-j <5$)	593	30.9	35	5.9	1.8	3	0.5	0.2
Backbone-Backbone	147	7.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	286	14.9	15	5.2	0.8	2	0.7	0.1
Sidechain-Sidechain	160	8.3	20	12.5	1.0	1	0.6	0.1
Long range ($ i-j \ge 5$)	293	15.3	15	5.1	0.8	0	0.0	0.0
Backbone-Backbone	6	0.3	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	128	6.7	7	5.5	0.4	0	0.0	0.0
Sidechain-Sidechain	159	8.3	8	5.0	0.4	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1917	100.0	79	4.1	4.1	6	0.3	0.3
Backbone-Backbone	261	13.6	2	0.8	0.1	0	0.0	0.0
Backbone-Sidechain	1211	63.2	47	3.9	2.5	4	0.3	0.2
Sidechain-Sidechain	445	23.2	30	6.7	1.6	2	0.4	0.1

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID	Number of violations					5	Mean (Å)	Morr (Å)	${ m SD}^6$ (Å)	Modion (Å)	
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (Å)	$SD^*(A)$	Median (Å)	
1	7	2	8	6	0	23	0.59	1.75	0.45	0.64	
2	8	4	15	7	0	34	0.47	1.77	0.44	0.2	
3	5	1	6	6	0	18	0.73	1.75	0.46	0.76	
4	9	3	9	6	0	27	0.53	1.75	0.44	0.32	
5	8	5	8	7	0	28	0.51	1.77	0.46	0.19	
6	7	1	6	6	0	20	0.68	1.75	0.45	0.78	
7	7	5	11	5	0	28	0.52	1.75	0.45	0.29	
8	8	3	10	7	0	28	0.51	1.75	0.44	0.34	
9	6	1	11	3	0	21	0.7	1.76	0.52	0.78	
10	6	2	7	6	0	21	0.64	1.76	0.44	0.74	
11	8	2	9	5	0	24	0.59	1.76	0.45	0.52	




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	6	3	8	3	0	20	0.62	1.75	0.48	0.52
13	7	2	11	5	0	25	0.57	1.74	0.44	0.41
14	4	1	6	4	0	15	0.81	1.75	0.45	0.83
15	7	5	9	6	0	27	0.56	1.76	0.45	0.34
16	8	1	9	6	0	24	0.56	1.73	0.42	0.35
17	7	4	10	10	0	31	0.49	1.75	0.43	0.29
18	4	1	5	3	0	13	0.87	1.76	0.37	0.8
19	7	1	5	4	0	17	0.68	1.74	0.44	0.76
20	9	4	14	7	0	34	0.46	1.79	0.41	0.23
21	7	5	13	6	0	31	0.51	1.74	0.43	0.33
22	6	1	7	3	0	17	0.66	1.75	0.49	0.74
23	4	1	6	3	0	14	0.8	1.72	0.37	0.77
24	8	6	11	7	0	32	0.49	1.67	0.45	0.22
25	4	1	6	3	0	14	0.89	1.75	0.39	0.86
26	8	1	6	3	0	18	0.71	1.75	0.5	0.8
27	7	3	10	7	0	27	0.5	1.74	0.46	0.28
28	4	1	5	4	0	14	0.91	1.74	0.36	0.86
29	9	5	12	5	0	31	0.5	1.78	0.45	0.24
30	8	2	6	7	0	23	0.6	1.76	0.47	0.56

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation



9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1838(IR:482, SQ:520, MR:558, LR:278, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	restra	aints	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Count ⁶	%		
4	7	15	3	0	29	1	3.3		
0	2	4	1	0	7	2	6.7		
0	3	3	2	0	8	3	10.0		
1	0	1	0	0	2	4	13.3		
0	1	0	1	0	2	5	16.7		
0	0	0	0	0	0	6	20.0		




Continued from previous page...

	,	$\frac{1}{\text{of vio}}$				Fraction	n of the ensemble
IR^1	SQ^2	$ m MR^3$	LR^4	IC^5	Total	Count ⁶	%
0	1	0	0	0	1	7	23.3
0	0	0	2	0	2	8	26.7
0	0	1	1	0	2	9	30.0
1	0	1	0	0	2	10	33.3
0	0	0	0	0	0	11	36.7
0	0	2	1	0	3	12	40.0
0	0	2	0	0	2	13	43.3
0	0	0	0	0	0	14	46.7
0	1	0	0	0	1	15	50.0
0	0	0	0	0	0	16	53.3
0	0	0	0	0	0	17	56.7
0	0	0	0	0	0	18	60.0
1	0	2	0	0	3	19	63.3
0	0	0	0	0	0	20	66.7
0	0	0	0	0	0	21	70.0
0	0	0	0	0	0	22	73.3
0	0	0	0	0	0	23	76.7
1	0	0	1	0	2	24	80.0
1	0	0	0	0	1	25	83.3
0	0	1	0	0	1	26	86.7
0	0	0	2	0	2	27	90.0
1	0	0	0	0	1	28	93.3
1	0	0	1	0	2	29	96.7
2	1	3	0	0	6	30	100.0

 $^{^1{\}rm Intra-residue}$ restraints, $^2{\rm Sequential}$ restraints, $^3{\rm Medium}$ range restraints, $^4{\rm Long}$ range restraints, $^5{\rm Inter-chain}$ restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	$Models^1$	Mean (Å)	SD^1 (Å)	Median (Å)
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	30	1.75	0.02	1.75
(1,1355)	1:A:27:MET:HB2	1:A:31:ASN:HD21	30	1.32	0.08	1.33
(1,1355)	1:A:27:MET:HB3	1:A:31:ASN:HD21	30	1.32	0.08	1.33
(1,1354)	1:A:30:ALA:HB1	1:A:31:ASN:HD21	30	1.04	0.08	1.04
(1,1354)	1:A:30:ALA:HB2	1:A:31:ASN:HD21	30	1.04	0.08	1.04
(1,1354)	1:A:30:ALA:HB3	1:A:31:ASN:HD21	30	1.04	0.08	1.04
(1,894)	1:A:42:ASN:H	1:A:42:ASN:HD21	30	0.81	0.03	0.82
(1,1350)	1:A:31:ASN:H	1:A:31:ASN:HD21	30	0.78	0.04	0.78
(1,1353)	1:A:28:LEU:HA	1:A:31:ASN:HD21	30	0.72	0.13	0.74
(1,1682)	1:A:45:LEU:HD11	1:A:65:GLN:HE21	29	1.02	0.12	1.02
(1,1682)	1:A:45:LEU:HD12	1:A:65:GLN:HE21	29	1.02	0.12	1.02
(1,1682)	1:A:45:LEU:HD13	1:A:65:GLN:HE21	29	1.02	0.12	1.02
(1,1682)	1:A:45:LEU:HD21	1:A:65:GLN:HE21	29	1.02	0.12	1.02
(1,1682)	1:A:45:LEU:HD22	1:A:65:GLN:HE21	29	1.02	0.12	1.02
(1,1682)	1:A:45:LEU:HD23	1:A:65:GLN:HE21	29	1.02	0.12	1.02
(1,1392)	1:A:65:GLN:HA	1:A:65:GLN:HE21	29	0.92	0.04	0.92



Continued from previous page...


Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	${ m SD}^1\ (m \AA)$	Median (Å)	
(1,1386)	1:A:66:GLN:HA	1:A:66:GLN:HE21	28	0.6	0.04	0.6	
(1,1402)	1:A:42:ASN:HD22	1:A:64:GLN:HE22	27	1.06	0.14	1.08	

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram: Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	20	1.79
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	29	1.78
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	2	1.77



Continued from previous page...

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	5	1.77
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	9	1.76
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	10	1.76
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	11	1.76
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	15	1.76
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	18	1.76
(1,1352)	1:A:27:MET:HA	1:A:31:ASN:HD21	30	1.76



10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle true	Count	$\%^{1}$	${f Violated}^3$			Consistently Violated ⁴		
Angle type	Count	/0	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	% ¹
PSI	54	50.0	9	16.7	8.3	0	0.0	0.0
PHI	54	50.0	10	18.5	9.3	0	0.0	0.0
Total	108	100.0	19	17.6	17.6	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models

10.1.1 Bar chart: Distribution of dihedral-angles and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories



10.2 Dihedral-angle violation statistics for each model (i)


Page 24

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

N/ 1 1 TD	Nun	nber o	of violations	N (0)	N (0)	GD (0)	N 1 1 (0)
Model ID	PSI	PHI	Total	$ Mean (^{\circ}) $	$\mathbf{Max} \ (^{\circ})$	\mathbf{SD} (°)	$ \ \mathbf{Median} (^{\circ}) $
1	1	1	2	3.3	4.9	1.6	3.3
2	5	2	7	1.94	4.0	0.93	1.6
3	2	0	2	4.5	4.9	0.4	4.5
4	3	1	4	3.6	5.1	1.03	3.5
5	6	0	6	2.88	4.7	1.39	2.8
6	1	1	2	4.35	5.0	0.65	4.35
7	2	0	2	4.25	4.8	0.55	4.25
8	3	1	4	3.1	4.9	1.41	3.2
9	2	1	3	2.87	4.9	1.48	2.3
10	1	1	2	4.35	4.6	0.25	4.35
11	2	2	4	3.05	4.8	1.47	3.15
12	0	2	2	3.55	4.7	1.15	3.55
13	2	2	4	2.35	4.7	1.36	1.65
14	0	0	0	0.0	0.0	0.0	0.0
15	5	2	7	2.76	4.8	0.99	2.6
16	3	3	6	2.67	4.8	1.28	2.45
17	2	2	4	3.32	4.6	1.06	3.35
18	1	0	1	1.9	1.9	0.0	1.9
19	0	1	1	4.6	4.6	0.0	4.6
20	3	3	6	2.7	4.9	1.12	2.15
21	3	0	3	2.73	4.4	1.31	2.6
22	1	1	2	3.3	4.6	1.3	3.3
23	0	1	1	4.9	4.9	0.0	4.9
24	3	1	4	3.88	4.6	0.86	4.25
25	1	0	1	5.1	5.1	0.0	5.1
26	1	2	3	4.03	4.7	0.47	3.7
27	2	1	3	3.43	4.9	1.54	4.1
28	0	0	0	0.0	0.0	0.0	0.0
29	5	3	8	2.68	4.7	1.25	2.55
30	2	1	3	3.53	5.0	1.23	3.6



10.2.1 Bar graph: Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Nun	nber o	f violated restraints	Fractio	n of the ensemble
PSI	PHI	Total	Count ¹	%
3	4	7	1	3.3
0	1	1	2	6.7
0	1	1	3	10.0
0	2	2	4	13.3
2	0	2	5	16.7
0	0	0	6	20.0
0	0	0	7	23.3
2	0	2	8	26.7
0	2	2	9	30.0
0	0	0	10	33.3
0	0	0	11	36.7




Continued from previous page...

Nun	nber o	f violated restraints	Fraction	n of the ensemble
PSI	PHI	Total	Count ¹	%
0	0	0	12	40.0
0	0	0	13	43.3
0	0	0	14	46.7
0	0	0	15	50.0
1	0	1	16	53.3
1	0	1	17	56.7
0	0	0	18	60.0
0	0	0	19	63.3
0	0	0	20	66.7
0	0	0	21	70.0
0	0	0	22	73.3
0	0	0	23	76.7
0	0	0	24	80.0
0	0	0	25	83.3
0	0	0	26	86.7
0	0	0	27	90.0
0	0	0	28	93.3
0	0	0	29	96.7
0	0	0	30	100.0

 $^{^{1}}$ Number of models with violations



10.3.1 Bar graph: Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram: Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,30)	1:A:37:LEU:N	1:A:37:LEU:CA	1:A:37:LEU:C	1:A:38:LEU:N	17	3.04	0.88	2.8
(1,68)	1:A:60:VAL:N	1:A:60:VAL:CA	1:A:60:VAL:C	1:A:61:LEU:N	16	3.11	1.54	2.9
(1,95)	1:A:73:GLN:C	1:A:74:GLU:N	1:A:74:GLU:CA	1:A:74:GLU:C	9	4.52	0.38	4.7
(1,107)	1:A:86:LEU:C	1:A:87:GLU:N	1:A:87:GLU:CA	1:A:87:GLU:C	9	4.3	0.55	4.1
(1,100)	1:A:76:ILE:N	1:A:76:ILE:CA	1:A:76:ILE:C	1:A:77:ARG:N	8	3.16	1.15	3.35
(1,92)	1:A:72:GLU:N	1:A:72:GLU:CA	1:A:72:GLU:C	1:A:73:GLN:N	8	3.14	1.37	3.15
(1,36)	1:A:40:GLU:N	1:A:40:GLU:CA	1:A:40:GLU:C	1:A:41:ARG:N	5	1.78	0.54	1.5
(1,26)	1:A:35:LEU:N	1:A:35:LEU:CA	1:A:35:LEU:C	1:A:36:SER:N	5	1.64	0.41	1.6
(1,105)	1:A:85:ASP:C	1:A:86:LEU:N	1:A:86:LEU:CA	1:A:86:LEU:C	4	2.45	1.3	2.05
(1,27)	1:A:35:LEU:C	1:A:36:SER:N	1:A:36:SER:CA	1:A:36:SER:C	4	1.85	0.48	1.95


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram: Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation (°)
(1,92)	1:A:72:GLU:N	1:A:72:GLU:CA	1:A:72:GLU:C	1:A:73:GLN:N	25	5.1
(1,107)	1:A:86:LEU:C	1:A:87:GLU:N	1:A:87:GLU:CA	1:A:87:GLU:C	4	5.1
(1,95)	1:A:73:GLN:C	1:A:74:GLU:N	1:A:74:GLU:CA	1:A:74:GLU:C	6	5.0
(1,68)	1:A:60:VAL:N	1:A:60:VAL:CA	1:A:60:VAL:C	1:A:61:LEU:N	30	5.0
(1,95)	1:A:73:GLN:C	1:A:74:GLU:N	1:A:74:GLU:CA	1:A:74:GLU:C	23	4.9
(1,89)	1:A:70:ARG:C	1:A:71:ARG:N	1:A:71:ARG:CA	1:A:71:ARG:C	1	4.9
(1,68)	1:A:60:VAL:N	1:A:60:VAL:CA	1:A:60:VAL:C	1:A:61:LEU:N	3	4.9
(1,68)	1:A:60:VAL:N	1:A:60:VAL:CA	1:A:60:VAL:C	1:A:61:LEU:N	27	4.9
(1,107)	1:A:86:LEU:C	1:A:87:GLU:N	1:A:87:GLU:CA	1:A:87:GLU:C	9	4.9
(1,107)	1:A:86:LEU:C	1:A:87:GLU:N	1:A:87:GLU:CA	1:A:87:GLU:C	20	4.9

