



wwPDB EM Validation Summary Report ⓘ

Apr 20, 2026 – 08:34 PM JST

PDB ID : 9K53 / pdb_00009k53
EMDB ID : EMD-62079
Title : Structure of substrate-engaged human 26S proteasome RP-CP subcomplex in state ED2.1
Authors : Wu, Z.; Chen, E.; Mao, Y.
Deposited on : 2024-10-21
Resolution : 2.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	7115 (2.00 - 3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	433	<div> <div>14%</div> <div> <div></div> <div>73%</div> <div>22%</div> <div>5%</div> </div> </div>
2	B	440	<div> <div>13%</div> <div> <div></div> <div>67%</div> <div>25%</div> <div>7%</div> </div> </div>
3	C	398	<div> <div>8%</div> <div> <div></div> <div>78%</div> <div>20%</div> <div>..</div> </div> </div>
4	D	418	<div> <div>10%</div> <div> <div></div> <div>64%</div> <div>26%</div> <div>9%</div> </div> </div>





















Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	403	
6	F	439	
7	G	246	
7	g	246	
8	H	234	
8	h	234	
9	I	261	
9	i	261	
10	J	248	
10	j	248	
11	K	241	
11	k	241	
12	L	263	
12	l	263	
13	M	255	
13	m	255	
14	N	239	
14	n	239	
15	O	277	
15	o	277	
16	P	205	
16	p	205	
17	Q	201	
17	q	201	
18	R	263	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
18	r	263	
19	S	241	
19	s	241	
20	T	264	
20	t	264	
21	U	953	
22	V	534	
23	W	456	
24	X	422	
25	Y	389	
26	Z	324	
27	a	376	
28	b	377	
29	c	310	
30	d	350	
31	e	70	
32	f	908	
33	v	36	
34	y	5	
34	z	5	

2 Entry composition

There are 38 unique types of molecules in this entry. The entry contains 106358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 26S proteasome regulatory subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	A	413	Total	C	N	O	S	0	0
			3229	2034	566	611	18		

- Molecule 2 is a protein called 26S proteasome regulatory subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	B	411	Total	C	N	O	S	0	0
			3207	2022	548	622	15		

- Molecule 3 is a protein called 26S proteasome regulatory subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	C	396	Total	C	N	O	S	0	0
			3105	1954	558	576	17		

- Molecule 4 is a protein called 26S proteasome regulatory subunit 6B.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	D	380	Total	C	N	O	S	0	0
			3040	1923	524	580	13		

- Molecule 5 is a protein called Proteasome 26S subunit, ATPase 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	389	Total	C	N	O	S	0	0
			3097	1947	552	581	17		

- Molecule 6 is a protein called 26S proteasome regulatory subunit 6A.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	395	Total	C	N	O	S	0	0
			3098	1951	533	596	18		

- Molecule 7 is a protein called Proteasome subunit alpha type-6.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	240	Total	C	N	O	S	0	0
			1867	1187	312	355	13		
7	g	244	Total	C	N	O	S	0	0
			1879	1193	318	355	13		

- Molecule 8 is a protein called Proteasome subunit alpha type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	H	232	Total	C	N	O	S	0	0
			1801	1149	304	342	6		
8	h	232	Total	C	N	O	S	0	0
			1805	1154	307	338	6		

- Molecule 9 is a protein called Proteasome subunit alpha type-4.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	I	248	Total	C	N	O	S	0	0
			1933	1222	330	371	10		
9	i	250	Total	C	N	O	S	0	0
			1955	1234	336	375	10		

- Molecule 10 is a protein called Proteasome subunit alpha type-7.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	J	239	Total	C	N	O	S	0	0
			1861	1166	327	363	5		
10	j	239	Total	C	N	O	S	0	0
			1861	1168	332	356	5		

- Molecule 11 is a protein called Proteasome subunit alpha type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	K	238	Total	C	N	O	S	0	0
			1813	1139	302	361	11		
11	k	234	Total	C	N	O	S	0	0
			1782	1119	295	357	11		

- Molecule 12 is a protein called Proteasome subunit alpha type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	L	240	Total	C	N	O	S	0	0
			1876	1175	338	352	11		
12	l	238	Total	C	N	O	S	0	0
			1861	1165	335	350	11		

- Molecule 13 is a protein called Proteasome subunit alpha type-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	M	242	Total	C	N	O	S	1	0
			1893	1202	323	356	12		
13	m	240	Total	C	N	O	S	0	0
			1881	1193	321	356	11		

- Molecule 14 is a protein called Proteasome subunit beta type-6.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	N	203	Total	C	N	O	S	0	0
			1521	954	259	296	12		
14	n	202	Total	C	N	O	S	0	0
			1510	947	258	293	12		

- Molecule 15 is a protein called Proteasome subunit beta type-7.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	O	220	Total	C	N	O	S	0	0
			1645	1035	278	320	12		
15	o	220	Total	C	N	O	S	0	0
			1659	1044	283	320	12		

- Molecule 16 is a protein called Proteasome subunit beta type-3.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	P	204	Total	C	N	O	S	0	0
			1587	1010	264	294	19		
16	p	204	Total	C	N	O	S	0	0
			1591	1013	265	294	19		

- Molecule 17 is a protein called Proteasome subunit beta type-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	Q	199	Total	C	N	O	S	1	0
			1591	1019	270	292	10		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf	Trace
17	q	199	Total	C	N	O	S	0	0
			1578	1012	267	290	9		

- Molecule 18 is a protein called Proteasome subunit beta type-5.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	R	201	Total	C	N	O	S	0	0
			1559	982	274	294	9		
18	r	201	Total	C	N	O	S	0	0
			1549	977	270	293	9		

- Molecule 19 is a protein called Proteasome subunit beta type-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	S	213	Total	C	N	O	S	0	0
			1641	1041	281	309	10		
19	s	213	Total	C	N	O	S	0	0
			1650	1044	283	313	10		

- Molecule 20 is a protein called Proteasome subunit beta type-4.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	T	216	Total	C	N	O	S	0	0
			1683	1062	291	318	12		
20	t	216	Total	C	N	O	S	0	0
			1687	1064	291	320	12		

- Molecule 21 is a protein called 26S proteasome non-ATPase regulatory subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	U	878	Total	C	N	O	S	0	0
			6867	4352	1163	1306	46		

- Molecule 22 is a protein called 26S proteasome non-ATPase regulatory subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	V	444	Total	C	N	O	S	0	0
			3612	2301	645	653	13		

- Molecule 23 is a protein called 26S proteasome non-ATPase regulatory subunit 12.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	W	441	Total	C	N	O	S	0	0
			3596	2277	613	682	24		

- Molecule 24 is a protein called 26S proteasome non-ATPase regulatory subunit 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	X	422	Total	C	N	O	S	0	0
			3335	2116	567	639	13		

- Molecule 25 is a protein called 26S proteasome non-ATPase regulatory subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	Y	389	Total	C	N	O	S	0	0
			3202	2041	545	598	18		

- Molecule 26 is a protein called 26S proteasome non-ATPase regulatory subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	Z	286	Total	C	N	O	S	0	0
			2281	1457	392	427	5		

- Molecule 27 is a protein called 26S proteasome non-ATPase regulatory subunit 13.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	a	373	Total	C	N	O	S	0	0
			2995	1911	510	559	15		

- Molecule 28 is a protein called 26S proteasome non-ATPase regulatory subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	b	191	Total	C	N	O	S	0	0
			1458	910	261	279	8		

- Molecule 29 is a protein called 26S proteasome non-ATPase regulatory subunit 14.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	c	287	Total	C	N	O	S	0	0
			2260	1430	389	422	19		

- Molecule 30 is a protein called 26S proteasome non-ATPase regulatory subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	d	257	Total	C	N	O	S	0	0
			2116	1371	346	390	9		

- Molecule 31 is a protein called 26S proteasome complex subunit SEM1.

Mol	Chain	Residues	Atoms				AltConf	Trace
31	e	50	Total	C	N	O	0	0
			425	260	65	100		

- Molecule 32 is a protein called 26S proteasome non-ATPase regulatory subunit 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
32	f	844	Total	C	N	O	S	0	0
			6529	4126	1108	1250	45		


- Molecule 33 is a protein called Substrate binding RP.

Mol	Chain	Residues	Atoms				AltConf	Trace
33	v	36	Total	C	N	O	0	0
			180	108	36	36		

- Molecule 34 is a protein called Substrates binding CP.

Mol	Chain	Residues	Atoms				AltConf	Trace
34	y	5	Total	C	N	O	0	0
			25	15	5	5		
34	z	5	Total	C	N	O	0	0
			25	15	5	5		

- Molecule 35 is ADENOSINE-5'-TRIPHOSPHATE (CCD ID: ATP) (formula: C₁₀H₁₆N₅O₁₃P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
35	A	1	Total	C	N	O	P	0
			31	10	5	13	3	
35	B	1	Total	C	N	O	P	0
			31	10	5	13	3	
35	C	1	Total	C	N	O	P	0
			31	10	5	13	3	
35	F	1	Total	C	N	O	P	0
			31	10	5	13	3	

- Molecule 36 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		AltConf
36	A	1	Total	Mg	0
			1	1	
36	B	1	Total	Mg	0
			1	1	
36	C	1	Total	Mg	0
			1	1	
36	D	1	Total	Mg	0
			1	1	
36	F	1	Total	Mg	0
			1	1	

- Molecule 37 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
37	D	1	Total	C	N	O	P	0
			27	10	5	10	2	


- Molecule 38 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
38	c	1	Total Zn 1 1	0


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 26S proteasome regulatory subunit 7





- Molecule 2: 26S proteasome regulatory subunit 4






• Molecule 3: 26S proteasome regulatory subunit 8




• Molecule 4: 26S proteasome regulatory subunit 6B




• Molecule 5: Proteasome 26S subunit, ATPase 6






• Molecule 6: 26S proteasome regulatory subunit 6A





• Molecule 7: Proteasome subunit alpha type-6








- Molecule 7: Proteasome subunit alpha type-6




- Molecule 8: Proteasome subunit alpha type-2




- Molecule 8: Proteasome subunit alpha type-2




- Molecule 9: Proteasome subunit alpha type-4





- Molecule 9: Proteasome subunit alpha type-4







- Molecule 10: Proteasome subunit alpha type-7





- Molecule 10: Proteasome subunit alpha type-7




- Molecule 11: Proteasome subunit alpha type-5





- Molecule 11: Proteasome subunit alpha type-5







- Molecule 12: Proteasome subunit alpha type-1




- Molecule 12: Proteasome subunit alpha type-1




- Molecule 13: Proteasome subunit alpha type-3




- Molecule 13: Proteasome subunit alpha type-3





- Molecule 14: Proteasome subunit beta type-6

Chain N:  75% 9% 15%




- Molecule 14: Proteasome subunit beta type-6

Chain n:  79% 6% 15%




- Molecule 15: Proteasome subunit beta type-7

Chain O:  71% 8% 21%




- Molecule 15: Proteasome subunit beta type-7

Chain o:  69% 10% 21%




- Molecule 16: Proteasome subunit beta type-3


Chain P:  85% 13% 1%




- Molecule 16: Proteasome subunit beta type-3




- Molecule 17: Proteasome subunit beta type-2




- Molecule 17: Proteasome subunit beta type-2



- Molecule 18: Proteasome subunit beta type-5




- Molecule 18: Proteasome subunit beta type-5




- Molecule 19: Proteasome subunit beta type-1

Chain S: 




- Molecule 19: Proteasome subunit beta type-1

Chain s: 




- Molecule 20: Proteasome subunit beta type-4

Chain T: 




- Molecule 20: Proteasome subunit beta type-4

Chain t: 





- Molecule 21: 26S proteasome non-ATPase regulatory subunit 1

Chain U: 




- Molecule 22: 26S proteasome non-ATPase regulatory subunit 3







• Molecule 23: 26S proteasome non-ATPase regulatory subunit 12





• Molecule 24: 26S proteasome non-ATPase regulatory subunit 11







• Molecule 25: 26S proteasome non-ATPase regulatory subunit 6




• Molecule 26: 26S proteasome non-ATPase regulatory subunit 7







- Molecule 27: 26S proteasome non-ATPase regulatory subunit 13




- Molecule 28: 26S proteasome non-ATPase regulatory subunit 4




- Molecule 29: 26S proteasome non-ATPase regulatory subunit 14

Chain c: 15% 64% 28% 7%




• Molecule 30: 26S proteasome non-ATPase regulatory subunit 8

Chain d: 29% 55% 19% 27%




• Molecule 31: 26S proteasome complex subunit SEM1

Chain e: 21% 51% 16% 29%




• Molecule 32: 26S proteasome non-ATPase regulatory subunit 2


Chain f: 31% 67% 26% 7%



- Molecule 34: Substrates binding CP



- Molecule 34: Substrates binding CP



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	856683	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	44	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	3500	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.026	Depositor
Minimum map value	-0.006	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.00594	Depositor
Map size (\AA)	411.0, 411.0, 411.0	wwPDB
Map dimensions	600, 600, 600	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	0.685, 0.685, 0.685	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, ATP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/3283	0.41	0/4433
2	B	0.19	0/3254	0.48	0/4388
3	C	0.20	0/3146	0.49	1/4226 (0.0%)
4	D	0.21	0/3090	0.54	2/4168 (0.0%)
5	E	0.19	0/3145	0.46	0/4233
6	F	0.19	0/3137	0.47	0/4223
7	G	0.13	0/1901	0.36	0/2572
7	g	0.17	0/1913	0.40	0/2589
8	H	0.14	0/1840	0.39	0/2495
8	h	0.13	0/1844	0.35	0/2497
9	I	0.20	0/1963	0.43	0/2650
9	i	0.14	0/1985	0.35	0/2677
10	J	0.17	0/1887	0.38	0/2553
10	j	0.14	0/1887	0.39	0/2549
11	K	0.14	0/1841	0.38	0/2486
11	k	0.15	0/1809	0.39	0/2444
12	L	0.14	0/1911	0.34	0/2584
12	l	0.14	0/1896	0.35	0/2565
13	M	0.13	0/1931	0.37	0/2600
13	m	0.13	0/1916	0.32	0/2580
14	N	0.20	0/1548	0.37	0/2097
14	n	0.13	0/1536	0.36	0/2080
15	O	0.13	0/1672	0.38	1/2267 (0.0%)
15	o	0.12	0/1686	0.36	0/2282
16	P	0.21	0/1616	0.42	0/2180
16	p	0.16	0/1620	0.42	0/2184
17	Q	0.13	0/1627	0.38	0/2202
17	q	0.18	0/1611	0.44	0/2182
18	R	0.15	0/1590	0.39	0/2147
18	r	0.14	0/1580	0.40	0/2135
19	S	0.16	0/1671	0.43	0/2252
19	s	0.13	0/1680	0.36	0/2264

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
20	T	0.16	0/1716	0.44	0/2323
20	t	0.17	0/1720	0.42	0/2328
21	U	0.16	0/6984	0.41	0/9435
22	V	0.18	0/3681	0.42	0/4969
23	W	0.14	0/3644	0.37	0/4901
24	X	0.17	0/3381	0.37	0/4558
25	Y	0.16	0/3261	0.39	1/4393 (0.0%)
26	Z	0.22	0/2324	0.49	0/3150
27	a	0.18	0/3053	0.46	0/4133
28	b	0.14	0/1478	0.42	0/2001
29	c	0.19	0/2302	0.51	0/3110
30	d	0.16	0/2162	0.44	0/2919
31	e	0.34	0/437	0.55	0/595
32	f	0.16	0/6640	0.48	2/8988 (0.0%)
All	All	0.17	0/107799	0.42	7/145587 (0.0%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
32	f	113	MET	CA-C-N	14.78	140.94	120.39
32	f	113	MET	C-N-CA	14.78	140.94	120.39
25	Y	236	LEU	N-CA-C	-5.83	105.06	111.82
3	C	221	GLN	CB-CA-C	-5.56	110.15	116.54
4	D	384	MET	CB-CG-SD	5.37	128.82	112.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3229	0	3261	75	0
2	B	3207	0	3277	93	0
3	C	3105	0	3219	64	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	D	3040	0	3076	89	0
5	E	3097	0	3174	95	0
6	F	3098	0	3187	69	0
7	G	1867	0	1867	23	0
7	g	1879	0	1872	22	0
8	H	1801	0	1773	20	0
8	h	1805	0	1798	15	0
9	I	1933	0	1923	28	0
9	i	1955	0	1955	24	0
10	J	1861	0	1846	31	0
10	j	1861	0	1865	50	0
11	K	1813	0	1796	11	0
11	k	1782	0	1766	37	0
12	L	1876	0	1856	30	0
12	l	1861	0	1839	28	0
13	M	1893	0	1885	21	0
13	m	1881	0	1868	27	0
14	N	1521	0	1494	16	0
14	n	1510	0	1483	10	0
15	O	1645	0	1648	20	0
15	o	1659	0	1681	18	0
16	P	1587	0	1598	19	0
16	p	1591	0	1609	32	0
17	Q	1591	0	1589	18	0
17	q	1578	0	1569	18	0
18	R	1559	0	1523	23	0
18	r	1549	0	1506	14	0
19	S	1641	0	1639	27	0
19	s	1650	0	1645	22	0
20	T	1683	0	1662	28	0
20	t	1687	0	1666	23	0
21	U	6867	0	6929	187	0
22	V	3612	0	3682	92	0
23	W	3596	0	3713	84	0
24	X	3335	0	3435	81	0
25	Y	3202	0	3204	81	0
26	Z	2281	0	2312	89	0
27	a	2995	0	3012	81	0
28	b	1458	0	1505	66	0
29	c	2260	0	2276	67	0
30	d	2116	0	2146	51	0
31	e	425	0	328	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
32	f	6529	0	6541	173	0
33	v	180	0	50	1	0
34	y	25	0	7	0	0
34	z	25	0	7	1	0
35	A	31	0	12	0	0
35	B	31	0	12	1	0
35	C	31	0	12	0	0
35	F	31	0	12	1	0
36	A	1	0	0	0	0
36	B	1	0	0	0	0
36	C	1	0	0	0	0
36	D	1	0	0	0	0
36	F	1	0	0	0	0
37	D	27	0	12	3	0
38	c	1	0	0	0	0
All	All	106358	0	106622	2008	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 2008 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
20:T:108:ASN:HB3	20:T:110:MET:HE1	1.36	1.08
16:p:88:MET:HE3	16:p:130:PRO:HB2	1.44	1.00
12:l:61:LYS:HE2	12:l:61:LYS:H	1.36	0.91
19:S:147:PRO:HB2	16:p:149:MET:HE1	1.55	0.89
29:c:113:HIS:NE2	29:c:115:HIS:CE1	2.41	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	411/433 (95%)	381 (93%)	30 (7%)	0	100	100
2	B	409/440 (93%)	362 (88%)	46 (11%)	1 (0%)	43	63
3	C	394/398 (99%)	357 (91%)	35 (9%)	2 (0%)	24	43
4	D	378/418 (90%)	318 (84%)	57 (15%)	3 (1%)	16	31
5	E	387/403 (96%)	334 (86%)	53 (14%)	0	100	100
6	F	391/439 (89%)	354 (90%)	34 (9%)	3 (1%)	16	31
7	G	238/246 (97%)	224 (94%)	14 (6%)	0	100	100
7	g	242/246 (98%)	235 (97%)	7 (3%)	0	100	100
8	H	230/234 (98%)	219 (95%)	11 (5%)	0	100	100
8	h	230/234 (98%)	222 (96%)	8 (4%)	0	100	100
9	I	246/261 (94%)	235 (96%)	10 (4%)	1 (0%)	30	49
9	i	248/261 (95%)	243 (98%)	5 (2%)	0	100	100
10	J	237/248 (96%)	227 (96%)	10 (4%)	0	100	100
10	j	237/248 (96%)	226 (95%)	10 (4%)	1 (0%)	30	49
11	K	236/241 (98%)	225 (95%)	11 (5%)	0	100	100
11	k	232/241 (96%)	224 (97%)	8 (3%)	0	100	100
12	L	238/263 (90%)	230 (97%)	8 (3%)	0	100	100
12	l	236/263 (90%)	227 (96%)	9 (4%)	0	100	100
13	M	241/255 (94%)	234 (97%)	7 (3%)	0	100	100
13	m	238/255 (93%)	236 (99%)	2 (1%)	0	100	100
14	N	201/239 (84%)	195 (97%)	6 (3%)	0	100	100
14	n	200/239 (84%)	194 (97%)	6 (3%)	0	100	100
15	O	218/277 (79%)	212 (97%)	6 (3%)	0	100	100
15	o	218/277 (79%)	210 (96%)	8 (4%)	0	100	100
16	P	202/205 (98%)	195 (96%)	7 (4%)	0	100	100
16	p	202/205 (98%)	196 (97%)	6 (3%)	0	100	100
17	Q	198/201 (98%)	189 (96%)	9 (4%)	0	100	100
17	q	197/201 (98%)	191 (97%)	6 (3%)	0	100	100
18	R	199/263 (76%)	192 (96%)	7 (4%)	0	100	100
18	r	199/263 (76%)	194 (98%)	5 (2%)	0	100	100
19	S	211/241 (88%)	202 (96%)	9 (4%)	0	100	100
19	s	211/241 (88%)	204 (97%)	7 (3%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
20	T	214/264 (81%)	208 (97%)	6 (3%)	0	100	100
20	t	214/264 (81%)	206 (96%)	8 (4%)	0	100	100
21	U	874/953 (92%)	814 (93%)	60 (7%)	0	100	100
22	V	442/534 (83%)	420 (95%)	22 (5%)	0	100	100
23	W	439/456 (96%)	432 (98%)	7 (2%)	0	100	100
24	X	420/422 (100%)	405 (96%)	15 (4%)	0	100	100
25	Y	387/389 (100%)	363 (94%)	24 (6%)	0	100	100
26	Z	284/324 (88%)	257 (90%)	27 (10%)	0	100	100
27	a	371/376 (99%)	344 (93%)	27 (7%)	0	100	100
28	b	189/377 (50%)	171 (90%)	18 (10%)	0	100	100
29	c	285/310 (92%)	243 (85%)	42 (15%)	0	100	100
30	d	255/350 (73%)	219 (86%)	36 (14%)	0	100	100
31	e	48/70 (69%)	39 (81%)	9 (19%)	0	100	100
32	f	840/908 (92%)	808 (96%)	32 (4%)	0	100	100
All	All	13417/14876 (90%)	12616 (94%)	790 (6%)	11 (0%)	49	68

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	D	157	ASP
6	F	73	ILE
6	F	86	LEU
9	I	53	HIS
2	B	319	PHE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/372 (94%)	340 (98%)	8 (2%)	44	72
2	B	357/385 (93%)	348 (98%)	9 (2%)	42	69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	340/346 (98%)	334 (98%)	6 (2%)	51	77
4	D	333/366 (91%)	329 (99%)	4 (1%)	63	83
5	E	341/353 (97%)	337 (99%)	4 (1%)	63	83
6	F	340/379 (90%)	334 (98%)	6 (2%)	51	77
7	G	202/210 (96%)	201 (100%)	1 (0%)	81	92
7	g	201/210 (96%)	196 (98%)	5 (2%)	42	69
8	H	187/191 (98%)	185 (99%)	2 (1%)	65	84
8	h	188/191 (98%)	185 (98%)	3 (2%)	55	79
9	I	202/221 (91%)	199 (98%)	3 (2%)	57	80
9	i	206/221 (93%)	205 (100%)	1 (0%)	81	92
10	J	197/211 (93%)	192 (98%)	5 (2%)	42	69
10	j	196/211 (93%)	195 (100%)	1 (0%)	81	92
11	K	197/203 (97%)	196 (100%)	1 (0%)	81	92
11	k	195/203 (96%)	193 (99%)	2 (1%)	68	86
12	L	202/224 (90%)	200 (99%)	2 (1%)	68	86
12	l	201/224 (90%)	199 (99%)	2 (1%)	68	86
13	M	199/212 (94%)	197 (99%)	2 (1%)	68	86
13	m	198/212 (93%)	197 (100%)	1 (0%)	81	92
14	N	158/181 (87%)	156 (99%)	2 (1%)	61	82
14	n	156/181 (86%)	156 (100%)	0	100	100
15	O	178/228 (78%)	177 (99%)	1 (1%)	78	91
15	o	181/228 (79%)	181 (100%)	0	100	100
16	P	172/174 (99%)	169 (98%)	3 (2%)	53	78
16	p	173/174 (99%)	169 (98%)	4 (2%)	44	72
17	Q	169/171 (99%)	169 (100%)	0	100	100
17	q	166/171 (97%)	164 (99%)	2 (1%)	63	83
18	R	156/202 (77%)	155 (99%)	1 (1%)	78	91
18	r	154/202 (76%)	152 (99%)	2 (1%)	61	82
19	S	175/199 (88%)	170 (97%)	5 (3%)	37	65
19	s	177/199 (89%)	177 (100%)	0	100	100
20	T	178/215 (83%)	177 (99%)	1 (1%)	78	91

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
20	t	179/215 (83%)	177 (99%)	2 (1%)	65	84
21	U	752/816 (92%)	739 (98%)	13 (2%)	53	78
22	V	390/460 (85%)	382 (98%)	8 (2%)	47	74
23	W	406/416 (98%)	405 (100%)	1 (0%)	87	95
24	X	362/362 (100%)	357 (99%)	5 (1%)	59	81
25	Y	344/344 (100%)	339 (98%)	5 (2%)	57	80
26	Z	257/295 (87%)	247 (96%)	10 (4%)	28	55
27	a	333/336 (99%)	325 (98%)	8 (2%)	43	70
28	b	167/312 (54%)	165 (99%)	2 (1%)	63	83
29	c	252/268 (94%)	249 (99%)	3 (1%)	63	83
30	d	231/294 (79%)	227 (98%)	4 (2%)	53	78
31	e	44/63 (70%)	40 (91%)	4 (9%)	9	19
32	f	711/763 (93%)	701 (99%)	10 (1%)	59	81
All	All	11451/12614 (91%)	11287 (99%)	164 (1%)	57	81

5 of 164 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
27	a	365	MET
7	g	66	VAL
29	c	269	GLN
32	f	340	MET
11	k	206	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 92 such sidechains are listed below:

Mol	Chain	Res	Type
29	c	274	ASN
9	i	235	GLN
30	d	46	GLN
32	f	737	ASN
12	l	59	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
35	ATP	B	501	36	29,33,33	0.29	0	44,52,52	0.52	1 (2%)
37	ADP	D	502	36	27,29,29	1.37	4 (14%)	42,45,45	2.01	9 (21%)
35	ATP	A	501	36	29,33,33	0.29	0	44,52,52	0.51	1 (2%)
35	ATP	C	501	36	29,33,33	0.31	0	44,52,52	0.54	1 (2%)
35	ATP	F	501	36	29,33,33	0.29	0	44,52,52	0.50	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
35	ATP	B	501	36	-	2/22/38/38	0/3/3/3
37	ADP	D	502	36	-	3/16/32/32	0/3/3/3
35	ATP	A	501	36	-	3/22/38/38	0/3/3/3
35	ATP	C	501	36	-	5/22/38/38	0/3/3/3
35	ATP	F	501	36	-	6/22/38/38	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
37	D	502	ADP	C5-C4	4.66	1.47	1.39
37	D	502	ADP	C5-C6	2.67	1.48	1.41
37	D	502	ADP	C8-N7	2.34	1.36	1.31
37	D	502	ADP	C5-N7	-2.28	1.34	1.39

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
37	D	502	ADP	C5-C4-N3	-6.59	118.15	126.75
37	D	502	ADP	N3-C4-N9	5.22	135.68	127.08
37	D	502	ADP	C2-N3-C4	3.97	121.12	111.75
37	D	502	ADP	PA-O3A-PB	-3.60	120.47	132.83
37	D	502	ADP	C4-C5-N7	-3.13	106.80	110.62

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
35	C	501	ATP	C5'-O5'-PA-O3A
35	C	501	ATP	C3'-C4'-C5'-O5'
35	F	501	ATP	C5'-O5'-PA-O1A
35	F	501	ATP	C5'-O5'-PA-O2A
35	C	501	ATP	O4'-C4'-C5'-O5'


There are no ring outliers.


3 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
35	B	501	ATP	1	0
37	D	502	ADP	3	0
35	F	501	ATP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient

equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-62079. These allow visual inspection of the internal detail of the map and identification of artifacts.

Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 300




Y Index: 300




Z Index: 300


6.2.2 Raw map



X Index: 300



Y Index: 300




Z Index: 300


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 310




Y Index: 280




Z Index: 275


6.3.2 Raw map



X Index: 275



Y Index: 280




Z Index: 381


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.00594. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 700 nm³; this corresponds to an approximate mass of 633 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.400 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.400 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.50	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	3.73	4.22	3.79

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.73 differs from the reported value 2.5 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-62079 and PDB model 9K53. Per-residue inclusion information can be found in section [3](#) on page [13](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.00594 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.00594).







































































9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 69% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ































The table lists the average atom inclusion at the recommended contour level (0.00594) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6880	 0.3910
A	 0.6520	 0.3820
B	 0.6450	 0.3880
C	 0.6960	 0.4050
D	 0.6680	 0.3760
E	 0.4010	 0.2570
F	 0.5800	 0.3500
G	 0.8170	 0.4710
H	 0.8410	 0.4730
I	 0.8010	 0.4670
J	 0.7830	 0.4450
K	 0.7840	 0.4560
L	 0.8340	 0.4700
M	 0.8120	 0.4690
N	 0.8550	 0.4870
O	 0.8570	 0.4880
P	 0.8680	 0.4780
Q	 0.8590	 0.4790
R	 0.8690	 0.4660
S	 0.8380	 0.4730
T	 0.8570	 0.4860
U	 0.5850	 0.3100
V	 0.5890	 0.3260
W	 0.5100	 0.3120
X	 0.5400	 0.3260
Y	 0.7380	 0.3670
Z	 0.6320	 0.3530
a	 0.5750	 0.2940
b	 0.5240	 0.3030
c	 0.6180	 0.3510
d	 0.4630	 0.2650
e	 0.5300	 0.3460
f	 0.5020	 0.2830
g	 0.7910	 0.4560
h	 0.7850	 0.4530



Continued on next page...

Continued from previous page...

Chain	Atom inclusion	Q-score
i	 0.7440	 0.4260
j	 0.7030	 0.3970
k	 0.7380	 0.4300
l	 0.8130	 0.4590
m	 0.8050	 0.4600
n	 0.8410	 0.4790
o	 0.8360	 0.4760
p	 0.8410	 0.4660
q	 0.8460	 0.4720
r	 0.8790	 0.4870
s	 0.8350	 0.4790
t	 0.8530	 0.4710
v	 0.2610	 0.3670
y	 0.5600	 0.3570
z	 0.5600	 0.3820