



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 22, 2024 – 04:08 PM JST

PDB ID : 8K62  
Title : Crystal structure of ALKBH1 and 13h complex.  
Authors : Liang, X.; Yinping, G.; Feng, L.; Jiang, Z.; Ke, X.; Shengyong, Y.  
Deposited on : 2023-07-24  
Resolution : 1.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.37.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 5242 atoms, of which 2560 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Nucleic acid dioxygenase ALKBH1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	326	5162	1675	2550	452	470	15	0	1	0

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	A	1	1	1	0	0

- Molecule 3 is 1-[5-[[3-(trifluoromethoxy)phenyl]methoxy]pyrimidin-2-yl]pyrazole-4-carboxylic acid (three-letter code: IAU) (formula: C<sub>16</sub>H<sub>11</sub>F<sub>3</sub>N<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	F	H	N	O		
3	A	1	37	16	3	10	4	4	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	42	Total	O	0	0
			42	42		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Nucleic acid dioxygenase ALKBH1



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	56.50Å 56.50Å 192.13Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	34.28 – 1.99 34.28 – 1.99	Depositor EDS
% Data completeness (in resolution range)	85.9 (34.28-1.99) 85.9 (34.28-1.99)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.28 (at 2.00Å)	Xtrriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, $R_{free}$	0.204 , 0.278 0.204 , 0.278	Depositor DCC
$R_{free}$ test set	1996 reflections (9.81%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.2	Xtrriage
Anisotropy	0.526	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.40 , 48.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.097 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5242	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	60.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.17% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IAU, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.68	0/2690	0.77	3/3640 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	195	ASP	CB-CG-OD1	5.17	122.95	118.30
1	A	233	ASP	CB-CG-OD1	-5.13	113.68	118.30
1	A	165	LEU	CA-CB-CG	5.07	126.96	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	352	ASN	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2612	2550	2549	48	1
2	A	1	0	0	0	0
3	A	27	10	0	0	0
4	A	42	0	0	0	0
All	All	2682	2560	2549	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:VAL:HB	1:A:85:VAL:HG21	1.76	0.67
1:A:42:GLY:O	1:A:55:LYS:HE3	1.97	0.64
1:A:133:LYS:HG3	1:A:134:HIS:H	1.69	0.57
1:A:110:TYR:HE2	1:A:319:VAL:HG11	1.69	0.56
1:A:42:GLY:HA2	1:A:55:LYS:HG3	1.88	0.56
1:A:237:LEU:O	1:A:238:ASP:HB2	2.05	0.56
1:A:42:GLY:C	1:A:55:LYS:HE3	2.26	0.55
1:A:110:TYR:CE2	1:A:319:VAL:HG11	2.42	0.55
1:A:121:LEU:O	1:A:124:GLN:HB2	2.06	0.54
1:A:44:ILE:HG13	1:A:55:LYS:HE2	1.89	0.53
1:A:55:LYS:HZ2	1:A:55:LYS:HB2	1.74	0.53
1:A:19:GLY:HA3	1:A:24:ARG:HB3	1.92	0.52
1:A:236:GLU:HG3	1:A:237:LEU:O	2.10	0.52
1:A:78:TYR:CE2	1:A:314:PRO:HG2	2.46	0.50
1:A:213:ARG:HG2	1:A:213:ARG:NH1	2.26	0.50
1:A:291:ARG:HG3	1:A:293:LEU:HD13	1.94	0.49
1:A:113:HIS:CG	1:A:319:VAL:HG23	2.48	0.49
1:A:166:GLU:HG2	1:A:223:ARG:NH2	2.28	0.48
1:A:55:LYS:HB2	1:A:55:LYS:NZ	2.28	0.48
1:A:44:ILE:CG1	1:A:55:LYS:HE2	2.45	0.47
1:A:133:LYS:HG3	1:A:134:HIS:N	2.29	0.46
1:A:170:TRP:HA	1:A:219:LEU:O	2.14	0.46
1:A:117:GLN:HB3	1:A:122:TYR:CZ	2.51	0.46
1:A:55:LYS:HD2	1:A:59:ALA:H	1.81	0.46
1:A:165:LEU:HD22	1:A:165:LEU:O	2.16	0.45
1:A:163:SER:HB3	1:A:166:GLU:HG3	1.98	0.45
1:A:113:HIS:ND1	1:A:319:VAL:HG23	2.32	0.45
1:A:133:LYS:CG	1:A:134:HIS:H	2.29	0.45

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:GLN:OE1	1:A:272:SER:OG	2.30	0.44
1:A:119:LEU:HD22	1:A:165:LEU:HD21	2.00	0.44
1:A:165:LEU:HD11	1:A:333:TYR:CD1	2.53	0.44
1:A:258:GLY:HA2	1:A:285:LEU:HD22	2.00	0.43
1:A:33:SER:OG	1:A:39:ASP:HB3	2.18	0.43
1:A:253:ILE:O	1:A:290:PRO:HD2	2.17	0.43
1:A:143:LEU:CB	1:A:164:LEU:HD22	2.47	0.43
1:A:317:SER:OG	1:A:319:VAL:HG12	2.19	0.43
1:A:42:GLY:HA2	1:A:55:LYS:CG	2.49	0.43
1:A:237:LEU:HD21	1:A:284:LEU:HD11	2.00	0.42
1:A:143:LEU:HD12	1:A:164:LEU:CD2	2.50	0.42
1:A:220:ASN:HB2	1:A:340:ASN:HB3	2.02	0.41
1:A:179:TRP:O	1:A:182:LYS:HE2	2.21	0.41
1:A:182:LYS:O	1:A:183:LYS:HG3	2.21	0.41
1:A:125:LYS:N	1:A:126:PRO:CD	2.84	0.41
1:A:78:TYR:CE2	1:A:314:PRO:CG	3.04	0.40
1:A:68:ASN:HB3	1:A:71:SER:OG	2.22	0.40
1:A:176:HIS:O	1:A:185:SER:N	2.42	0.40
1:A:143:LEU:HD12	1:A:164:LEU:HD23	2.04	0.40
1:A:237:LEU:CD2	1:A:284:LEU:HD11	2.52	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:GLU:OE2	1:A:153:TYR:OH[5_665]	1.92	0.28

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	321/389 (82%)	298 (93%)	22 (7%)	1 (0%)	41 37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	353	PHE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	280/331 (85%)	265 (95%)	15 (5%)	22 18

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	23	PHE
1	A	50	HIS
1	A	53	ARG
1	A	60	GLN
1	A	124	GLN
1	A	132	ASP
1	A	148	LYS
1	A	154	LYS
1	A	165	LEU
1	A	293	LEU
1	A	309	LEU
1	A	315	ARG
1	A	323	SER
1	A	325	GLU
1	A	347	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	IAU	A	402	2	27,29,29	1.08	1 (3%)	36,41,41	2.60	9 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	IAU	A	402	2	-	7/16/18/18	0/3/3/3

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	IAU	C13-N14	-2.03	1.30	1.34

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	IAU	C08-N07-N06	9.96	110.95	103.70
3	A	402	IAU	C11-C12-C13	5.48	120.18	116.56
3	A	402	IAU	C12-C11-N10	-5.08	116.74	122.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	IAU	N10-C09-N06	5.00	120.53	116.00
3	A	402	IAU	C11-N10-C09	3.83	119.06	115.61
3	A	402	IAU	C16-C17-C22	-2.22	114.99	120.29
3	A	402	IAU	C16-C17-C18	2.19	125.78	120.66
3	A	402	IAU	N14-C09-N06	-2.13	114.06	116.00
3	A	402	IAU	C16-O15-C12	-2.07	112.54	117.65

There are no chirality outliers.


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	402	IAU	F25-C24-O23-C21
3	A	402	IAU	F26-C24-O23-C21
3	A	402	IAU	F27-C24-O23-C21
3	A	402	IAU	C13-C12-O15-C16
3	A	402	IAU	C11-C12-O15-C16
3	A	402	IAU	C20-C21-O23-C24
3	A	402	IAU	C22-C21-O23-C24

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	326/389 (83%)	1.01	57 (17%) <b>1</b> <b>1</b>	31, 47, 104, 134	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	52	ALA	17.6
1	A	54	GLY	7.7
1	A	53	ARG	6.9
1	A	57	PRO	6.5
1	A	51	ALA	6.4
1	A	186	ALA	6.2
1	A	133	LYS	6.1
1	A	138	GLU	6.1
1	A	187	ASP	5.9
1	A	179	TRP	5.8
1	A	135	MET	5.7
1	A	180	ASP	5.4
1	A	154	LYS	5.3
1	A	134	HIS	5.1
1	A	153	TYR	4.7
1	A	136	SER	4.6
1	A	56	GLY	4.6
1	A	178	ASN	4.5
1	A	49	ALA	4.4
1	A	188	HIS	4.2
1	A	315	ARG	3.9
1	A	181	SER	3.9
1	A	60	GLN	3.8
1	A	137	LYS	3.7
1	A	55	LYS	3.7
1	A	139	GLU	3.4
1	A	140	THR	3.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	50	HIS	3.3
1	A	142	ASP	3.3
1	A	19	GLY	3.2
1	A	164	LEU	3.2
1	A	125	LYS	3.1
1	A	58	GLY	3.0
1	A	149	GLU	2.9
1	A	20	GLU	2.8
1	A	151	LEU	2.7
1	A	78	TYR	2.7
1	A	183	LYS	2.6
1	A	352	ASN	2.6
1	A	132	ASP	2.6
1	A	152	ARG	2.5
1	A	246	PHE	2.5
1	A	343	VAL	2.4
1	A	190	THR	2.4
1	A	311	ALA	2.3
1	A	161	PRO	2.3
1	A	162	ARG	2.3
1	A	247	SER	2.3
1	A	141	GLN	2.3
1	A	106	PHE	2.2
1	A	185	SER	2.2
1	A	339	VAL	2.2
1	A	182	LYS	2.2
1	A	189	TYR	2.1
1	A	275	ILE	2.1
1	A	347	LEU	2.0
1	A	248	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IAU	A	402	27/27	0.88	0.15	33,56,80,83	0
2	MN	A	401	1/1	1.00	0.13	34,34,34,34	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.