



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 30, 2024 – 07:54 PM EST

PDB ID : 1KAQ
Title : Structure of Bacillus subtilis Nicotinic Acid Mononucleotide Adenylyl Transferase
Authors : Olland, A.M.; Underwood, K.W.; Czerwinski, R.M.; Lo, M.C.; Aulabaugh, A.; Bard, J.; Stahl, M.L.; Somers, W.S.; Sullivan, F.X.; Chopra, R.
Deposited on : 2001-11-02
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	194	38% 45% 10% . .
1	B	194	31% 46% 14% . .
1	C	194	39% 44% 11% . .
1	D	194	36% 48% 8% . .
1	E	194	30% 50% 10% . 9%
1	F	194	24% 44% 12% 7% 14%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DND	A	601	X	-	-	-
2	DND	B	501	X	-	-	-
2	DND	C	401	X	-	-	-
2	DND	D	301	X	-	-	-
2	DND	E	201	X	-	-	-
2	DND	F	190	X	-	-	-

2 Entry composition i

There are 2 unique types of molecules in this entry. The entry contains 9268 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	186	1542	1010	246	280	6	0	0	0
1	B	186	1542	1010	246	280	6	0	0	0
1	C	186	1542	1010	246	280	6	0	0	0
1	D	186	1542	1010	246	280	6	0	0	0
1	E	176	1456	957	231	262	6	0	0	0
1	F	167	1380	906	216	252	6	0	0	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	HIS	-	expression tag	UNP P54455
A	-3	MET	-	expression tag	UNP P54455
A	-2	PRO	-	expression tag	UNP P54455
A	-1	GLY	-	expression tag	UNP P54455
A	0	GLY	-	expression tag	UNP P54455
A	1	SER	MET	cloning artifact	UNP P54455
B	-4	HIS	-	expression tag	UNP P54455
B	-3	MET	-	expression tag	UNP P54455
B	-2	PRO	-	expression tag	UNP P54455
B	-1	GLY	-	expression tag	UNP P54455
B	0	GLY	-	expression tag	UNP P54455
B	1	SER	MET	cloning artifact	UNP P54455
C	-4	HIS	-	expression tag	UNP P54455
C	-3	MET	-	expression tag	UNP P54455
C	-2	PRO	-	expression tag	UNP P54455
C	-1	GLY	-	expression tag	UNP P54455
C	0	GLY	-	expression tag	UNP P54455

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	1	SER	MET	cloning artifact	UNP P54455
D	-4	HIS	-	expression tag	UNP P54455
D	-3	MET	-	expression tag	UNP P54455
D	-2	PRO	-	expression tag	UNP P54455
D	-1	GLY	-	expression tag	UNP P54455
D	0	GLY	-	expression tag	UNP P54455
D	1	SER	MET	cloning artifact	UNP P54455
E	-4	HIS	-	expression tag	UNP P54455
E	-3	MET	-	expression tag	UNP P54455
E	-2	PRO	-	expression tag	UNP P54455
E	-1	GLY	-	expression tag	UNP P54455
E	0	GLY	-	expression tag	UNP P54455
E	1	SER	MET	cloning artifact	UNP P54455
F	-4	HIS	-	expression tag	UNP P54455
F	-3	MET	-	expression tag	UNP P54455
F	-2	PRO	-	expression tag	UNP P54455
F	-1	GLY	-	expression tag	UNP P54455
F	0	GLY	-	expression tag	UNP P54455
F	1	SER	MET	cloning artifact	UNP P54455

- Molecule 2 is NICOTINIC ACID ADENINE DINUCLEOTIDE (three-letter code: DND) (formula: C₂₁H₂₇N₆O₁₅P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
2	A	1	44	21	6	15	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	B	1	Total 44	C 21	N 6	O 15	P 2	0	0
2	C	1	Total 44	C 21	N 6	O 15	P 2	0	0
2	D	1	Total 44	C 21	N 6	O 15	P 2	0	0
2	E	1	Total 44	C 21	N 6	O 15	P 2	0	0
2	F	1	Total 44	C 21	N 6	O 15	P 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE

Chain A: 




- Molecule 1: NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE


Chain B: 



- Molecule 1: NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE

Chain C: 








- Molecule 1: NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE

Chain D: 36% 48% 8%






- Molecule 1: NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE

Chain E: 30% 50% 10% 9%



- Molecule 1: NICOTINATE-NUCLEOTIDE ADENYLYLTRANSFERASE

Chain F: 24% 44% 12% 7% 14%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	78.31Å 108.78Å 177.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 3.20	Depositor
% Data completeness (in resolution range)	(Not available) (15.00-3.20)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.279 , 0.280	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	9268	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DND

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/1588	0.83	3/2151 (0.1%)
1	B	0.68	4/1588 (0.3%)	0.99	12/2151 (0.6%)
1	C	0.52	0/1588	0.80	1/2151 (0.0%)
1	D	0.57	0/1588	0.81	2/2151 (0.1%)
1	E	0.38	0/1497	0.69	2/2024 (0.1%)
1	F	0.45	0/1416	0.98	8/1914 (0.4%)
All	All	0.53	4/9265 (0.0%)	0.86	28/12542 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	46	GLN	CB-CG	-7.89	1.31	1.52
1	B	45	LYS	C-O	7.04	1.36	1.23
1	B	46	GLN	C-O	5.13	1.33	1.23
1	B	45	LYS	CA-CB	-5.04	1.42	1.53

The worst 5 of 28 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	149	VAL	C-N-CD	-17.77	81.50	120.60
1	F	113	LEU	C-N-CD	-11.90	94.42	120.60
1	B	49	ASP	CB-CG-OD2	-11.83	107.65	118.30
1	F	81	GLY	N-CA-C	-9.92	88.31	113.10
1	B	55	HIS	N-CA-C	-8.17	88.94	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1542	0	1523	135	0
1	B	1542	0	1523	155	0
1	C	1542	0	1523	123	0
1	D	1542	0	1523	120	0
1	E	1456	0	1448	137	0
1	F	1380	0	1372	158	0
2	A	44	0	25	8	0
2	B	44	0	25	4	0
2	C	44	0	25	4	0
2	D	44	0	25	8	0
2	E	44	0	25	9	0
2	F	44	0	25	5	0
All	All	9268	0	9062	802	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 44.

The worst 5 of 802 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:149:VAL:CG1	1:F:150:PRO:HD3	1.68	1.23
1:B:10:THR:HG22	1:B:39:ASN:HD22	0.99	1.16
1:B:10:THR:HG22	1:B:39:ASN:ND2	1.68	1.08
1:F:113:LEU:HB3	1:F:114:PRO:HD2	1.32	1.08
1:F:149:VAL:HG12	1:F:150:PRO:CD	1.82	1.08

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	184/194 (95%)	139 (76%)	31 (17%)	14 (8%)	1	7
1	B	184/194 (95%)	145 (79%)	18 (10%)	21 (11%)	0	2
1	C	184/194 (95%)	151 (82%)	19 (10%)	14 (8%)	1	7
1	D	184/194 (95%)	144 (78%)	27 (15%)	13 (7%)	1	8
1	E	172/194 (89%)	142 (83%)	21 (12%)	9 (5%)	2	15
1	F	161/194 (83%)	111 (69%)	24 (15%)	26 (16%)	0	0
All	All	1069/1164 (92%)	832 (78%)	140 (13%)	97 (9%)	1	3

5 of 97 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	47	ASN
1	A	51	THR
1	A	52	ASP
1	A	79	ARG
1	A	80	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	171/177 (97%)	149 (87%)	22 (13%)	4	19
1	B	171/177 (97%)	149 (87%)	22 (13%)	4	19
1	C	171/177 (97%)	148 (86%)	23 (14%)	4	18
1	D	171/177 (97%)	147 (86%)	24 (14%)	3	16
1	E	161/177 (91%)	143 (89%)	18 (11%)	6	25
1	F	154/177 (87%)	127 (82%)	27 (18%)	2	9
All	All	999/1062 (94%)	863 (86%)	136 (14%)	3	17

5 of 136 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	60	LEU
1	F	77	MET
1	F	162	ARG
1	C	46	GLN
1	C	44	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	67	ASN
1	E	184	ASN
1	F	55	HIS
1	B	184	ASN
1	B	94	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DND	B	501	-	42,48,48	1.41	5 (11%)	50,73,73	1.26	5 (10%)
2	DND	E	201	-	42,48,48	1.41	5 (11%)	50,73,73	1.39	8 (16%)
2	DND	C	401	-	42,48,48	1.43	6 (14%)	50,73,73	1.38	7 (14%)
2	DND	A	601	-	42,48,48	1.39	5 (11%)	50,73,73	1.28	7 (14%)
2	DND	F	190	-	42,48,48	1.37	3 (7%)	50,73,73	1.43	7 (14%)
2	DND	D	301	-	42,48,48	1.38	6 (14%)	50,73,73	1.37	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DND	B	501	-	3/3/11/11	9/26/62/62	0/5/5/5
2	DND	E	201	-	3/3/11/11	10/26/62/62	0/5/5/5
2	DND	C	401	-	3/3/11/11	8/26/62/62	0/5/5/5
2	DND	A	601	-	3/3/11/11	10/26/62/62	0/5/5/5
2	DND	F	190	-	3/3/11/11	12/26/62/62	0/5/5/5
2	DND	D	301	-	3/3/11/11	9/26/62/62	0/5/5/5

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	201	DND	O7N-C7N	4.74	1.36	1.22
2	B	501	DND	O7N-C7N	4.64	1.36	1.22
2	A	601	DND	O7N-C7N	4.51	1.36	1.22
2	D	301	DND	O7N-C7N	4.50	1.36	1.22
2	C	401	DND	O7N-C7N	4.47	1.35	1.22

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	401	DND	C6N-N1N-C2N	-5.57	116.89	121.97
2	E	201	DND	C6N-N1N-C2N	-5.46	117.00	121.97
2	D	301	DND	C6N-N1N-C2N	-5.08	117.34	121.97
2	F	190	DND	C6N-N1N-C2N	-5.03	117.39	121.97
2	A	601	DND	C6N-N1N-C2N	-4.91	117.50	121.97

5 of 18 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	601	DND	C3B
2	A	601	DND	C2B
2	A	601	DND	C1B
2	B	501	DND	C3B
2	B	501	DND	C2B

5 of 58 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	601	DND	C5D-O5D-PN-O12
2	A	601	DND	C5D-O5D-PN-O3P
2	A	601	DND	C5B-O5B-PA-O3P
2	A	601	DND	C5B-O5B-PA-O13
2	A	601	DND	C5B-O5B-PA-O14


There are no ring outliers.


6 monomers are involved in 38 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	501	DND	4	0
2	E	201	DND	9	0
2	C	401	DND	4	0
2	A	601	DND	8	0
2	F	190	DND	5	0
2	D	301	DND	8	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.