



# Full wwPDB X-ray Structure Validation Report ⓘ

May 13, 2020 – 08:47 am BST

PDB ID : 1KCE  
Title : E. COLI THYMIDYLATE SYNTHASE MUTANT E58Q IN COMPLEX WITH CB3717 AND 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (DUMP)  
Authors : Sage, C.R.; Rutenber, E.E.; Stout, T.J.; Stroud, R.M.  
Deposited on : 1996-10-22  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	264	
1	B	264	

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 4657 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called THYMIDYLATE SYNTHASE.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	264	Total	C	N	O	S	0	0	0
			2153	1375	371	395	12			
1	B	264	Total	C	N	O	S	0	0	0
			2153	1375	371	395	12			

- Molecule 2 is 2'-DEOXYURIDINE 5'-MONOPHOSPHATE (three-letter code: UMP) (formula: C<sub>9</sub>H<sub>13</sub>N<sub>2</sub>O<sub>8</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total	C	N	O	P	0	0
			20	9	2	8	1		
2	B	1	Total	C	N	O	P	0	0
			20	9	2	8	1		

- Molecule 3 is 10-PROPARGYL-5,8-DIDEAZAFOLIC ACID (three-letter code: CB3) (formula: C<sub>24</sub>H<sub>23</sub>N<sub>5</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
3	A	1	Total	C	N	O	0	0
			35	24	5	6		
3	B	1	Total	C	N	O	0	0
			35	24	5	6		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	143	Total	O	0	0
			143	143		
4	B	98	Total	O	0	0
			98	98		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: THYMIDYLATE SYNTHASE

Chain A:  91% 9%



- Molecule 1: THYMIDYLATE SYNTHASE

Chain B:  85% 13%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	126.80Å 126.80Å 67.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	7.00 – 2.00	Depositor
% Data completeness (in resolution range)	80.8 (7.00-2.00)	Depositor
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	X-PLOR 3.1	Depositor
R, $R_{free}$	0.172 , 0.222	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4657	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CB3, UMP, CXM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.40	0/2202	0.67	1/2990 (0.0%)
1	B	0.40	0/2202	0.68	1/2990 (0.0%)
All	All	0.40	0/4404	0.67	2/5980 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	A	27	LEU	N-CA-C	-5.52	96.08	111.00
1	B	27	LEU	N-CA-C	-5.01	97.46	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	243	ARG	Sidechain

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2153	0	2079	27	0
1	B	2153	0	2078	19	0
2	A	20	0	10	0	0
2	B	20	0	10	0	0
3	A	35	0	21	0	0
3	B	35	0	21	1	0
4	A	143	0	0	3	1
4	B	98	0	0	1	1
All	All	4657	0	4219	47	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:HIS:NE2	1:A:53:ARG:CB	1.76	1.43
1:A:51:HIS:NE2	1:A:53:ARG:HB3	1.07	1.37
1:A:51:HIS:HE1	1:A:53:ARG:NH1	1.51	1.08
1:A:51:HIS:NE2	1:A:53:ARG:HB2	1.80	0.97
1:A:51:HIS:CE1	1:A:53:ARG:NH1	2.33	0.95
1:A:51:HIS:CD2	1:A:53:ARG:CB	2.51	0.94
1:A:51:HIS:HD2	1:A:53:ARG:H	1.20	0.86
1:A:51:HIS:CE1	1:A:53:ARG:HB3	2.08	0.85
1:A:51:HIS:CD2	1:A:53:ARG:H	2.01	0.78
1:A:51:HIS:CD2	1:A:53:ARG:HB2	2.17	0.78
1:A:51:HIS:CD2	1:A:53:ARG:N	2.54	0.76
1:A:51:HIS:HD2	1:A:53:ARG:N	1.86	0.73
1:A:51:HIS:CE1	1:A:53:ARG:HH11	2.13	0.66
1:A:229:LYS:NZ	4:A:904:HOH:O	2.32	0.63
1:A:51:HIS:CD2	1:A:51:HIS:C	2.72	0.62
1:A:5:LEU:HD11	1:A:47:THR:HG21	1.83	0.61
1:B:51:HIS:CD2	1:B:51:HIS:O	2.56	0.59
1:B:2:LYS:HE3	1:B:6:GLU:OE2	2.04	0.58
1:A:110:ASP:O	1:A:114:THR:HG23	2.08	0.53
1:A:51:HIS:CD2	1:A:53:ARG:CA	2.94	0.50
1:B:5:LEU:HD11	1:B:47:THR:HG21	1.96	0.48
1:A:86:GLU:H	1:A:86:GLU:CD	2.16	0.47
1:B:1:CXM:HG3	1:B:227:LEU:HD21	1.97	0.47

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:147:HIS:HB2	1:A:163:LEU:HD11	1.97	0.46
1:B:61:TRP:HH2	1:B:90:LEU:HD12	1.80	0.46
1:B:115:VAL:HA	1:B:118:GLN:HE21	1.81	0.46
1:A:108:HIS:HE1	4:A:778:HOH:O	1.98	0.46
1:A:51:HIS:NE2	1:A:53:ARG:CA	2.70	0.44
1:A:108:HIS:HD2	4:A:779:HOH:O	2.00	0.44
1:B:51:HIS:CG	1:B:51:HIS:O	2.70	0.44
1:B:147:HIS:HB2	1:B:163:LEU:HD11	1.98	0.44
1:B:61:TRP:CD1	1:B:66:ASP:HB3	2.53	0.43
1:A:51:HIS:HE1	1:A:53:ARG:CZ	2.26	0.43
1:B:133:TRP:CZ2	1:B:138:LEU:HD21	2.53	0.43
1:B:26:THR:HG22	1:B:209:TYR:CD1	2.53	0.43
1:B:1:CXM:ON1	1:B:45:VAL:HG13	2.18	0.43
1:B:7:LEU:O	1:B:11:VAL:HG23	2.18	0.43
1:B:109:ILE:HA	4:B:789:HOH:O	2.18	0.42
1:A:105:ASP:OD1	1:A:107:ARG:HD3	2.19	0.42
1:B:231:ILE:HG23	1:B:248:GLU:HB3	2.01	0.42
1:B:96:LYS:HE2	1:B:102:PRO:HD3	2.02	0.41
3:B:300:CB3:H13	3:B:300:CB3:CP2	2.50	0.41
1:B:181:TYR:O	1:B:185:VAL:HG23	2.21	0.41
1:A:1:CXM:HG3	1:A:227:LEU:HD21	2.03	0.41
1:A:7:LEU:HD13	1:A:32:HIS:CD2	2.56	0.41
1:B:92:PRO:HD2	1:B:140:LYS:O	2.20	0.41
1:B:77:VAL:HG12	1:B:79:ILE:HG12	2.02	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:903:HOH:O	4:B:906:HOH:O[2_655]	0.27	1.93

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	262/264 (99%)	255 (97%)	6 (2%)	1 (0%)	34	30
1	B	262/264 (99%)	255 (97%)	6 (2%)	1 (0%)	34	30
All	All	524/528 (99%)	510 (97%)	12 (2%)	2 (0%)	34	30

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	93	VAL
1	B	93	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	232/232 (100%)	225 (97%)	7 (3%)	41	41
1	B	232/232 (100%)	223 (96%)	9 (4%)	32	30
All	All	464/464 (100%)	448 (97%)	16 (3%)	37	36

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	LEU
1	A	38	LEU
1	A	39	GLN
1	A	51	HIS
1	A	116	LEU
1	A	176	PHE
1	A	239	ILE
1	B	17	GLN
1	B	39	GLN
1	B	51	HIS
1	B	53	ARG
1	B	96	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	138	LEU
1	B	166	ARG
1	B	218	LEU
1	B	231	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	GLN
1	A	51	HIS
1	A	108	HIS
1	A	118	GLN
1	A	151	GLN
1	A	217	HIS
1	B	17	GLN
1	B	51	HIS
1	B	118	GLN
1	B	151	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CXM	B	1	1	6,10,11	0.38	0	5,11,13	0.73	0
1	CXM	A	1	1	6,10,11	0.52	0	5,11,13	0.84	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CXM	B	1	1	-	1/7/10/12	-
1	CXM	A	1	1	-	1/7/10/12	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	1	CXM	O-C-CA-CB
1	A	1	CXM	O-C-CA-CB

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	1	CXM	2	0
1	A	1	CXM	1	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	CB3	B	300	-	30,37,37	1.56	4 (13%)	38,51,51	2.75	10 (26%)
3	CB3	A	300	-	30,37,37	1.42	4 (13%)	38,51,51	2.40	9 (23%)
2	UMP	A	301	1	18,21,21	3.65	5 (27%)	21,31,31	1.61	5 (23%)
2	UMP	B	301	1	18,21,21	3.67	6 (33%)	21,31,31	2.09	6 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	CB3	B	300	-	-	0/21/28/28	0/3/3/3
3	CB3	A	300	-	-	6/21/28/28	0/3/3/3
2	UMP	A	301	1	-	1/7/22/22	0/2/2/2
2	UMP	B	301	1	-	1/7/22/22	0/2/2/2

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	UMP	C6-N1	11.98	1.50	1.35
2	B	301	UMP	C6-N1	11.91	1.50	1.35
2	B	301	UMP	C6-C5	7.27	1.54	1.38
2	A	301	UMP	C6-C5	7.25	1.54	1.38
3	B	300	CB3	O4-C4	5.41	1.38	1.24
3	A	300	CB3	O4-C4	4.85	1.36	1.24
2	B	301	UMP	O4'-C1'	3.95	1.51	1.42
2	A	301	UMP	C4-N3	3.68	1.39	1.33
2	A	301	UMP	O4'-C1'	3.67	1.50	1.42
2	B	301	UMP	C4-N3	3.66	1.39	1.33
3	A	300	CB3	C9-N10	2.63	1.50	1.46
3	B	300	CB3	C8-C7	2.48	1.41	1.36
3	A	300	CB3	C5-C6	2.36	1.42	1.37
2	A	301	UMP	P-OP3	-2.28	1.46	1.54
3	A	300	CB3	C8-C7	2.25	1.41	1.36
3	B	300	CB3	C11-C	-2.24	1.45	1.50
2	B	301	UMP	P-OP2	-2.19	1.46	1.54
2	B	301	UMP	P-OP3	-2.01	1.47	1.54
3	B	300	CB3	C5-C6	2.01	1.42	1.37

All (30) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	300	CB3	C4A-C8A-N1	-8.67	118.89	123.60
3	B	300	CB3	CP1-N10-C9	-7.93	109.55	117.10
3	B	300	CB3	C4A-C8A-N1	-7.46	119.55	123.60
3	B	300	CB3	C4A-C4-N3	-6.34	119.98	124.40
3	B	300	CB3	N1-C2-N3	-6.21	118.94	127.22
3	A	300	CB3	N1-C2-N3	-5.92	119.33	127.22
2	B	301	UMP	C5-C6-N1	-5.89	107.53	120.68
3	A	300	CB3	C4A-C4-N3	-5.69	120.44	124.40
3	B	300	CB3	CA-N-C	4.52	128.17	122.34
3	B	300	CB3	C9-N10-C14	3.93	127.53	120.78
2	B	301	UMP	P-O5'-C5'	3.86	128.92	118.30
3	A	300	CB3	C4-N3-C2	3.64	121.72	115.93
3	B	300	CB3	C4-N3-C2	3.63	121.69	115.93
2	A	301	UMP	C5-C6-N1	-3.54	112.78	120.68
2	B	301	UMP	O4'-C4'-C3'	-3.10	98.44	105.67
2	A	301	UMP	P-O5'-C5'	2.98	126.51	118.30
2	A	301	UMP	O4'-C4'-C3'	-2.60	99.59	105.67
2	B	301	UMP	O4'-C1'-C2'	-2.60	101.35	106.25
3	A	300	CB3	CB-CA-N	-2.58	106.44	110.19
2	B	301	UMP	C5-C4-N3	-2.54	117.72	123.31
3	B	300	CB3	C2-N1-C8A	2.43	122.34	116.33
3	A	300	CB3	C2-N1-C8A	2.41	122.30	116.33
2	A	301	UMP	O4'-C1'-C2'	-2.30	101.91	106.25
3	A	300	CB3	C8-C8A-N1	2.27	122.15	118.69
2	B	301	UMP	C4'-O4'-C1'	2.22	114.82	109.45
3	B	300	CB3	C9-C6-C7	-2.19	116.65	120.77
3	A	300	CB3	C13-C14-N10	-2.13	118.44	121.38
2	A	301	UMP	C6-N1-C2	2.06	124.47	121.20
3	B	300	CB3	CP1-N10-C14	2.02	122.86	119.01
3	A	300	CB3	O-C-N	-2.02	118.73	122.45

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	300	CB3	CT-CA-N-C
3	A	300	CB3	N-CA-CB-CG
3	A	300	CB3	CT-CA-CB-CG
3	A	300	CB3	C11-C-N-CA
3	A	300	CB3	O-C-N-CA
3	A	300	CB3	CB-CA-N-C
2	B	301	UMP	O4'-C4'-C5'-O5'
2	A	301	UMP	O4'-C4'-C5'-O5'


There are no ring outliers.

1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	300	CB3	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

### 6.4 Ligands

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers

EDS was not executed - this section is therefore empty.