

# wwPDB X-ray Structure Validation Summary Report (i)

#### Aug 28, 2023 – 10:26 AM EDT

PDB ID : 3KGD

Title : Crystal structure of E. coli RNA 3' cyclase

Authors: Shuman, S.; Tanaka, N.; Smith, P.

Deposited on : 2009-10-28

Resolution : 1.68 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{aligned}  ext{Similar resolution} \ (\# ext{Entries, resolution range}( ext{Å})) \end{aligned}$		
$R_{free}$	130704	6780 (1.70-1.66)		
Clashscore	141614	7310 (1.70-1.66)		
Ramachandran outliers	138981	7173 (1.70-1.66)		
Sidechain outliers	138945	7172 (1.70-1.66)		
RSRZ outliers	127900	6661 (1.70-1.66)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	358	79%	14%	6%
1	В	358	85%	119	% •
1	С	358	80%	15%	6%
1	D	358	80%	14%	• 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard



residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	С	341	-	-	X	-
3	GOL	D	341	-	-	X	-



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 12204 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called RNA 3'-terminal phosphate cyclase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	336	Total	С	N	О	S	0	21	0
1	A	330	2615	1659	465	483	8	0	21	0
1	В	342	Total	С	N	О	S	0	21	0
1	Ъ	342	2658	1681	480	489	8		21	U
1	С	338	Total	С	N	О	S	0	16	0
1			2605	1649	463	484	9	0	10	
1	D	340	Total	С	N	О	S	0	16	0
1		340	2620	1653	472	486	9		10	

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP P46849
A	-17	HIS	-	expression tag	UNP P46849
A	-16	HIS	-	expression tag	UNP P46849
A	-15	HIS	-	expression tag	UNP P46849
A	-14	HIS	-	expression tag	UNP P46849
A	-13	HIS	-	expression tag	UNP P46849
A	-12	HIS	-	expression tag	UNP P46849
A	-11	HIS	-	expression tag	UNP P46849
A	-10	HIS	-	expression tag	UNP P46849
A	-9	HIS	-	expression tag	UNP P46849
A	-8	HIS	-	expression tag	UNP P46849
A	-7	SER	-	expression tag	UNP P46849
A	-6	SER	-	expression tag	UNP P46849
A	-5	GLY	-	expression tag	UNP P46849
A	-4	HIS	-	expression tag	UNP P46849
A	-3	ILE	-	expression tag	UNP P46849
A	-2	GLU		expression tag	UNP P46849
A	-1	GLY	-	expression tag	UNP P46849
A	0	ARG	-	expression tag	UNP P46849
A	1	HIS	-	expression tag	UNP P46849
В	-19	GLY	_	expression tag	UNP P46849



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled  Modelled	Actual	Comment	Reference
В	-18	HIS	-	expression tag	UNP P46849
В	-17	HIS	-	expression tag	UNP P46849
В	-16	HIS	-	expression tag	UNP P46849
В	-15	HIS	-	expression tag	UNP P46849
В	-14	HIS	-	expression tag	UNP P46849
В	-13	HIS	_	expression tag	UNP P46849
В	-12	HIS	-	expression tag	UNP P46849
В	-11	HIS	-	expression tag	UNP P46849
В	-10	HIS	_	expression tag	UNP P46849
В	-9	HIS	-	expression tag	UNP P46849
В	-7	SER	_	expression tag	UNP P46849
В	-6	SER	-	expression tag	UNP P46849
В	-5	GLY	_	expression tag	UNP P46849
В	-4	HIS	_	expression tag	UNP P46849
В	-3	ILE	_	expression tag	UNP P46849
В	-2	GLU	_	expression tag	UNP P46849
В	-1	GLY	-	expression tag	UNP P46849
В	0	ARG	-	expression tag	UNP P46849
В	1	HIS	_	expression tag	UNP P46849
С	-18	GLY	-	expression tag	UNP P46849
С	-17	HIS	-	expression tag	UNP P46849
С	-16	HIS	-	expression tag	UNP P46849
С	-15	HIS	-	expression tag	UNP P46849
С	-14	HIS	-	expression tag	UNP P46849
С	-13	HIS	-	expression tag	UNP P46849
С	-12	HIS	-	expression tag	UNP P46849
С	-11	HIS	-	expression tag	UNP P46849
С	-10	HIS	-	expression tag	UNP P46849
С	-9	HIS	-	expression tag	UNP P46849
С	-8	HIS	-	expression tag	UNP P46849
С	-7	SER	-	expression tag	UNP P46849
С	-6	SER	-	expression tag	UNP P46849
С	-5	GLY	-	expression tag	UNP P46849
С	-4	HIS	-	expression tag	UNP P46849
С	-3	ILE	-	expression tag	UNP P46849
С	-2	GLU	-	expression tag	UNP P46849
С	-1	GLY	_	expression tag	UNP P46849
С	0	ARG	-	expression tag	UNP P46849
С	1	HIS	-	expression tag	UNP P46849
D	-18	GLY	-	expression tag	UNP P46849
D	-17	HIS	-	expression tag	UNP P46849
D	-16	HIS	-	expression tag	UNP P46849



 $Continued\ from\ previous\ page...$ 

Chain	Residue	Modelled	Actual	Comment	Reference
D	-15	HIS	-	expression tag	UNP P46849
D	-14	HIS	-	expression tag	UNP P46849
D	-13	HIS	-	expression tag	UNP P46849
D	-12	HIS	-	expression tag	UNP P46849
D	-11	HIS	-	expression tag	UNP P46849
D	-10	HIS	-	expression tag	UNP P46849
D	-9	HIS	-	expression tag	UNP P46849
D	-8	HIS	-	expression tag	UNP P46849
D	-7	SER	-	expression tag	UNP P46849
D	-6	SER	-	expression tag	UNP P46849
D	-5	GLY	-	expression tag	UNP P46849
D	-4	HIS	-	expression tag	UNP P46849
D	-3	ILE	-	expression tag	UNP P46849
D	-2	GLU	-	expression tag	UNP P46849
D	-1	GLY	-	expression tag	UNP P46849
D	0	ARG	-	expression tag	UNP P46849
D	1	HIS	-	expression tag	UNP P46849

 $\bullet \ \ Molecule\ 2\ is\ ADENOSINE\ MONOPHOSPHATE\ (three-letter\ code:\ AMP)\ (formula:\ C_{10}H_{14}N_5O_7P).$ 


Mol	Chain	Residues	${f Atoms}$				ZeroOcc	AltConf		
2	Λ	A 1	Total	С	N	О	Р	0	0	
	$Z \mid A$		22	10	5	6	1	U		
2	B	1	Total	С	N	О	Р	0	0	
	2 D	1	22	10	5	6	1			
9	С	1	Total	С	N	О	Р	0	0	
	C		22	10	5	6	1	U		



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
9	D	1	Total	С	N	О	Р	0	0
2	D	1	22	10	5	6	1	U	0

• Molecule 3 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 3 3	0	0
3	A	1	Total C O 6 3 3	0	0
3	В	1	Total C O 6 3 3	0	0
3	В	1	Total C O 6 3 3	0	0
3	С	1	Total C O 6 3 3	0	0
3	С	1	Total C O 6 3 3	0	0
3	С	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0
3	D	1	Total C O 6 3 3	0	0



• Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Na 1 1	0	0
4	В	1	Total Na 1 1	0	0
4	С	2	Total Na 2 2	0	0
4	D	1	Total Na 1 1	0	0

 $\bullet$  Molecule 5 is SULFATE ION (three-letter code: SO4) (formula:  $\mathrm{O_4S}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	A	1	Total O S 5 4 1	0	0
5	В	1	Total O S 5 4 1	0	0
5	В	1	Total O S 5 4 1	0	0
5	С	1	Total O S 5 4 1	0	0
5	С	1	Total O S 5 4 1	0	0



 $Continued\ from\ previous\ page...$ 

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
5	D	1	Total 5	O 4	S 1	0	0

#### • Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	375	Total O 376 376	0	1
6	В	411	Total O 413 413	0	2
6	С	363	Total O 364 364	0	1
6	D	359	Total O 360 360	0	1




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: RNA 3'-terminal phosphate cyclase







# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	83.73Å 81.93Å 105.18Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 103.44° 90.00°	Depositor
Resolution (Å)	35.14 - 1.68	Depositor
rtesolution (A)	48.71 - 1.60	EDS
% Data completeness	98.9 (35.14-1.68)	Depositor
(in resolution range)	98.4 (48.71-1.60)	EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.63 (at 1.60Å)	Xtriage
Refinement program	PHENIX 1.4_125	Depositor
P. P.	0.168 , 0.207	Depositor
$R, R_{free}$	0.164 , 0.203	DCC
$R_{free}$ test set	9058 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	22.8	Xtriage
Anisotropy	0.237	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.34, 51.7	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	12204	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 23.39 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.7110e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $<L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, AMP, GOL, SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	Bond lengths		ond angles
IVIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.58	0/2709	0.72	0/3681
1	В	0.57	0/2752	0.72	$2/3736 \ (0.1\%)$
1	С	0.56	0/2680	0.70	0/3636
1	D	0.56	0/2694	0.71	$1/3656 \ (0.0\%)$
All	All	0.57	0/10835	0.71	3/14709 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	186	ARG	NE-CZ-NH2	-6.67	116.97	120.30
1	В	319	ARG	NE-CZ-NH1	-5.55	117.52	120.30
1	D	198	LEU	CA-CB-CG	5.03	126.88	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2615	0	2717	41	0
1	В	2658	0	2725	26	0
1	С	2605	0	2683	55	0
1	D	2620	0	2683	61	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
2	A	22	0	12	0	0
2	В	22	0	12	0	0
2	С	22	0	12	0	0
2	D	22	0	12	1	0
3	A	12	0	16	1	0
3	В	12	0	16	0	0
3	С	18	0	24	9	0
3	D	18	0	24	7	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	2	0	0	0	0
4	D	1	0	0	0	0
5	A	15	0	0	0	0
5	В	10	0	0	0	0
5	С	10	0	0	0	0
5	D	5	0	0	0	0
6	A	376	0	0	10	0
6	В	413	0	0	9	0
6	С	364	0	0	11	0
6	D	360	0	0	12	0
All	All	12204	0	10936	177	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 177 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} \operatorname{Clash} \ \operatorname{overlap}\ (\mathring{\mathbf{A}}) \end{aligned}$
1:A:85:ARG:HH22	1:C:339[B]:GLU:HG2	0.98	1.15
1:A:85:ARG:NH2	1:C:339[B]:GLU:HG2	1.78	0.98
1:D:11[A]:ALA:HA	6:D:766:HOH:O	1.66	0.92
1:D:197:LEU:HD11	1:D:227:LEU:HD23	1.52	0.92
3:C:342:GOL:H11	6:C:1171:HOH:O	1.77	0.85

There are no symmetry-related clashes.



## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percent	iles
1	A	355/358~(99%)	351 (99%)	4 (1%)	0	100 1	.00
1	В	359/358 (100%)	353 (98%)	6 (2%)	0	100 1	.00
1	С	351/358 (98%)	343 (98%)	8 (2%)	0	100 1	.00
1	D	354/358 (99%)	347 (98%)	7 (2%)	0	100 1	.00
All	All	1419/1432 (99%)	1394 (98%)	25 (2%)	0	100 1	.00

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	283/283 (100%)	280 (99%)	3 (1%)	73	61
1	В	283/283 (100%)	282 (100%)	1 (0%)	91	86
1	С	278/283 (98%)	274 (99%)	4 (1%)	67	51
1	D	278/283 (98%)	276 (99%)	2 (1%)	84	76
All	All	1122/1132 (99%)	1112 (99%)	10 (1%)	78	69

5 of 10 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	С	339[B]	GLU
1	D	116	PRO
1	D	126	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	В	231	GLN
1	С	154	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	18	GLN
1	D	104	GLN
1	D	204	HIS
1	С	18	GLN
1	A	18	GLN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry (i)

Of 27 ligands modelled in this entry, 5 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trunc	Chain	Res	Link	Bond lengths			В	ond ang	les
	Type				Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	D	342	-	5,5,5	0.37	0	5,5,5	0.24	0
3	GOL	A	341	-	5,5,5	0.32	0	5,5,5	0.36	0



Mol	Trme	Chain	Res	Link	Во	ond leng	ths	В	ond ang	eles
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	GOL	A	340	-	5,5,5	0.41	0	5,5,5	0.47	0
3	GOL	В	340	-	5,5,5	0.41	0	5,5,5	0.18	0
2	AMP	A	501	1	18,24,25	0.92	1 (5%)	18,35,38	1.60	3 (16%)
2	AMP	С	501	1	18,24,25	0.98	2 (11%)	18,35,38	1.55	5 (27%)
3	GOL	С	342	-	5,5,5	0.29	0	5,5,5	0.77	0
5	SO4	В	343	4	4,4,4	0.26	0	6,6,6	0.56	0
5	SO4	С	346	_	4,4,4	0.14	0	6,6,6	0.43	0
3	GOL	D	340	-	5,5,5	0.39	0	5,5,5	0.43	0
5	SO4	D	344	4	4,4,4	0.26	0	6,6,6	0.27	0
2	AMP	D	501	1	18,24,25	0.92	1 (5%)	18,35,38	1.51	3 (16%)
3	GOL	С	340	-	5,5,5	0.36	0	5,5,5	0.32	0
3	GOL	В	341	-	5,5,5	0.36	0	5,5,5	0.26	0
3	GOL	С	341	-	5,5,5	0.35	0	5,5,5	0.29	0
5	SO4	С	345	-	4,4,4	0.16	0	6,6,6	0.33	0
5	SO4	A	343	-	4,4,4	0.16	0	6,6,6	0.22	0
5	SO4	В	344	-	4,4,4	0.14	0	6,6,6	0.14	0
5	SO4	A	344	_	4,4,4	0.15	0	6,6,6	0.33	0
5	SO4	A	345	-	4,4,4	0.19	0	6,6,6	0.15	0
3	GOL	D	341	_	5,5,5	0.44	0	5,5,5	0.73	0
2	AMP	В	501	1	18,24,25	1.07	1 (5%)	18,35,38	1.68	3 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	D	342	-	-	2/4/4/4	-
3	GOL	A	341	-	-	2/4/4/4	-
2	AMP	D	501	1	-	0/3/25/26	0/3/3/3
3	GOL	С	342	-	-	4/4/4/4	-
3	GOL	A	340	-	-	0/4/4/4	-
3	GOL	С	340	-	-	2/4/4/4	-
3	GOL	В	340	-	-	2/4/4/4	-
3	GOL	В	341	-	-	2/4/4/4	-
3	GOL	С	341	-	-	2/4/4/4	-
3	GOL	D	340	-	-	2/4/4/4	-
2	AMP	A	501	1	-	0/3/25/26	0/3/3/3
2	AMP	С	501	1	-	0/3/25/26	0/3/3/3
3	GOL	D	341	-	-	4/4/4/4	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	В	501	1	-	0/3/25/26	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	Ideal(Å)
2	A	501	AMP	C5-C4	2.26	1.46	1.40
2	D	501	AMP	C5-C4	2.19	1.46	1.40
2	С	501	AMP	C5-C4	2.15	1.46	1.40
2	С	501	AMP	C2-N3	2.14	1.35	1.32
2	В	501	AMP	C5-C4	2.14	1.46	1.40

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$Ideal(^{o})$
2	В	501	AMP	N3-C2-N1	-4.35	121.88	128.68
2	A	501	AMP	N3-C2-N1	-3.88	122.61	128.68
2	D	501	AMP	N3-C2-N1	-3.70	122.89	128.68
2	В	501	AMP	C2-N1-C6	3.12	124.09	118.75
2	С	501	AMP	N6-C6-N1	2.92	124.63	118.57

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	341	GOL	O1-C1-C2-C3
3	В	341	GOL	O1-C1-C2-O2
3	В	341	GOL	O1-C1-C2-C3
3	С	340	GOL	O1-C1-C2-C3
3	С	341	GOL	O1-C1-C2-C3


There are no ring outliers.

6 monomers are involved in 18 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	340	GOL	1	0
3	С	342	GOL	1	0
3	D	340	GOL	3	0
2	D	501	AMP	1	0
3	С	341	GOL	8	0
3	D	341	GOL	4	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



## 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	A	336/358~(93%)	-0.44	1 (0%) 94 94	16, 25, 45, 84	0
1	В	342/358 (95%)	-0.37	7 (2%) 65 69	16, 26, 49, 124	0
1	С	338/358 (94%)	-0.31	8 (2%) 59 62	15, 28, 53, 111	0
1	D	340/358 (94%)	-0.29	6 (1%) 68 72	15, 27, 75, 113	0
All	All	1356/1432 (94%)	-0.35	22 (1%) 72 75	15, 26, 55, 124	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	230	ASP	9.9
1	С	2	MET	5.9
1	В	-10	HIS	3.9
1	С	161	TYR	3.5
1	A	161	TYR	3.2

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

## 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	GOL	В	340	6/6	0.58	0.24	62,66,68,68	0
3	GOL	С	340	6/6	0.68	0.20	73,74,76,77	0
3	GOL	D	342	6/6	0.78	0.24	55,63,65,68	0
3	GOL	A	341	6/6	0.81	0.14	72,74,74,75	0
3	GOL	В	341	6/6	0.81	0.13	80,83,84,85	0
3	GOL	С	342	6/6	0.82	0.21	34,38,43,46	0
3	GOL	D	340	6/6	0.84	0.15	58,66,68,69	0
3	GOL	С	341	6/6	0.84	0.13	49,51,55,59	0
5	SO4	В	344	5/5	0.88	0.10	45,47,49,50	5
3	GOL	D	341	6/6	0.89	0.19	43,46,50,53	0
5	SO4	A	344	5/5	0.93	0.14	26,31,33,38	5
5	SO4	A	345	5/5	0.94	0.17	36,38,38,41	5
5	SO4	С	346	5/5	0.94	0.14	30,32,34,35	5
3	GOL	A	340	6/6	0.95	0.20	24,32,34,38	0
4	NA	С	343	1/1	0.95	0.07	35,35,35,35	0
4	NA	D	343	1/1	0.96	0.12	32,32,32,32	0
5	SO4	В	343	5/5	0.96	0.10	63,67,70,83	0
4	NA	В	342	1/1	0.97	0.10	32,32,32,32	0
2	AMP	D	501	22/23	0.97	0.07	19,22,28,34	0
5	SO4	D	344	5/5	0.97	0.09	52,53,72,74	0
4	NA	A	342	1/1	0.98	0.07	43,43,43,43	0
2	AMP	С	501	22/23	0.98	0.07	19,21,25,30	0
2	AMP	A	501	22/23	0.98	0.07	17,21,25,27	0
4	NA	С	344	1/1	0.98	0.09	32,32,32,32	0
5	SO4	С	345	5/5	0.98	0.09	42,43,46,103	0
2	AMP	В	501	22/23	0.98	0.05	19,20,25,26	0
5	SO4	A	343	5/5	0.98	0.09	51,53,60,67	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




### Electron density around AMP D 501:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)










#### Electron density around AMP C 501:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 0.7 rmsd) in gray  $\mathrm{mF}_o\text{-}\mathrm{DF}_c$  (at 3 rmsd) in purple (negative) and green (positive)













# 6.5 Other polymers (i)

There are no such residues in this entry.

