



# wwPDB NMR Structure Validation Summary Report ⓘ

Apr 20, 2024 – 10:00 PM EDT

PDB ID : 2KWD  
Title : Supramolecular Protein Structure Determination by Site-Specific Long-Range Intermolecular Solid State NMR Spectroscopy  
Authors : Nieuwkoop, A.J.; Rienstra, C.M.  
Deposited on : 2010-04-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
wwPDB-RCI : v\_1n\_11\_5\_13\_A (Berjanski et al., 2005)  
PANAV : Wang et al. (2010)  
wwPDB-ShiftChecker : v1.2  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*SOLID-STATE NMR*

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	56	36% 55% 7% .
1	B	56	46% 48% . .
1	C	56	39% 54% 7%
1	D	56	45% 50% 5%
1	E	56	43% 48% 9%

## 2 Ensemble composition and analysis i

This entry contains 10 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:56, B:1-B:56, C:1-C:56, D:1-D:56, E:1-E:56 (280)	0.53	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	2, 3, 4, 5, 7, 8
2	1, 6, 9, 10

### 3 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 4290 atoms, of which 2100 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Immunoglobulin G-binding protein G.

Mol	Chain	Residues	Atoms					Trace	
1	A	56	Total	C	H	N	O	S	0
			858	274	420	68	95	1	
1	B	56	Total	C	H	N	O	S	0
			858	274	420	68	95	1	
1	C	56	Total	C	H	N	O	S	0
			858	274	420	68	95	1	
1	D	56	Total	C	H	N	O	S	0
			858	274	420	68	95	1	
1	E	56	Total	C	H	N	O	S	0
			858	274	420	68	95	1	

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP P19909
A	2	GLN	-	expression tag	UNP P19909
B	1	MET	-	expression tag	UNP P19909
B	2	GLN	-	expression tag	UNP P19909
C	1	MET	-	expression tag	UNP P19909
C	2	GLN	-	expression tag	UNP P19909
D	1	MET	-	expression tag	UNP P19909
D	2	GLN	-	expression tag	UNP P19909
E	1	MET	-	expression tag	UNP P19909
E	2	GLN	-	expression tag	UNP P19909

## 4 Residue-property plots [i](#)


### 4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Immunoglobulin G-binding protein G




- Molecule 1: Immunoglobulin G-binding protein G




- Molecule 1: Immunoglobulin G-binding protein G



- Molecule 1: Immunoglobulin G-binding protein G



- Molecule 1: Immunoglobulin G-binding protein G




## 4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: Immunoglobulin G-binding protein G

Chain A: 





- Molecule 1: Immunoglobulin G-binding protein G

Chain B: 



- Molecule 1: Immunoglobulin G-binding protein G

Chain C: 




- Molecule 1: Immunoglobulin G-binding protein G

Chain D: 



- Molecule 1: Immunoglobulin G-binding protein G

Chain E: 



## 5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 600 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR NIH	structure solution	

No chemical shift data was provided. Note: This is a solid-state NMR structure, where hydrogen atoms are typically not assigned a chemical shift value, which may lead to lower completeness of assignment measure.

## 6 Model quality [i](#)

### 6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.89±0.00	0±0/444 ( 0.0± 0.0%)	0.93±0.00	1±0/600 ( 0.2± 0.0%)
1	B	0.86±0.01	0±0/444 ( 0.0± 0.0%)	0.86±0.01	1±0/600 ( 0.1± 0.1%)
1	C	0.86±0.00	0±0/444 ( 0.0± 0.0%)	0.88±0.01	0±0/600 ( 0.0± 0.0%)
1	D	0.84±0.00	0±0/444 ( 0.0± 0.0%)	0.81±0.01	0±0/600 ( 0.0± 0.0%)
1	E	0.84±0.00	0±0/444 ( 0.0± 0.0%)	0.82±0.00	0±0/600 ( 0.0± 0.0%)
All	All	0.86	0/22200 ( 0.0%)	0.86	18/30000 ( 0.1%)

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	11	THR	CA-CB-CG2	-5.27	105.03	112.40	3	10
1	B	11	THR	CA-CB-CG2	-5.26	105.03	112.40	3	8

There are no chirality outliers.

There are no planarity outliers.

### 6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	438	420	420	61±5
1	B	438	420	420	36±3
1	C	438	420	420	35±3
1	D	438	420	420	33±3

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	E	438	420	420	34±2
All	All	21900	21000	21000	1682

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 39.

5 of 347 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:12:LEU:N	1:B:13:LYS:HZ3	0.83	1.72	1	10
1:A:12:LEU:N	1:A:13:LYS:HZ3	0.82	1.72	10	10
1:D:9:GLY:C	1:D:13:LYS:HZ1	0.80	1.79	7	10
1:E:9:GLY:C	1:E:13:LYS:HZ1	0.80	1.79	5	10
1:C:9:GLY:C	1:C:13:LYS:HZ1	0.79	1.81	10	10

## 6.3 Torsion angles [i](#)

### 6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	54/56 (96%)	54±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	B	54/56 (96%)	54±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	C	54/56 (96%)	54±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	D	54/56 (96%)	54±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
1	E	54/56 (96%)	54±0 (100±0%)	0±0 (0±0%)	0±0 (0±0%)	100	100
All	All	2700/2800 (96%)	2700 (100%)	0 (0%)	0 (0%)	100	100

There are no Ramachandran outliers.

### 6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	46/46 (100%)	36±1 (78±3%)	10±1 (22±3%)	3	30
1	B	46/46 (100%)	37±2 (81±3%)	9±2 (19±3%)	4	36
1	C	46/46 (100%)	37±2 (81±3%)	9±2 (19±3%)	4	36
1	D	46/46 (100%)	38±1 (82±3%)	8±1 (18±3%)	4	39
1	E	46/46 (100%)	37±1 (80±3%)	9±1 (20±3%)	3	33
All	All	2300/2300 (100%)	1849 (80%)	451 (20%)	4	34

5 of 66 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	1	MET	10
1	A	11	THR	10
1	A	13	LYS	10
1	A	16	THR	10
1	A	21	VAL	10

### 6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 6.7 Other polymers [i](#)

There are no such molecules in this entry.

## 6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 7 Chemical shift validation

No chemical shift data were provided